



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 12:43 PM UTC

PDB ID : 5YS2 / pdb_00005ys2
Title : Structure of the domain IV(D_IV) of Pseudorabies virus glycoprotein B(PRV gB)
Authors : Hu, X.L.; Yang, F.L.
Deposited on : 2017-11-12
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	254	
1	B	254	
1	C	254	
1	D	254	
1	E	254	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	254	% 46%16%6%32%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 8134 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein B,Envelope glycoprotein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	175	Total	C	N	O	S	0	0	0
			1365	849	246	258	12			
1	B	169	Total	C	N	O	S	0	0	0
			1324	825	236	251	12			
1	C	177	Total	C	N	O	S	0	0	0
			1380	858	248	262	12			
1	D	174	Total	C	N	O	S	0	0	0
			1361	848	244	257	12			
1	E	172	Total	C	N	O	S	0	0	0
			1347	838	241	256	12			
1	F	173	Total	C	N	O	S	0	0	0
			1357	846	243	256	12			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	484	GLY	-	linker	UNP A0A0U3FH21
A	485	GLY	-	linker	UNP A0A0U3FH21
A	486	SER	-	linker	UNP A0A0U3FH21
A	487	GLY	-	linker	UNP A0A0U3FH21
A	643	HIS	-	expression tag	UNP A0A1Q0AKY1
A	644	HIS	-	expression tag	UNP A0A1Q0AKY1
A	645	HIS	-	expression tag	UNP A0A1Q0AKY1
A	646	HIS	-	expression tag	UNP A0A1Q0AKY1
A	647	HIS	-	expression tag	UNP A0A1Q0AKY1
A	648	HIS	-	expression tag	UNP A0A1Q0AKY1
A	649	HIS	-	expression tag	UNP A0A1Q0AKY1
A	650	HIS	-	expression tag	UNP A0A1Q0AKY1
B	484	GLY	-	linker	UNP A0A0U3FH21
B	485	GLY	-	linker	UNP A0A0U3FH21
B	486	SER	-	linker	UNP A0A0U3FH21
B	487	GLY	-	linker	UNP A0A0U3FH21
B	643	HIS	-	expression tag	UNP A0A1Q0AKY1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	644	HIS	-	expression tag	UNP A0A1Q0AKY1
B	645	HIS	-	expression tag	UNP A0A1Q0AKY1
B	646	HIS	-	expression tag	UNP A0A1Q0AKY1
B	647	HIS	-	expression tag	UNP A0A1Q0AKY1
B	648	HIS	-	expression tag	UNP A0A1Q0AKY1
B	649	HIS	-	expression tag	UNP A0A1Q0AKY1
B	650	HIS	-	expression tag	UNP A0A1Q0AKY1
C	484	GLY	-	linker	UNP A0A0U3FH21
C	485	GLY	-	linker	UNP A0A0U3FH21
C	486	SER	-	linker	UNP A0A0U3FH21
C	487	GLY	-	linker	UNP A0A0U3FH21
C	643	HIS	-	expression tag	UNP A0A1Q0AKY1
C	644	HIS	-	expression tag	UNP A0A1Q0AKY1
C	645	HIS	-	expression tag	UNP A0A1Q0AKY1
C	646	HIS	-	expression tag	UNP A0A1Q0AKY1
C	647	HIS	-	expression tag	UNP A0A1Q0AKY1
C	648	HIS	-	expression tag	UNP A0A1Q0AKY1
C	649	HIS	-	expression tag	UNP A0A1Q0AKY1
C	650	HIS	-	expression tag	UNP A0A1Q0AKY1
D	484	GLY	-	linker	UNP A0A0U3FH21
D	485	GLY	-	linker	UNP A0A0U3FH21
D	486	SER	-	linker	UNP A0A0U3FH21
D	487	GLY	-	linker	UNP A0A0U3FH21
D	643	HIS	-	expression tag	UNP A0A1Q0AKY1
D	644	HIS	-	expression tag	UNP A0A1Q0AKY1
D	645	HIS	-	expression tag	UNP A0A1Q0AKY1
D	646	HIS	-	expression tag	UNP A0A1Q0AKY1
D	647	HIS	-	expression tag	UNP A0A1Q0AKY1
D	648	HIS	-	expression tag	UNP A0A1Q0AKY1
D	649	HIS	-	expression tag	UNP A0A1Q0AKY1
D	650	HIS	-	expression tag	UNP A0A1Q0AKY1
E	484	GLY	-	linker	UNP A0A0U3FH21
E	485	GLY	-	linker	UNP A0A0U3FH21
E	486	SER	-	linker	UNP A0A0U3FH21
E	487	GLY	-	linker	UNP A0A0U3FH21
E	643	HIS	-	expression tag	UNP A0A1Q0AKY1
E	644	HIS	-	expression tag	UNP A0A1Q0AKY1
E	645	HIS	-	expression tag	UNP A0A1Q0AKY1
E	646	HIS	-	expression tag	UNP A0A1Q0AKY1
E	647	HIS	-	expression tag	UNP A0A1Q0AKY1
E	648	HIS	-	expression tag	UNP A0A1Q0AKY1
E	649	HIS	-	expression tag	UNP A0A1Q0AKY1

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	650	HIS	-	expression tag	UNP A0A1Q0AKY1
F	484	GLY	-	linker	UNP A0A0U3FH21
F	485	GLY	-	linker	UNP A0A0U3FH21
F	486	SER	-	linker	UNP A0A0U3FH21
F	487	GLY	-	linker	UNP A0A0U3FH21
F	643	HIS	-	expression tag	UNP A0A1Q0AKY1
F	644	HIS	-	expression tag	UNP A0A1Q0AKY1
F	645	HIS	-	expression tag	UNP A0A1Q0AKY1
F	646	HIS	-	expression tag	UNP A0A1Q0AKY1
F	647	HIS	-	expression tag	UNP A0A1Q0AKY1
F	648	HIS	-	expression tag	UNP A0A1Q0AKY1
F	649	HIS	-	expression tag	UNP A0A1Q0AKY1
F	650	HIS	-	expression tag	UNP A0A1Q0AKY1


- Molecule 1: Envelope glycoprotein B, Envelope glycoprotein B




• Molecule 1: Envelope glycoprotein B,Envelope glycoprotein B



• Molecule 1: Envelope glycoprotein B,Envelope glycoprotein B



• Molecule 1: Envelope glycoprotein B,Envelope glycoprotein B



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	91.91Å 119.85Å 123.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.52 – 2.70 46.52 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.6 (46.52-2.70) 99.6 (46.52-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.227 , 0.291 0.227 , 0.288	Depositor DCC
R_{free} test set	1914 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	63.0	Xtriage
Anisotropy	0.497	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 35.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for -h,l,k	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8134	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/1390	0.99	1/1883 (0.1%)
1	B	0.57	0/1348	0.94	1/1825 (0.1%)
1	C	0.60	0/1406	1.05	6/1906 (0.3%)
1	D	0.72	0/1386	1.02	5/1877 (0.3%)
1	E	0.56	0/1371	0.93	1/1857 (0.1%)
1	F	0.72	0/1382	1.08	8/1872 (0.4%)
All	All	0.63	0/8283	1.00	22/11220 (0.2%)

There are no bond length outliers.

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	69	CYS	CA-C-N	-8.29	113.69	119.66
1	C	69	CYS	C-N-CA	-8.29	113.69	119.66
1	C	81	GLU	CA-C-N	7.89	128.95	120.11
1	C	81	GLU	C-N-CA	7.89	128.95	120.11
1	F	562	ARG	N-CA-C	-6.68	101.68	110.43

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1365	0	1337	44	0
1	B	1324	0	1295	35	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1380	0	1348	43	0
1	D	1361	0	1331	37	0
1	E	1347	0	1316	45	0
1	F	1357	0	1328	45	0
All	All	8134	0	7955	199	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 199 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:633:ILE:HD12	1:C:78:VAL:HG21	1.21	1.11
1:B:633:ILE:CD1	1:C:78:VAL:HG21	1.81	1.09
1:A:575:HIS:CG	1:A:576:ASN:H	1.76	1.03
1:D:605:HIS:CE1	1:D:621:TYR:CE2	2.49	1.00
1:D:605:HIS:ND1	1:D:621:TYR:CZ	2.31	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	171/254 (67%)	159 (93%)	12 (7%)	0	100	100
1	B	163/254 (64%)	153 (94%)	10 (6%)	0	100	100
1	C	173/254 (68%)	159 (92%)	14 (8%)	0	100	100
1	D	168/254 (66%)	156 (93%)	12 (7%)	0	100	100
1	E	166/254 (65%)	155 (93%)	11 (7%)	0	100	100
1	F	167/254 (66%)	154 (92%)	13 (8%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1008/1524 (66%)	936 (93%)	72 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	149/205 (73%)	127 (85%)	22 (15%)	3	8
1	B	145/205 (71%)	127 (88%)	18 (12%)	4	11
1	C	151/205 (74%)	126 (83%)	25 (17%)	2	6
1	D	149/205 (73%)	125 (84%)	24 (16%)	2	6
1	E	148/205 (72%)	123 (83%)	25 (17%)	2	6
1	F	149/205 (73%)	128 (86%)	21 (14%)	3	9
All	All	891/1230 (72%)	756 (85%)	135 (15%)	3	7

5 of 135 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	77	VAL
1	F	522	THR
1	F	610	LYS
1	C	522	THR
1	C	515	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	554	ASN
1	C	527	GLN
1	E	605	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	175/254 (68%)	-0.05	1 (0%) 85 85	19, 51, 107, 140	0
1	B	169/254 (66%)	-0.12	4 (2%) 59 56	17, 44, 91, 130	0
1	C	177/254 (69%)	-0.28	1 (0%) 85 85	9, 36, 98, 138	0
1	D	174/254 (68%)	-0.28	3 (1%) 69 67	15, 37, 90, 115	0
1	E	172/254 (67%)	-0.22	2 (1%) 76 75	15, 39, 86, 129	0
1	F	173/254 (68%)	-0.28	2 (1%) 76 75	10, 33, 94, 122	0
All	All	1040/1524 (68%)	-0.20	13 (1%) 75 73	9, 41, 98, 140	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	574	GLU	5.2
1	D	579	GLY	3.1
1	D	575	HIS	2.8
1	A	575	HIS	2.8
1	E	598	ILE	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.