



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:55 PM UTC

PDB ID : 5YSD / pdb_00005ysd
Title : Crystal structure of beta-1,2-glucooligosaccharide binding protein in complex with sophorotriose
Authors : Abe, K.; Nakajima, M.; Taguchi, H.; Arakawa, T.; Fushinobu, S.
Deposited on : 2017-11-14
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	7% 90% 7%
1	B	397	28% 91% 7%
2	C	3	100%
2	D	3	67% 33%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6538 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lin1841 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	1	0
			3065	1958	497	593	17			
1	B	389	Total	C	N	O	S	0	0	0
			3078	1965	502	595	16			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	26	MET	-	expression tag	UNP Q92AS8
A	415	LEU	-	expression tag	UNP Q92AS8
A	416	GLU	-	expression tag	UNP Q92AS8
A	417	HIS	-	expression tag	UNP Q92AS8
A	418	HIS	-	expression tag	UNP Q92AS8
A	419	HIS	-	expression tag	UNP Q92AS8
A	420	HIS	-	expression tag	UNP Q92AS8
A	421	HIS	-	expression tag	UNP Q92AS8
A	422	HIS	-	expression tag	UNP Q92AS8
B	26	MET	-	expression tag	UNP Q92AS8
B	415	LEU	-	expression tag	UNP Q92AS8
B	416	GLU	-	expression tag	UNP Q92AS8
B	417	HIS	-	expression tag	UNP Q92AS8
B	418	HIS	-	expression tag	UNP Q92AS8
B	419	HIS	-	expression tag	UNP Q92AS8
B	420	HIS	-	expression tag	UNP Q92AS8
B	421	HIS	-	expression tag	UNP Q92AS8
B	422	HIS	-	expression tag	UNP Q92AS8

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-2)-beta-D-glucopyranose-(1-2)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	3	Total	C	O	0	0	0
			34	18	16			
2	D	3	Total	C	O	0	0	0
			34	18	16			

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		
3	A	1	Total	C	O	0	0
			8	6	2		
3	A	1	Total	C	O	0	0
			8	6	2		
3	B	1	Total	C	O	0	0
			8	6	2		
3	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Mg 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	176	Total 176	O 176	0	0
5	B	110	Total 110	O 110	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lin1841 protein



- Molecule 1: Lin1841 protein



- Molecule 2: beta-D-glucopyranose-(1-2)-beta-D-glucopyranose-(1-2)-alpha-D-glucopyranose



- Molecule 2: beta-D-glucopyranose-(1-2)-beta-D-glucopyranose-(1-2)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	36.08Å 125.56Å 91.78Å 90.00° 101.61° 90.00°	Depositor
Resolution (Å)	44.95 – 2.10 44.95 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.4 (44.95-2.10) 96.4 (44.95-2.10)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0158, Coot	Depositor
R, R_{free}	0.169 , 0.207 0.203 , 0.230	Depositor DCC
R_{free} test set	2264 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	28.4	Xtriage
Anisotropy	0.166	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.033 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6538	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BGC, GLC, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.26	5/3142 (0.2%)	1.12	4/4253 (0.1%)
1	B	1.18	2/3157 (0.1%)	1.08	2/4273 (0.0%)
All	All	1.22	7/6299 (0.1%)	1.10	6/8526 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	292	LYS	CA-C	6.36	1.55	1.52
1	A	49	LEU	C-O	-6.15	1.17	1.24
1	A	36	VAL	C-O	-5.73	1.18	1.24
1	A	365	LYS	C-O	-5.45	1.17	1.24
1	B	314	ASP	C-O	-5.41	1.17	1.24
1	A	305	SER	N-CA	5.41	1.52	1.45
1	B	145	TYR	C-O	5.03	1.29	1.23

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	223	LYS	CA-C-N	-6.24	112.13	119.05
1	A	223	LYS	C-N-CA	-6.24	112.13	119.05
1	B	223	LYS	CA-C-N	-5.70	112.82	119.32
1	B	223	LYS	C-N-CA	-5.70	112.82	119.32
1	A	119	ASN	N-CA-C	5.31	120.41	113.30
1	A	205	ARG	CB-CG-CD	5.09	123.01	111.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3065	0	2985	11	0
1	B	3078	0	2991	8	0
2	C	34	0	30	0	0
2	D	34	0	30	0	0
3	A	24	0	42	6	0
3	B	16	0	28	0	0
4	A	1	0	0	0	0
5	A	176	0	0	1	0
5	B	110	0	0	1	0
All	All	6538	0	6106	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:503:MPD:H12	3:A:503:MPD:H52	1.67	0.74
1:B:50:LYS:HD2	1:B:66:VAL:HB	1.88	0.55
1:A:246:LEU:HG	3:A:503:MPD:H13	1.91	0.52
1:A:163:TYR:HA	3:A:501:MPD:HM1	1.93	0.50
1:A:319:PHE:CE1	1:A:323:MET:HE3	2.47	0.50
1:B:183:ARG:CZ	1:B:187:MET:HE2	2.42	0.50
1:B:73:ASN:HB3	5:B:697:HOH:O	2.12	0.49
1:A:42[B]:MET:CE	1:A:71:TRP:CH2	2.97	0.48
1:A:42[B]:MET:HE3	1:A:71:TRP:CZ3	2.49	0.47
1:A:113:ASP:HB3	1:A:317:LEU:HD21	1.97	0.47
3:A:503:MPD:H12	3:A:503:MPD:C5	2.43	0.47
1:B:113:ASP:HB3	1:B:317:LEU:HD21	1.96	0.46
1:B:319:PHE:CE1	1:B:323:MET:HE3	2.52	0.44
3:A:501:MPD:HM2	5:A:653:HOH:O	2.18	0.44
1:A:183:ARG:CZ	1:A:187:MET:HE2	2.47	0.43
1:A:361:GLY:HA2	1:A:364:MET:HE2	2.00	0.43
1:B:322:TYR:CD1	1:B:322:TYR:C	2.97	0.42
1:B:111:THR:O	1:B:115:GLU:HG2	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:42:MET:HE3	1:B:71:TRP:CZ3	2.56	0.41
1:A:160:LYS:HE2	1:A:160:LYS:HA	2.03	0.40
1:A:165:GLU:H	3:A:501:MPD:HM3	1.86	0.40
1:A:42[B]:MET:HE2	1:A:71:TRP:CH2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/397 (97%)	378 (98%)	8 (2%)	0	100	100
1	B	387/397 (98%)	380 (98%)	7 (2%)	0	100	100
All	All	773/794 (97%)	758 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/341 (98%)	326 (98%)	7 (2%)	47	54
1	B	334/341 (98%)	322 (96%)	12 (4%)	31	34
All	All	667/682 (98%)	648 (97%)	19 (3%)	37	43

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	ASP
1	A	55	LYS
1	A	112	LYS
1	A	118	LYS
1	A	160	LYS
1	A	168	LYS
1	A	351	LEU
1	B	35	LYS
1	B	50	LYS
1	B	51	GLU
1	B	112	LYS
1	B	121	ASN
1	B	168	LYS
1	B	246	LEU
1	B	308	LYS
1	B	311	ASN
1	B	330	LEU
1	B	351	LEU
1	B	411	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	325	GLN
1	B	67	GLN
1	B	325	GLN
1	B	381	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	C	1	2	12,12,12	0.58	0	17,17,17	1.85	6 (35%)
2	BGC	C	2	2	11,11,12	0.60	0	15,15,17	1.27	2 (13%)
2	BGC	C	3	2	11,11,12	0.83	0	15,15,17	1.17	2 (13%)
2	GLC	D	1	2	12,12,12	0.47	0	17,17,17	1.33	2 (11%)
2	BGC	D	2	2	11,11,12	0.53	0	15,15,17	0.71	0
2	BGC	D	3	2	11,11,12	0.64	0	15,15,17	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	BGC	C	2	2	-	0/2/19/22	0/1/1/1
2	BGC	C	3	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	BGC	D	2	2	-	0/2/19/22	0/1/1/1
2	BGC	D	3	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GLC	O5-C1-C2	4.23	117.73	110.30
2	C	2	BGC	C1-O5-C5	3.06	116.29	112.19
2	D	1	GLC	C1-O5-C5	2.92	119.30	113.65
2	C	1	GLC	O3-C3-C2	-2.51	104.45	110.38
2	C	1	GLC	O2-C2-C3	-2.44	104.62	110.38
2	C	1	GLC	C1-O5-C5	2.28	118.07	113.65
2	C	3	BGC	O4-C4-C3	-2.14	105.34	110.38
2	C	2	BGC	O4-C4-C3	-2.13	105.35	110.38
2	C	1	GLC	O1-C1-O5	-2.12	104.12	110.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	BGC	C1-C2-C3	2.04	112.61	109.64
2	C	1	GLC	O2-C2-C1	2.03	113.94	109.25
2	D	1	GLC	O2-C2-C3	-2.00	105.66	110.38


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MPD	A	502	-	7,7,7	0.36	0	9,10,10	0.51	0
3	MPD	B	501	-	7,7,7	0.56	0	9,10,10	1.19	0
3	MPD	A	503	-	7,7,7	0.91	0	9,10,10	1.20	0
3	MPD	B	502	-	7,7,7	0.35	0	9,10,10	1.11	1 (11%)
3	MPD	A	501	-	7,7,7	1.05	0	9,10,10	1.67	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. ' ' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	A	502	-	-	2/5/5/5	-
3	MPD	B	501	-	-	1/5/5/5	-
3	MPD	A	503	-	-	1/5/5/5	-
3	MPD	B	502	-	-	2/5/5/5	-
3	MPD	A	501	-	-	4/5/5/5	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	MPD	CM-C2-C1	-3.18	103.51	110.63
3	A	501	MPD	O4-C4-C5	-2.62	98.17	109.45
3	B	502	MPD	CM-C2-C1	-2.46	105.11	110.63

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	MPD	O2-C2-C3-C4
3	A	501	MPD	CM-C2-C3-C4
3	A	501	MPD	C2-C3-C4-C5
3	B	502	MPD	C1-C2-C3-C4
3	B	502	MPD	CM-C2-C3-C4
3	A	503	MPD	C2-C3-C4-C5
3	B	501	MPD	C2-C3-C4-C5
3	A	501	MPD	C1-C2-C3-C4
3	A	502	MPD	C1-C2-C3-C4
3	A	502	MPD	CM-C2-C3-C4

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	503	MPD	3	0
3	A	501	MPD	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	387/397 (97%)	0.44	27 (6%)	22 24	22, 45, 87, 116	1 (0%)
1	B	389/397 (97%)	1.36	112 (28%)	1 1	33, 66, 104, 135	0
All	All	776/794 (97%)	0.90	139 (17%)	3 4	22, 52, 100, 135	1 (0%)

All (139) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	322	TYR	6.1
1	A	410	ALA	5.0
1	B	319	PHE	4.9
1	A	408	VAL	4.9
1	B	170	TRP	4.7
1	A	412	LEU	4.7
1	A	411	LEU	4.6
1	B	328	VAL	4.6
1	B	83	ALA	4.6
1	B	320	MET	4.6
1	B	79	LEU	4.4
1	B	330	LEU	4.3
1	B	90	VAL	4.3
1	B	280	ILE	4.2
1	B	36	VAL	4.2
1	B	326	PRO	4.2
1	A	415	LEU	4.2
1	B	82	VAL	4.1
1	B	166	ALA	4.0
1	B	321	ASP	4.0
1	B	278	PRO	3.8
1	B	35	LYS	3.8
1	B	78	LEU	3.7
1	B	161	VAL	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	115	GLU	3.5
1	B	81	ALA	3.4
1	B	329	GLN	3.4
1	B	91	VAL	3.4
1	A	351	LEU	3.4
1	A	414	LYS	3.4
1	B	169	THR	3.3
1	B	137	GLY	3.3
1	B	53	ALA	3.3
1	B	246	LEU	3.3
1	B	171	ASP	3.3
1	B	324	SER	3.3
1	B	159	LYS	3.3
1	B	359	VAL	3.2
1	B	167	PRO	3.2
1	A	418	HIS	3.2
1	A	420	HIS	3.2
1	B	318	LYS	3.2
1	B	323	MET	3.1
1	B	356	TYR	3.1
1	A	407	GLN	3.1
1	B	89	ASP	3.1
1	B	347	GLU	3.1
1	A	34	SER	3.1
1	B	325	GLN	3.0
1	A	406	ASP	3.0
1	B	168	LYS	3.0
1	B	309	TYR	3.0
1	B	282	GLY	3.0
1	B	279	ASP	3.0
1	B	139	THR	2.9
1	B	259	ILE	2.9
1	B	327	ASP	2.9
1	B	50	LYS	2.9
1	A	35	LYS	2.9
1	B	163	TYR	2.8
1	B	275	ASP	2.8
1	B	118	LYS	2.8
1	B	334	LYS	2.8
1	A	413	LYS	2.7
1	B	162	GLY	2.7
1	B	353	ASN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	170	TRP	2.7
1	B	360	PHE	2.7
1	B	273	LEU	2.7
1	B	54	GLN	2.6
1	B	55	LYS	2.6
1	B	357	TYR	2.6
1	A	216	GLY	2.6
1	B	247	GLY	2.6
1	B	307	PHE	2.6
1	B	361	GLY	2.6
1	B	244	THR	2.6
1	B	122	SER	2.6
1	B	341	ALA	2.5
1	B	331	SER	2.5
1	B	52	LEU	2.5
1	A	350	MET	2.5
1	A	86	SER	2.5
1	B	85	LYS	2.5
1	B	311	ASN	2.5
1	B	84	SER	2.4
1	B	88	PRO	2.4
1	B	355	PRO	2.4
1	B	422	HIS	2.4
1	B	76	ASP	2.4
1	B	349	ASP	2.4
1	B	114	VAL	2.4
1	B	270	VAL	2.4
1	B	420	HIS	2.4
1	B	56	PHE	2.3
1	B	116	LYS	2.3
1	A	82	VAL	2.3
1	B	47	LYS	2.3
1	A	309	TYR	2.3
1	B	256	GLY	2.3
1	B	283	LYS	2.3
1	B	80	THR	2.3
1	A	85	LYS	2.3
1	B	124	LEU	2.3
1	A	409	GLU	2.2
1	B	34	SER	2.2
1	B	126	PHE	2.2
1	B	156	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	279	ASP	2.2
1	B	61	GLY	2.2
1	B	277	ALA	2.2
1	B	131	LYS	2.2
1	A	311	ASN	2.2
1	A	215	ASP	2.2
1	B	93	MET	2.2
1	B	249	ASP	2.2
1	A	417	HIS	2.2
1	B	108	LEU	2.2
1	B	164	ASN	2.2
1	B	187	MET	2.2
1	B	281	ASP	2.2
1	B	250	ALA	2.1
1	B	37	LEU	2.1
1	B	276	THR	2.1
1	B	92	GLN	2.1
1	B	46	ALA	2.1
1	A	217	THR	2.1
1	B	332	TRP	2.1
1	B	352	LYS	2.1
1	B	86	SER	2.1
1	B	248	LEU	2.1
1	B	350	MET	2.1
1	B	165	GLU	2.0
1	B	77	LYS	2.0
1	B	284	TRP	2.0
1	B	285	ALA	2.0
1	B	158	LEU	2.0
1	B	160	LYS	2.0
1	B	340	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	C	1	12/12	0.92	0.09	19,22,23,28	0
2	BGC	D	2	11/12	0.92	0.09	25,28,31,33	0
2	GLC	D	1	12/12	0.93	0.08	26,28,32,39	0
2	BGC	C	2	11/12	0.95	0.08	18,20,21,21	0
2	BGC	C	3	11/12	0.95	0.07	17,18,19,21	0
2	BGC	D	3	11/12	0.95	0.08	22,25,26,27	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MPD	A	501	8/8	0.77	0.14	29,34,44,50	0
3	MPD	B	501	8/8	0.82	0.16	51,55,63,67	0
3	MPD	B	502	8/8	0.84	0.16	43,47,48,49	0
3	MPD	A	502	8/8	0.86	0.14	43,51,59,63	0
3	MPD	A	503	8/8	0.87	0.14	36,45,50,50	0
4	MG	A	504	1/1	0.94	0.17	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.