



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2026 – 02:17 AM UTC

PDB ID : 5YSE / pdb_00005yse
Title : Crystal structure of beta-1,2-glucooligosaccharide binding protein in complex with sophorotetraose
Authors : Abe, K.; Nakajima, M.; Taguchi, H.; Arakawa, T.; Fushinobu, S.
Deposited on : 2017-11-14
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	7% 91% 6% .
1	B	397	45% 93% 5% .
2	C	4	75% 25%
2	D	4	75% 25%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6899 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lin1841 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	3	0
			3081	1968	502	595	16			
1	B	387	Total	C	N	O	S	0	1	0
			3071	1961	502	592	16			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	26	MET	-	expression tag	UNP Q92AS8
A	415	LEU	-	expression tag	UNP Q92AS8
A	416	GLU	-	expression tag	UNP Q92AS8
A	417	HIS	-	expression tag	UNP Q92AS8
A	418	HIS	-	expression tag	UNP Q92AS8
A	419	HIS	-	expression tag	UNP Q92AS8
A	420	HIS	-	expression tag	UNP Q92AS8
A	421	HIS	-	expression tag	UNP Q92AS8
A	422	HIS	-	expression tag	UNP Q92AS8
B	26	MET	-	expression tag	UNP Q92AS8
B	415	LEU	-	expression tag	UNP Q92AS8
B	416	GLU	-	expression tag	UNP Q92AS8
B	417	HIS	-	expression tag	UNP Q92AS8
B	418	HIS	-	expression tag	UNP Q92AS8
B	419	HIS	-	expression tag	UNP Q92AS8
B	420	HIS	-	expression tag	UNP Q92AS8
B	421	HIS	-	expression tag	UNP Q92AS8
B	422	HIS	-	expression tag	UNP Q92AS8

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-2)-beta-D-glucopyranose-(1-2)-beta-D-glucopyranose-(1-2)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	4	Total	C	O	0	0	0
			45	24	21			
2	D	4	Total	C	O	0	0	0
			45	24	21			

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		
3	A	1	Total	C	O	0	0
			8	6	2		
3	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	427	Total 427	O 427	0	0
5	B	205	Total 205	O 205	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Lin1841 protein



• Molecule 1: Lin1841 protein



• Molecule 2: beta-D-glucopyranose-(1-2)-beta-D-glucopyranose-(1-2)-alpha-D-glucopyranose





- Molecule 2: beta-D-glucopyranose-(1-2)-beta-D-glucopyranose-(1-2)-beta-D-glucopyranose-(1-2)-alpha-D-glucopyranose

Chain D:  75% 25%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	36.01Å 63.28Å 87.45Å 89.99° 82.24° 86.08°	Depositor
Resolution (Å)	43.32 – 1.60 43.32 – 1.60	Depositor EDS
% Data completeness (in resolution range)	96.4 (43.32-1.60) 96.4 (43.32-1.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.158 , 0.188 0.204 , 0.227	Depositor DCC
R_{free} test set	4783 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.251	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6899	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, MPD, MG, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.35	9/3165 (0.3%)	1.09	2/4284 (0.0%)
1	B	1.08	3/3153 (0.1%)	1.01	1/4268 (0.0%)
All	All	1.22	12/6318 (0.2%)	1.05	3/8552 (0.0%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	331	SER	CB-OG	8.62	1.59	1.42
1	A	142	VAL	CA-CB	-8.28	1.48	1.54
1	A	292	LYS	CA-C	8.00	1.56	1.52
1	A	69	ILE	CA-CB	-7.94	1.48	1.54
1	B	227	ASP	CG-OD2	7.70	1.40	1.25
1	A	279	ASP	CB-CG	6.96	1.69	1.52
1	B	227	ASP	CG-OD1	-6.90	1.12	1.25
1	A	362	GLU	C-O	-6.52	1.16	1.24
1	B	421	HIS	N-CA	-6.26	1.38	1.46
1	A	69	ILE	CA-C	5.67	1.56	1.52
1	A	235	PHE	N-CA	5.34	1.52	1.46
1	A	409	GLU	C-O	-5.26	1.18	1.24

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	331	SER	CA-CB-OG	6.94	124.98	111.10
1	A	279	ASP	CB-CG-OD1	5.48	131.00	118.40
1	B	227	ASP	CB-CG-OD2	-5.13	106.60	118.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3081	0	3002	8	0
1	B	3071	0	2986	7	0
2	C	45	0	39	1	0
2	D	45	0	39	1	0
3	A	24	0	42	2	0
4	A	1	0	0	0	0
5	A	427	0	0	2	1
5	B	205	0	0	0	0
All	All	6899	0	6108	16	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:503:MPD:HM1	3:A:503:MPD:H52	1.92	0.52
1:A:35:LYS:N	5:A:605:HOH:O	2.43	0.51
1:B:69:ILE:HD13	1:B:77:LYS:HD3	1.94	0.50
1:A:263:PHE:CE2	1:A:269:MET:HG2	2.48	0.48
1:A:40:TRP:CD1	1:A:88:PRO:HB3	2.49	0.48
5:A:733:HOH:O	1:B:422:HIS:HE1	1.98	0.47
1:B:263:PHE:CE2	1:B:269:MET:HG2	2.50	0.45
1:A:268:TRP:CD2	2:C:3:BGC:H5	2.51	0.45
1:B:40:TRP:CD1	1:B:88:PRO:HB3	2.52	0.45
1:A:374:PRO:HG2	1:A:415:LEU:HD21	1.99	0.44
1:A:294:ASN:OD1	1:A:296[B]:GLU:HG3	2.18	0.43
3:A:503:MPD:HM1	3:A:503:MPD:C5	2.49	0.42
1:A:281:ASP:OD2	1:B:309:TYR:OH	2.22	0.41
1:A:263:PHE:HE2	1:A:269:MET:HG2	1.85	0.41
1:B:289:LEU:O	1:B:295:ASN:HB2	2.20	0.41
1:B:268:TRP:CD2	2:D:3:BGC:H5	2.56	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:931:HOH:O	5:A:938:HOH:O[1_565]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/397 (98%)	380 (98%)	8 (2%)	0	100	100
1	B	386/397 (97%)	378 (98%)	8 (2%)	0	100	100
All	All	774/794 (98%)	758 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/341 (98%)	330 (98%)	5 (2%)	57	35
1	B	333/341 (98%)	327 (98%)	6 (2%)	51	28
All	All	668/682 (98%)	657 (98%)	11 (2%)	53	33

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	MET
1	A	58	LYS
1	A	214	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	291	LYS
1	A	413	LYS
1	B	42	MET
1	B	352	LYS
1	B	358	LYS
1	B	364	MET
1	B	413	LYS
1	B	417	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	271	ASN
1	A	400	GLN
1	B	400	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	C	1	2	12,12,12	0.71	0	17,17,17	1.12	1 (5%)
2	BGC	C	2	2	11,11,12	0.80	0	15,15,17	1.09	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	C	3	2	11,11,12	0.79	0	15,15,17	0.88	1 (6%)
2	BGC	C	4	2	11,11,12	1.07	0	15,15,17	0.99	1 (6%)
2	GLC	D	1	2	12,12,12	0.83	0	17,17,17	1.03	1 (5%)
2	BGC	D	2	2	11,11,12	1.03	0	15,15,17	1.15	1 (6%)
2	BGC	D	3	2	11,11,12	0.95	0	15,15,17	0.71	1 (6%)
2	BGC	D	4	2	11,11,12	1.06	1 (9%)	15,15,17	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	BGC	C	2	2	-	0/2/19/22	0/1/1/1
2	BGC	C	3	2	-	0/2/19/22	0/1/1/1
2	BGC	C	4	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	BGC	D	2	2	-	0/2/19/22	0/1/1/1
2	BGC	D	3	2	-	0/2/19/22	0/1/1/1
2	BGC	D	4	2	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	4	BGC	O5-C1	-2.20	1.40	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	GLC	C3-C4-C5	-2.55	105.61	110.23
2	C	1	GLC	O6-C6-C5	-2.53	102.71	111.33
2	D	2	BGC	C6-C5-C4	-2.35	107.25	113.02
2	C	3	BGC	C1-O5-C5	2.19	115.12	112.19
2	C	4	BGC	C1-O5-C5	2.16	115.08	112.19
2	C	2	BGC	C1-C2-C3	-2.08	106.62	109.64
2	D	3	BGC	O5-C5-C4	2.03	115.77	110.83

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3	BGC	1	0
2	D	3	BGC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MPD	A	501	-	7,7,7	0.56	0	9,10,10	0.67	0
3	MPD	A	502	-	7,7,7	0.49	0	9,10,10	0.60	0
3	MPD	A	503	-	7,7,7	0.47	0	9,10,10	1.18	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	A	501	-	-	3/5/5/5	-
3	MPD	A	502	-	-	2/5/5/5	-
3	MPD	A	503	-	-	1/5/5/5	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	503	MPD	CM-C2-C1	-2.11	105.89	110.63

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	MPD	C2-C3-C4-C5
3	A	503	MPD	C2-C3-C4-C5
3	A	501	MPD	CM-C2-C3-C4
3	A	501	MPD	O2-C2-C3-C4
3	A	501	MPD	C1-C2-C3-C4
3	A	502	MPD	C1-C2-C3-C4

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	503	MPD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	387/397 (97%)	0.60	27 (6%)	22 23	15, 33, 60, 87	3 (0%)
1	B	387/397 (97%)	2.05	178 (45%)	0 0	25, 56, 102, 144	1 (0%)
All	All	774/794 (97%)	1.32	205 (26%)	1 1	15, 43, 94, 144	4 (0%)

All (205) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	397	VAL	6.5
1	B	248	LEU	6.4
1	B	351	LEU	6.4
1	B	258	GLY	6.0
1	B	403	THR	5.7
1	B	243	ASP	5.6
1	B	235	PHE	5.6
1	B	391	TYR	5.6
1	B	226	VAL	5.5
1	B	395	ALA	5.5
1	B	390	ILE	5.5
1	B	387	TRP	5.5
1	B	399	THR	5.4
1	B	244	THR	5.3
1	B	259	ILE	5.3
1	B	239	GLY	5.1
1	B	161	VAL	5.1
1	B	246	LEU	5.1
1	B	410	ALA	5.0
1	A	117	SER	4.9
1	B	236	ILE	4.9
1	B	396	ASP	4.9
1	B	260	VAL	4.8
1	B	188	TYR	4.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	405	ASN	4.7
1	B	230	THR	4.7
1	B	254	PHE	4.7
1	B	402	ASP	4.6
1	B	247	GLY	4.5
1	B	227	ASP	4.4
1	B	253	SER	4.3
1	B	404	PHE	4.3
1	B	400	GLN	4.3
1	B	394	GLY	4.2
1	B	241	ALA	4.2
1	B	250	ALA	4.2
1	B	184	GLY	4.2
1	B	256	GLY	4.1
1	B	349	ASP	4.0
1	B	242	PRO	3.9
1	B	121	ASN	3.9
1	B	401	MET	3.9
1	B	384	GLY	3.8
1	B	255	GLY	3.8
1	B	234	SER	3.8
1	B	251	SER	3.8
1	B	189	GLY	3.8
1	B	412	LEU	3.8
1	B	380	ALA	3.8
1	B	382	LEU	3.7
1	B	350	MET	3.7
1	B	57	THR	3.7
1	B	240	SER	3.6
1	B	252	GLN	3.5
1	B	261	PRO	3.5
1	B	233	ASP	3.4
1	B	408	VAL	3.4
1	B	136	ASP	3.4
1	B	293	GLU	3.4
1	B	238	ASN	3.4
1	B	422	HIS	3.4
1	A	118	LYS	3.4
1	A	83	ALA	3.4
1	B	181	SER	3.4
1	B	417	HIS	3.4
1	B	383	TYR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	393	GLY	3.3
1	B	386	SER	3.3
1	B	182	LYS	3.3
1	B	352	LYS	3.3
1	B	398	GLN	3.3
1	B	231	TYR	3.2
1	B	245	ASP	3.2
1	B	137	GLY	3.2
1	A	136	ASP	3.2
1	B	59	ASP	3.2
1	B	406	ASP	3.2
1	B	180	LEU	3.2
1	B	331	SER	3.2
1	B	114	VAL	3.2
1	B	257	ASP	3.2
1	B	158	LEU	3.2
1	B	174	SER	3.2
1	B	62	ILE	3.1
1	A	417	HIS	3.1
1	B	249	ASP	3.1
1	B	407	GLN	3.1
1	B	170	TRP	3.0
1	B	124	LEU	3.0
1	B	186	ASP	3.0
1	A	137	GLY	3.0
1	B	116	LYS	3.0
1	B	163	TYR	3.0
1	A	415	LEU	3.0
1	B	421	HIS	3.0
1	B	311	ASN	3.0
1	B	36	VAL	2.9
1	B	219	VAL	2.9
1	B	60	THR	2.9
1	B	346	TRP	2.9
1	B	388	GLU	2.9
1	B	178	LEU	2.9
1	B	61	GLY	2.9
1	B	374	PRO	2.9
1	B	237	LYS	2.9
1	B	117	SER	2.9
1	B	353	ASN	2.9
1	A	104	ALA	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	208	GLY	2.8
1	B	223	LYS	2.8
1	B	126	PHE	2.8
1	B	278	PRO	2.8
1	B	112	LYS	2.8
1	B	327	ASP	2.8
1	B	228	THR	2.7
1	B	389	GLN	2.7
1	B	207	ASN	2.7
1	B	364	MET	2.7
1	B	127	PRO	2.7
1	B	330	LEU	2.7
1	B	111	THR	2.7
1	B	294	ASN	2.7
1	B	292	LYS	2.7
1	B	135	PHE	2.7
1	B	347	GLU	2.7
1	A	82	VAL	2.7
1	B	376	PHE	2.6
1	A	80	THR	2.6
1	B	58	LYS	2.6
1	A	216	GLY	2.6
1	B	279	ASP	2.6
1	B	281	ASP	2.6
1	B	217	THR	2.6
1	B	187	MET	2.6
1	B	68	VAL	2.5
1	A	84	SER	2.5
1	A	309	TYR	2.5
1	B	150	VAL	2.5
1	B	229	VAL	2.5
1	B	366	THR	2.5
1	B	381	GLN	2.5
1	B	155	THR	2.5
1	B	392	ARG	2.5
1	B	56	PHE	2.4
1	B	157	LEU	2.4
1	B	159	LYS	2.4
1	A	105	GLY	2.4
1	B	64	VAL	2.4
1	A	109	ASP	2.4
1	B	348	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	118	LYS	2.4
1	B	415	LEU	2.4
1	B	263	PHE	2.4
1	B	416	GLU	2.4
1	B	162	GLY	2.4
1	B	224	PRO	2.4
1	B	169	THR	2.4
1	B	110	ILE	2.4
1	B	214	LYS	2.3
1	B	418	HIS	2.3
1	A	121	ASN	2.3
1	B	220	PHE	2.3
1	A	119	ASN	2.3
1	A	86	SER	2.3
1	B	179	LYS	2.3
1	B	213	ASP	2.3
1	B	185	LYS	2.3
1	A	279	ASP	2.2
1	B	153	TYR	2.2
1	A	420	HIS	2.2
1	B	66	VAL	2.2
1	B	177	ALA	2.2
1	B	420	HIS	2.2
1	B	379	ILE	2.2
1	B	221	ASN	2.2
1	B	328	VAL	2.2
1	A	106	ALA	2.2
1	B	343	MET	2.2
1	B	345	ALA	2.2
1	B	317	LEU	2.2
1	B	385	LYS	2.2
1	B	171	ASP	2.1
1	A	102	VAL	2.1
1	B	183	ARG	2.1
1	B	232	LEU	2.1
1	A	35	LYS	2.1
1	A	116	LYS	2.1
1	B	290	PRO	2.1
1	B	55	LYS	2.1
1	B	411	LEU	2.1
1	B	409	GLU	2.1
1	B	225	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	277	ALA	2.1
1	B	291	LYS	2.1
1	B	54	GLN	2.1
1	A	79	LEU	2.1
1	B	326	PRO	2.1
1	B	378	GLU	2.0
1	A	349	ASP	2.0
1	B	102	VAL	2.0
1	B	215	ASP	2.0
1	B	218	PRO	2.0
1	B	361	GLY	2.0
1	B	130	VAL	2.0
1	A	112	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	C	1	12/12	0.95	0.10	15,17,21,26	0
2	BGC	D	2	11/12	0.95	0.10	20,23,28,28	0
2	BGC	C	3	11/12	0.96	0.09	13,13,14,15	0
2	BGC	C	4	11/12	0.96	0.09	13,14,16,17	0
2	GLC	D	1	12/12	0.96	0.10	26,30,33,37	0
2	BGC	C	2	11/12	0.96	0.10	14,14,16,17	0
2	BGC	D	3	11/12	0.96	0.10	20,22,24,25	0
2	BGC	D	4	11/12	0.97	0.09	19,20,22,23	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MPD	A	503	8/8	0.79	0.16	48,50,51,54	0
3	MPD	A	502	8/8	0.84	0.12	33,37,47,47	0
3	MPD	A	501	8/8	0.89	0.13	27,33,35,38	0
4	MG	A	504	1/1	0.94	0.21	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.