



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:24 PM UTC

PDB ID : 5YSJ / pdb_00005ysj
Title : SdeA mART-C domain WT apo
Authors : Kim, L.; Kwon, D.H.; Song, H.K.
Deposited on : 2017-11-14
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	152	7% 81% 9% • 9%
1	B	152	12% 82% 9% • 8%
1	C	152	9% 82% 9% 9%
1	D	152	4% 88% 5% • 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8673 atoms, of which 4197 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitinating/deubiquitinating enzyme SdeA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	138	Total	C	H	N	O	S	0	0	0
			2133	690	1046	181	213	3			
1	B	140	Total	C	H	N	O	S	0	0	0
			2156	703	1050	183	217	3			
1	C	138	Total	C	H	N	O	S	0	0	0
			2142	695	1051	181	212	3			
1	D	144	Total	C	H	N	O	S	0	0	0
			2183	715	1050	192	223	3			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	754	GLY	-	expression tag	UNP Q5ZTK4
A	755	SER	-	expression tag	UNP Q5ZTK4
B	754	GLY	-	expression tag	UNP Q5ZTK4
B	755	SER	-	expression tag	UNP Q5ZTK4
C	754	GLY	-	expression tag	UNP Q5ZTK4
C	755	SER	-	expression tag	UNP Q5ZTK4
D	754	GLY	-	expression tag	UNP Q5ZTK4
D	755	SER	-	expression tag	UNP Q5ZTK4


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	16	Total	O	0	0
			16	16		
2	B	13	Total	O	0	0
			13	13		
2	C	8	Total	O	0	0
			8	8		
2	D	22	Total	O	0	0
			22	22		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Ubiquitinating/deubiquitinating enzyme SdeA





- Molecule 1: Ubiquitinating/deubiquitinating enzyme SdeA



- Molecule 1: Ubiquitinating/deubiquitinating enzyme SdeA



- Molecule 1: Ubiquitinating/deubiquitinating enzyme SdeA



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.96Å 86.68Å 71.03Å 90.00° 97.90° 90.00°	Depositor
Resolution (Å)	43.54 – 2.06 43.54 – 2.06	Depositor EDS
% Data completeness (in resolution range)	99.0 (43.54-2.06) 99.0 (43.54-2.06)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.40 (at 2.06Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.225 , 0.269 0.227 , 0.270	Depositor DCC
R_{free} test set	2000 reflections (6.11%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.178	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 41.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8673	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.21 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4411e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.15	0/1106	0.37	0/1495
1	B	0.15	0/1126	0.37	0/1522
1	C	0.14	0/1111	0.37	0/1502
1	D	0.13	0/1154	0.48	2/1560 (0.1%)
All	All	0.14	0/4497	0.40	2/6079 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	847	LEU	CA-C-N	7.93	136.69	121.54
1	D	847	LEU	C-N-CA	7.93	136.69	121.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1087	1046	1093	9	0
1	B	1106	1050	1106	6	0
1	C	1091	1051	1095	4	0
1	D	1133	1050	1130	5	0
2	A	16	0	0	0	0
2	B	13	0	0	0	0
2	C	8	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	22	0	0	0	0
All	All	4476	4197	4424	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:785:MET:HE1	1:D:810:LEU:HG	1.70	0.73
1:B:766:ARG:NH2	1:B:811:SER:O	2.28	0.66
1:A:813:MET:HE1	1:A:840:MET:SD	2.36	0.66
1:A:766:ARG:HD3	1:A:813:MET:HE2	1.88	0.56
1:A:813:MET:HE1	1:A:840:MET:CE	2.38	0.54
1:B:766:ARG:NH2	1:B:813:MET:HG2	2.24	0.53
1:D:847:LEU:HB3	1:D:865:VAL:HG13	1.91	0.52
1:C:766:ARG:HD3	1:C:813:MET:HE1	1.91	0.51
1:A:847:LEU:HB3	1:A:865:VAL:HG13	1.93	0.50
1:A:759:LYS:HA	1:A:759:LYS:HE2	1.93	0.49
1:D:799:PRO:O	1:D:803:LYS:HG2	2.13	0.48
1:B:759:LYS:N	1:B:759:LYS:HD3	2.29	0.47
1:D:785:MET:HE1	1:D:810:LEU:CG	2.40	0.47
1:A:860:GLU:H	1:A:860:GLU:HG3	1.41	0.44
1:C:813:MET:HE3	1:C:840:MET:SD	2.59	0.43
1:A:877:LYS:HG2	1:C:879:THR:HG21	2.00	0.43
1:A:877:LYS:CG	1:C:879:THR:HG21	2.49	0.42
1:B:852:VAL:O	1:B:853:GLY:C	2.63	0.42
1:A:852:VAL:O	1:A:853:GLY:C	2.63	0.42
1:B:766:ARG:HG3	1:B:821:THR:HG22	2.02	0.41
1:B:859:THR:O	1:B:859:THR:HG23	2.21	0.40
1:D:759:LYS:HA	1:D:759:LYS:HE2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	134/152 (88%)	129 (96%)	5 (4%)	0	100	100
1	B	136/152 (90%)	131 (96%)	5 (4%)	0	100	100
1	C	134/152 (88%)	129 (96%)	5 (4%)	0	100	100
1	D	142/152 (93%)	137 (96%)	5 (4%)	0	100	100
All	All	546/608 (90%)	526 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/134 (92%)	118 (95%)	6 (5%)	23	16
1	B	126/134 (94%)	116 (92%)	10 (8%)	11	5
1	C	124/134 (92%)	114 (92%)	10 (8%)	11	5
1	D	128/134 (96%)	124 (97%)	4 (3%)	35	30
All	All	502/536 (94%)	472 (94%)	30 (6%)	17	10

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	762	THR
1	A	781	GLN
1	A	804	GLN
1	A	826	LYS
1	A	850	LYS
1	A	860	GLU
1	B	759	LYS
1	B	789	THR
1	B	791	GLU
1	B	827	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	859	THR
1	B	860	GLU
1	B	869	GLU
1	B	877	LYS
1	B	888	GLU
1	B	902	ASP
1	C	776	LYS
1	C	781	GLN
1	C	791	GLU
1	C	807	LEU
1	C	827	LEU
1	C	850	LYS
1	C	852	VAL
1	C	886	LYS
1	C	899	LYS
1	C	902	ASP
1	D	776	LYS
1	D	803	LYS
1	D	806	LYS
1	D	852	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	781	GLN
1	C	788	ASN
1	D	855	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/152 (90%)	0.72	11 (7%) 18 19	22, 36, 58, 80	0
1	B	140/152 (92%)	0.76	18 (12%) 7 7	18, 32, 57, 78	0
1	C	138/152 (90%)	0.77	13 (9%) 14 14	22, 36, 61, 78	0
1	D	144/152 (94%)	0.44	6 (4%) 40 41	18, 31, 54, 82	0
All	All	560/608 (92%)	0.67	48 (8%) 16 17	18, 34, 57, 82	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	853	GLY	6.0
1	B	859	THR	5.3
1	A	853	GLY	4.9
1	D	788	ASN	4.6
1	B	869	GLU	4.4
1	C	869	GLU	3.8
1	A	788	ASN	3.8
1	C	903	PHE	3.8
1	A	803	LYS	3.7
1	C	827	LEU	3.6
1	B	772	GLU	3.4
1	C	902	ASP	3.3
1	C	852	VAL	3.3
1	B	903	PHE	3.1
1	C	886	LYS	3.1
1	C	788	ASN	3.0
1	B	852	VAL	2.8
1	A	759	LYS	2.8
1	D	854	ARG	2.8
1	B	759	LYS	2.8
1	C	807	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	852	VAL	2.7
1	A	860	GLU	2.7
1	A	901	PRO	2.7
1	B	888	GLU	2.6
1	A	787	ALA	2.6
1	D	886	LYS	2.6
1	B	860	GLU	2.6
1	B	861	SER	2.5
1	D	900	SER	2.5
1	D	902	ASP	2.5
1	B	788	ASN	2.5
1	B	795	THR	2.4
1	A	827	LEU	2.4
1	B	885	GLN	2.4
1	B	870	ASP	2.3
1	A	859	THR	2.3
1	C	808	ASN	2.3
1	B	812	LYS	2.3
1	C	787	ALA	2.2
1	C	810	LEU	2.2
1	A	813	MET	2.2
1	C	758	ALA	2.2
1	C	900	SER	2.1
1	B	900	SER	2.1
1	B	818	ASN	2.1
1	B	776	LYS	2.1
1	D	856	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.