



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 09:50 AM UTC

PDB ID : 4YT0 / pdb_00004yt0
Title : Crystal structure of Mitochondrial rhodoquinol-fumarate reductase from Ascaris suum with 2-methyl-N-[3-(1-methylethoxy)phenyl]benzamide.
Authors : Harada, S.; Shiba, T.; Sato, D.; Yamamoto, A.; Nagahama, M.; Yone, A.; Inaoka, D.K.; Sakamoto, K.; Inoue, M.; Honma, T.; Kita, K.
Deposited on : 2015-03-17
Resolution : 3.66 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1062 (3.78-3.54)
Clashscore	190562	1009 (3.76-3.56)
Ramachandran outliers	187476	1054 (3.78-3.54)
Sidechain outliers	187428	1052 (3.78-3.54)
RSRZ outliers	180081	1061 (3.78-3.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	645	 74% 20% . .
1	E	645	 73% 21% . .
2	B	282	 65% 21% . 11%
2	F	282	 62% 23% . 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	188	
3	G	188	
4	D	156	
4	H	156	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MLI	E	701	-	-	X	-
8	SF4	B	302	-	-	X	-
8	SF4	F	302	-	-	X	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 18236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinate dehydrogenase flavoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	616	Total	C	N	O	S	0	0	0
			4787	3004	855	900	28			
1	E	616	Total	C	N	O	S	0	0	0
			4787	3004	855	900	28			

- Molecule 2 is a protein called Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	250	Total	C	N	O	S	0	0	0
			1985	1263	338	361	23			
2	F	250	Total	C	N	O	S	0	0	0
			1985	1263	338	361	23			


- Molecule 3 is a protein called Cytochrome b-large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	153	Total	C	N	O	S	0	0	0
			1217	813	204	194	6			
3	G	150	Total	C	N	O	S	0	0	0
			1195	798	201	190	6			


- Molecule 4 is a protein called Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	129	Total	C	N	O	S	0	0	0
			998	659	165	169	5			
4	H	129	Total	C	N	O	S	0	0	0
			998	659	165	169	5			

- Molecule 5 is MALONATE ION (CCD ID: MLI) (formula: C₃H₂O₄).




- Molecule 7 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	Fe	S	0	0
			4	2	2		
7	F	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 8 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			8	4	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	F	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 9 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe_3S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	Fe	S	0	0
			7	3	4		
9	F	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 10 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $\text{C}_{34}\text{H}_{32}\text{FeN}_4\text{O}_4$).



-
- The chemical structure is 2-(4-isobutoxyphenyl)benzoic acid. It consists of a central biphenyl system. The left ring is a benzoic acid moiety with a carboxylic acid group (-COOH) at the 1-position and a phenyl ring at the 2-position. The right ring is a 4-isobutoxyphenyl moiety with an isobutoxy group (-OCH₂CH(CH₃)₂) at the 4-position and a phenyl ring at the 1-position. The atoms are labeled as follows: C1-C7 for the left ring, C8 for the carbonyl carbon, O1 for the carbonyl oxygen, N for the amide nitrogen, C9-C14 for the right ring, C15-C17 for the isobutoxy group, and O2 for the ether oxygen.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	C	1	Total	C	N	O	0	0
			20	17	1	2		




Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	G	1	Total	C	N	O	0	0
			20	17	1	2		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Succinate dehydrogenase flavoprotein




• Molecule 1: Succinate dehydrogenase flavoprotein







- Molecule 2: Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial




- Molecule 2: Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial




- Molecule 3: Cytochrome b-large subunit




- Molecule 3: Cytochrome b-large subunit



- Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial



- Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	122.80Å 123.39Å 219.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.66 20.00 – 3.66	Depositor EDS
% Data completeness (in resolution range)	89.0 (20.00-3.66) 88.7 (20.00-3.66)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 3.62Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.193 , 0.262 0.196 , 0.260	Depositor DCC
R_{free} test set	1644 reflections (4.41%)	wwPDB-VP
Wilson B-factor (Å ²)	52.9	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 26.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.058 for k,h,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	18236	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, MLI, FAD, HEM, FES, MRN, F3S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/4889	0.79	0/6605
1	E	0.49	0/4889	0.80	3/6605 (0.0%)
2	B	0.51	0/2029	0.80	0/2739
2	F	0.49	0/2029	0.79	0/2739
3	C	0.52	0/1255	0.84	0/1709
3	G	0.50	0/1232	0.80	0/1676
4	D	0.55	0/1030	0.85	0/1406
4	H	0.53	0/1030	0.86	0/1406
All	All	0.50	0/18383	0.81	3/24885 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	383	ILE	CA-C-N	-5.37	114.23	119.76
1	E	383	ILE	C-N-CA	-5.37	114.23	119.76
1	E	425	HIS	N-CA-C	5.33	118.00	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4787	0	4720	79	0
1	E	4787	0	4720	85	0
2	B	1985	0	2002	48	0
2	F	1985	0	2003	46	0
3	C	1217	0	1265	26	0
3	G	1195	0	1240	16	0
4	D	998	0	985	18	0
4	H	998	0	985	15	0
5	A	7	0	2	0	0
5	E	7	0	2	2	0
6	A	53	0	31	7	0
6	E	53	0	31	6	0
7	B	4	0	0	0	0
7	F	4	0	0	0	0
8	B	8	0	0	2	0
8	F	8	0	0	3	0
9	B	7	0	0	0	0
9	F	7	0	0	0	0
10	C	43	0	30	8	0
10	G	43	0	30	4	0
11	C	20	0	19	1	0
11	G	20	0	19	2	0
All	All	18236	0	18084	311	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 311 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:HIS:NE2	6:A:702:FAD:HM83	1.73	1.03
2:B:182:CYS:HG	8:B:302:SF4:FE3	0.68	0.96
2:F:185:CYS:HG	8:F:302:SF4:FE1	0.77	0.88
1:A:79:HIS:NE2	6:A:702:FAD:C8M	2.44	0.79
2:F:185:CYS:SG	8:F:302:SF4:FE1	1.74	0.79

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	614/645 (95%)	568 (92%)	43 (7%)	3 (0%)	24	55
1	E	614/645 (95%)	563 (92%)	51 (8%)	0	100	100
2	B	248/282 (88%)	215 (87%)	28 (11%)	5 (2%)	6	29
2	F	248/282 (88%)	219 (88%)	22 (9%)	7 (3%)	4	23
3	C	151/188 (80%)	139 (92%)	11 (7%)	1 (1%)	18	48
3	G	148/188 (79%)	134 (90%)	11 (7%)	3 (2%)	6	29
4	D	127/156 (81%)	119 (94%)	5 (4%)	3 (2%)	4	25
4	H	127/156 (81%)	114 (90%)	11 (9%)	2 (2%)	7	32
All	All	2277/2542 (90%)	2071 (91%)	182 (8%)	24 (1%)	11	39

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	80	THR
2	B	140	ASN
2	F	140	ASN
2	F	243	MET
4	D	54	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	502/527 (95%)	473 (94%)	29 (6%)	18	43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	502/527 (95%)	471 (94%)	31 (6%)	16	41
2	B	220/242 (91%)	203 (92%)	17 (8%)	12	36
2	F	220/242 (91%)	204 (93%)	16 (7%)	13	38
3	C	127/158 (80%)	117 (92%)	10 (8%)	11	35
3	G	124/158 (78%)	112 (90%)	12 (10%)	8	29
4	D	98/119 (82%)	90 (92%)	8 (8%)	10	34
4	H	98/119 (82%)	93 (95%)	5 (5%)	21	46
All	All	1891/2092 (90%)	1763 (93%)	128 (7%)	14	40

5 of 128 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	98	LEU
3	G	136	ILE
3	C	61	THR
3	C	60	LEU
3	G	168	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	276	HIS
1	E	497	GLN
1	E	347	GLN
1	E	431	ASN
1	E	558	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	MRN	C	202	-	21,21,21	2.06	3 (14%)	28,28,28	1.11	2 (7%)
7	FES	B	301	2	0,4,4	-	-	-		
8	SF4	B	302	2	0,12,12	-	-	-		
9	F3S	B	303	2	0,9,9	-	-	-		
7	FES	F	301	2	0,4,4	-	-	-		
5	MLI	E	701	-	6,6,6	1.13	0	7,7,7	1.03	0
6	FAD	A	702	-	58,58,58	1.51	10 (17%)	85,89,89	1.65	18 (21%)
5	MLI	A	701	-	6,6,6	1.15	0	7,7,7	1.20	0
8	SF4	F	302	2	0,12,12	-	-	-		
10	HEM	G	201	4,3	50,50,50	1.58	8 (16%)	67,82,82	1.93	16 (23%)
11	MRN	G	202	-	21,21,21	2.06	3 (14%)	28,28,28	1.34	3 (10%)
10	HEM	C	201	4,3	50,50,50	1.63	9 (18%)	67,82,82	1.89	19 (28%)
6	FAD	E	702	-	58,58,58	1.51	9 (15%)	85,89,89	1.74	21 (24%)
9	F3S	F	303	2	0,9,9	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	MRN	C	202	-	-	7/12/12/12	0/2/2/2
7	FES	B	301	2	-	-	0/1/1/1
8	SF4	B	302	2	-	-	0/6/5/5
9	F3S	B	303	2	-	-	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	FES	F	301	2	-	-	0/1/1/1
5	MLI	E	701	-	-	2/4/4/4	-
6	FAD	A	702	-	-	8/34/50/50	0/6/6/6
5	MLI	A	701	-	-	0/4/4/4	-
10	HEM	G	201	4,3	-	7/14/54/54	-
11	MRN	G	202	-	-	6/12/12/12	0/2/2/2
8	SF4	F	302	2	-	-	0/6/5/5
10	HEM	C	201	4,3	-	4/14/54/54	-
6	FAD	E	702	-	-	7/34/50/50	0/6/6/6
9	F3S	F	303	2	-	-	0/3/3/3

The worst 5 of 42 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	C	202	MRN	C3-C8	-6.05	1.38	1.50
11	G	202	MRN	C3-C8	-5.53	1.39	1.50
11	C	202	MRN	C1-C2	-5.50	1.40	1.51
11	G	202	MRN	C1-C2	-5.36	1.41	1.51
6	A	702	FAD	C9A-C5X	5.31	1.49	1.41

The worst 5 of 79 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	G	201	HEM	CAD-C3D-C4D	6.09	135.32	124.70
10	C	201	HEM	CAD-C3D-C4D	5.65	134.54	124.70
6	E	702	FAD	C5A-C4A-N3A	-5.35	119.35	126.72
6	A	702	FAD	C5A-C4A-N3A	-5.03	119.79	126.72
11	G	202	MRN	C9-N-C8	-4.94	113.61	126.61

There are no chirality outliers.

5 of 41 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	702	FAD	N10-C1'-C2'-O2'
6	E	702	FAD	N10-C1'-C2'-O2'
6	E	702	FAD	N10-C1'-C2'-C3'
10	C	201	HEM	C2D-C3D-CAD-CBD
10	C	201	HEM	C4D-C3D-CAD-CBD


There are no ring outliers.



9 monomers are involved in 35 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	C	202	MRN	1	0
8	B	302	SF4	2	0
5	E	701	MLI	2	0
6	A	702	FAD	7	0
8	F	302	SF4	3	0
10	G	201	HEM	4	0
11	G	202	MRN	2	0
10	C	201	HEM	8	0
6	E	702	FAD	6	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	616/645 (95%)	-0.51	1 (0%) 91 80	18, 50, 76, 96	1 (0%)
1	E	616/645 (95%)	-0.49	1 (0%) 91 80	20, 59, 85, 119	1 (0%)
2	B	250/282 (88%)	-0.41	0 100 100	30, 53, 87, 115	0
2	F	250/282 (88%)	-0.48	0 100 100	30, 51, 75, 100	0
3	C	153/188 (81%)	-0.31	0 100 100	44, 61, 106, 126	0
3	G	150/188 (79%)	-0.31	0 100 100	42, 67, 133, 161	0
4	D	129/156 (82%)	-0.24	0 100 100	44, 64, 90, 125	0
4	H	129/156 (82%)	-0.30	0 100 100	45, 70, 112, 135	0
All	All	2293/2542 (90%)	-0.44	2 (0%) 92 86	18, 57, 91, 161	2 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	499	THR	2.8
1	A	499	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
11	MRN	C	202	20/20	0.91	0.15	62,68,83,83	0
11	MRN	G	202	20/20	0.91	0.20	83,90,127,129	0
10	HEM	G	201	43/43	0.95	0.12	69,75,77,78	0
5	MLI	E	701	7/7	0.95	0.08	48,49,50,51	0
10	HEM	C	201	43/43	0.95	0.13	66,72,76,78	0
6	FAD	A	702	53/53	0.97	0.07	24,26,29,30	0
6	FAD	E	702	53/53	0.97	0.07	33,35,39,40	0
5	MLI	A	701	7/7	0.97	0.09	41,43,46,46	0
9	F3S	F	303	7/7	0.98	0.05	38,41,45,47	0
9	F3S	B	303	7/7	0.99	0.04	39,41,45,46	0
7	FES	B	301	4/4	0.99	0.04	22,22,23,23	0
7	FES	F	301	4/4	1.00	0.03	29,29,30,31	0
8	SF4	B	302	8/8	1.00	0.02	26,27,27,28	0
8	SF4	F	302	8/8	1.00	0.03	28,28,29,29	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around MRN C 202:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around MRN G 202:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM G 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM C 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD A 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD E 702:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.