



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2026 – 11:21 AM UTC

PDB ID : 1YUT / pdb_00001yut
Title : Solution structure of Calcium-S100A13 (minimized mean structure)
Authors : Arnesano, F.; Banci, L.; Bertini, I.; Fantoni, A.; Tenori, L.; Viezzoli, M.S.;
Structural Proteomics in Europe (SPINE)
Deposited on : 2005-02-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	98	 76% 22% ..
1	B	98	 66% 31% .

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3284 atoms, of which 1666 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called S100 calcium-binding protein A13.

Mol	Chain	Residues	Atoms						Trace
1	A	98	Total	C	H	N	O	S	0
			1640	512	833	136	157	2	
1	B	98	Total	C	H	N	O	S	0
			1640	512	833	136	157	2	

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	
2	A	2	Total	Ca
			2	2
2	B	2	Total	Ca
			2	2

4 Residue-property plots

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: S100 calcium-binding protein A13

Chain A:  76% 22% ..



- Molecule 1: S100 calcium-binding protein A13

Chain B:  66% 31% .



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *Restrained energy minimization*.

Of the 300 calculated structures, 1 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	1.5
Amber	refinement	6

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.62	0/817 (0.0%)	1.40	7/1091 (0.6%)
1	B	0.64	0/817 (0.0%)	1.46	6/1091 (0.5%)
All	All	0.63	0/1634 (0.0%)	1.43	13/2182 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	2	5
1	B	1	11
All	All	3	16

There are no bond-length outliers.

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	96	ARG	CA-C-N	6.51	133.42	121.70
1	A	96	ARG	C-N-CA	6.51	133.42	121.70
1	B	80	ILE	CA-C-N	6.33	126.96	120.00
1	B	80	ILE	C-N-CA	6.33	126.96	120.00
1	A	95	ILE	CA-C-N	6.31	133.05	121.70
1	A	95	ILE	C-N-CA	6.31	133.05	121.70
1	B	73	PHE	CA-C-N	6.30	133.58	121.54
1	B	73	PHE	C-N-CA	6.30	133.58	121.54
1	A	97	LYS	CB-CA-C	5.88	121.28	110.10
1	A	97	LYS	CA-C-N	5.87	132.26	121.70
1	A	97	LYS	C-N-CA	5.87	132.26	121.70
1	B	27	GLU	CA-C-N	5.06	127.97	121.13

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	27	GLU	C-N-CA	5.06	127.97	121.13

All chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms
1	A	97	LYS	CA
1	A	98	LYS	CA
1	B	94	LYS	CA

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	A	1	MET	Peptide
1	A	38	PHE	Sidechain
1	A	78	ARG	Sidechain
1	A	89	LYS	Peptide
1	A	96	ARG	Sidechain
1	B	2	ALA	Peptide
1	B	3	ALA	Peptide
1	B	21	PHE	Sidechain
1	B	28	GLY	Peptide
1	B	35	VAL	Peptide
1	B	46	LEU	Peptide
1	B	64	ASP	Peptide
1	B	74	ASN	Peptide
1	B	76	TYR	Sidechain
1	B	78	ARG	Sidechain
1	B	91	LYS	Peptide

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	807	833	833	11
1	B	807	833	832	12
All	All	1618	1666	1665	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:77:TRP:CZ2	1:B:77:TRP:CZ2	0.64	2.85
1:A:77:TRP:CZ2	1:B:77:TRP:CE2	0.56	2.93
1:A:77:TRP:CZ2	1:B:81:GLY:CA	0.55	2.90
1:A:77:TRP:CE2	1:B:77:TRP:CZ2	0.55	2.95
1:A:81:GLY:CA	1:B:77:TRP:CE2	0.54	2.91
1:A:77:TRP:CE2	1:B:81:GLY:CA	0.54	2.90
1:B:7:THR:HG22	1:B:10:GLU:CD	0.53	2.29
1:A:81:GLY:N	1:B:77:TRP:CZ2	0.53	2.77
1:A:77:TRP:CE2	1:B:77:TRP:CH2	0.52	2.97
1:A:77:TRP:CZ2	1:B:81:GLY:N	0.47	2.82
1:A:81:GLY:HA2	1:B:77:TRP:CE2	0.44	2.47
1:A:77:TRP:CE2	1:B:81:GLY:HA3	0.44	2.47

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	96/98 (98%)	72 (75%)	17 (18%)	7 (7%)	1	15
1	B	96/98 (98%)	69 (72%)	20 (21%)	7 (7%)	1	15
All	All	192/196 (98%)	141 (73%)	37 (19%)	14 (7%)	1	15

All 14 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	28	GLY
1	A	29	ARG
1	A	52	ASP
1	A	92	ASP
1	A	93	LEU
1	A	96	ARG
1	A	97	LYS
1	B	3	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	29	ARG
1	B	51	LYS
1	B	52	ASP
1	B	65	VAL
1	B	74	ASN
1	B	95	ILE

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	91/91 (100%)	81 (89%)	10 (11%)	8	51
1	B	91/91 (100%)	80 (88%)	11 (12%)	7	49
All	All	182/182 (100%)	161 (88%)	21 (12%)	8	50

All 21 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	4	GLU
1	A	25	ARG
1	A	26	GLN
1	A	32	SER
1	A	46	LEU
1	A	51	LYS
1	A	59	LYS
1	A	72	LYS
1	A	88	ARG
1	A	91	LYS
1	B	4	GLU
1	B	26	GLN
1	B	45	GLN
1	B	51	LYS
1	B	59	LYS
1	B	72	LYS
1	B	83	LEU
1	B	85	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	93	LEU
1	B	94	LYS
1	B	98	LYS

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided