



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 04:29 PM UTC

PDB ID : 4YVW / pdb_00004yvw
Title : crystal structure of an enterovirus 71/coxsackievirus A16 chimeric virus-like particle
Authors : Chen, R.; Lyu, K.
Deposited on : 2015-03-20
Resolution : 3.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1065 (3.96-3.64)
Clashscore	190562	1012 (3.94-3.66)
Ramachandran outliers	187476	1048 (3.96-3.64)
Sidechain outliers	187428	1043 (3.96-3.64)
RSRZ outliers	180081	1064 (3.96-3.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	2% 50% 25% • 24%
1	D	297	2% 46% 28% • 24%
1	E	297	2% 46% 28% • 24%
1	J	297	2% 48% 27% • 24%
1	M	297	2% 43% 30% • 24%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	242	
2	F	242	
2	H	242	
2	K	242	
2	N	242	
3	C	323	
3	G	323	
3	I	323	
3	L	323	
3	O	323	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 26525 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Capsid protein VP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	225	Total	C	N	O	S	0	0	0
			1773	1134	299	329	11			
1	A	225	Total	C	N	O	S	0	0	0
			1773	1134	299	329	11			
1	D	225	Total	C	N	O	S	0	0	0
			1773	1134	299	329	11			
1	J	225	Total	C	N	O	S	0	0	0
			1773	1134	299	329	11			
1	M	225	Total	C	N	O	S	0	0	0
			1773	1134	299	329	11			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	215	LEU	LYS	engineered mutation	UNP F6KTB0
E	217	ALA	GLU	engineered mutation	UNP F6KTB0
E	218	ASN	LYS	engineered mutation	UNP F6KTB0
E	221	ASP	GLU	engineered mutation	UNP F6KTB0
A	215	LEU	LYS	engineered mutation	UNP F6KTB0
A	217	ALA	GLU	engineered mutation	UNP F6KTB0
A	218	ASN	LYS	engineered mutation	UNP F6KTB0
A	221	ASP	GLU	engineered mutation	UNP F6KTB0
D	215	LEU	LYS	engineered mutation	UNP F6KTB0
D	217	ALA	GLU	engineered mutation	UNP F6KTB0
D	218	ASN	LYS	engineered mutation	UNP F6KTB0
D	221	ASP	GLU	engineered mutation	UNP F6KTB0
J	215	LEU	LYS	engineered mutation	UNP F6KTB0
J	217	ALA	GLU	engineered mutation	UNP F6KTB0
J	218	ASN	LYS	engineered mutation	UNP F6KTB0
J	221	ASP	GLU	engineered mutation	UNP F6KTB0
M	215	LEU	LYS	engineered mutation	UNP F6KTB0
M	217	ALA	GLU	engineered mutation	UNP F6KTB0
M	218	ASN	LYS	engineered mutation	UNP F6KTB0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
M	221	ASP	GLU	engineered mutation	UNP F6KTB0

- Molecule 2 is a protein called Capsid protein VP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	224	Total	C	N	O	S	0	0	0
			1712	1103	281	317	11			
2	B	224	Total	C	N	O	S	0	0	0
			1712	1103	281	317	11			
2	H	224	Total	C	N	O	S	0	0	0
			1712	1103	281	317	11			
2	K	224	Total	C	N	O	S	0	0	0
			1712	1103	281	317	11			
2	N	224	Total	C	N	O	S	0	0	0
			1712	1103	281	317	11			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	227	GLN	LYS	engineered mutation	UNP F6KTB0
B	227	GLN	LYS	engineered mutation	UNP F6KTB0
H	227	GLN	LYS	engineered mutation	UNP F6KTB0
K	227	GLN	LYS	engineered mutation	UNP F6KTB0
N	227	GLN	LYS	engineered mutation	UNP F6KTB0


- Molecule 3 is a protein called Capsid protein VP0.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	235	Total	C	N	O	S	0	0	0
			1820	1170	299	343	8			
3	C	235	Total	C	N	O	S	0	0	0
			1820	1170	299	343	8			
3	I	235	Total	C	N	O	S	0	0	0
			1820	1170	299	343	8			
3	L	235	Total	C	N	O	S	0	0	0
			1820	1170	299	343	8			
3	O	235	Total	C	N	O	S	0	0	0
			1820	1170	299	343	8			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Capsid protein VP1




• Molecule 1: Capsid protein VP1




• Molecule 1: Capsid protein VP1







• Molecule 1: Capsid protein VP1




• Molecule 1: Capsid protein VP1




• Molecule 2: Capsid protein VP3







• Molecule 2: Capsid protein VP3



• Molecule 2: Capsid protein VP3




• Molecule 2: Capsid protein VP3









• Molecule 2: Capsid protein VP3




• Molecule 3: Capsid protein VP0




• Molecule 3: Capsid protein VP0







• Molecule 3: Capsid protein VP0




• Molecule 3: Capsid protein VP0



• Molecule 3: Capsid protein VP0





4 Data and refinement statistics

Property	Value	Source
Space group	P 42 3 2	Depositor
Cell constants a, b, c, α , β , γ	349.15Å 349.15Å 349.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.96 – 3.80 47.96 – 3.80	Depositor EDS
% Data completeness (in resolution range)	78.7 (47.96-3.80) 61.3 (47.96-3.80)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.52 (at 3.77Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1839)	Depositor
R, R_{free}	0.286 , 0.324 0.289 , 0.326	Depositor DCC
R_{free} test set	2000 reflections (3.52%)	wwPDB-VP
Wilson B-factor (Å ²)	94.0	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	26525	wwPDB-VP
Average B, all atoms (Å ²)	112.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/1828	0.90	3/2495 (0.1%)
1	D	0.36	0/1828	0.89	2/2495 (0.1%)
1	E	0.34	0/1828	0.94	7/2495 (0.3%)
1	J	0.34	0/1828	0.87	3/2495 (0.1%)
1	M	0.36	0/1828	0.90	5/2495 (0.2%)
2	B	0.35	0/1759	0.93	8/2408 (0.3%)
2	F	0.35	0/1759	0.94	5/2408 (0.2%)
2	H	0.35	0/1759	0.94	8/2408 (0.3%)
2	K	0.34	0/1759	0.92	2/2408 (0.1%)
2	N	0.35	0/1759	0.90	8/2408 (0.3%)
3	C	0.34	0/1875	1.00	10/2573 (0.4%)
3	G	0.36	0/1875	0.94	9/2573 (0.3%)
3	I	0.39	0/1875	0.99	15/2573 (0.6%)
3	L	0.37	0/1875	1.02	19/2573 (0.7%)
3	O	0.38	0/1875	0.95	5/2573 (0.2%)
All	All	0.36	0/27310	0.94	109/37380 (0.3%)

There are no bond length outliers.

The worst 5 of 109 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	220	LEU	N-CA-C	-8.63	102.51	113.72
2	F	169	ILE	CA-C-N	8.53	127.84	118.97
2	F	169	ILE	C-N-CA	8.53	127.84	118.97
3	C	88	GLU	N-CA-C	-7.97	104.01	112.93
3	C	232	ILE	CA-C-N	7.75	128.04	119.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1773	0	1712	72	0
1	D	1773	0	1712	82	0
1	E	1773	0	1712	85	0
1	J	1773	0	1712	84	0
1	M	1773	0	1712	94	0
2	B	1712	0	1699	71	0
2	F	1712	0	1699	93	0
2	H	1712	0	1699	79	2
2	K	1712	0	1699	67	0
2	N	1712	0	1699	75	0
3	C	1820	0	1758	67	0
3	G	1820	0	1758	82	0
3	I	1820	0	1758	95	1
3	L	1820	0	1758	83	0
3	O	1820	0	1758	94	0
All	All	26525	0	25845	1014	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 1014 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:189:GLN:HE21	2:N:21:SER:HB3	1.28	0.95
3:L:26:GLN:HE21	3:L:26:GLN:HA	1.33	0.93
1:J:262:ILE:HG21	3:L:128:PRO:HG2	1.52	0.90
3:G:132:ILE:HG12	3:G:150:GLN:HE21	1.40	0.87
1:E:262:ILE:HG21	3:G:128:PRO:HG2	1.57	0.87

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:60:THR:OG1	2:H:60:THR:OG1[14_777]	2.02	0.18

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:158:ASP:OD2	3:I:113:ASN:ND2[5_636]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	223/297 (75%)	199 (89%)	22 (10%)	2 (1%)	14	45
1	D	223/297 (75%)	202 (91%)	21 (9%)	0	100	100
1	E	223/297 (75%)	198 (89%)	23 (10%)	2 (1%)	14	45
1	J	223/297 (75%)	199 (89%)	23 (10%)	1 (0%)	30	62
1	M	223/297 (75%)	199 (89%)	22 (10%)	2 (1%)	14	45
2	B	220/242 (91%)	198 (90%)	20 (9%)	2 (1%)	14	45
2	F	220/242 (91%)	201 (91%)	17 (8%)	2 (1%)	14	45
2	H	220/242 (91%)	200 (91%)	20 (9%)	0	100	100
2	K	220/242 (91%)	201 (91%)	19 (9%)	0	100	100
2	N	220/242 (91%)	199 (90%)	20 (9%)	1 (0%)	24	57
3	C	233/323 (72%)	196 (84%)	34 (15%)	3 (1%)	9	38
3	G	233/323 (72%)	197 (84%)	32 (14%)	4 (2%)	7	34
3	I	233/323 (72%)	198 (85%)	32 (14%)	3 (1%)	9	38
3	L	233/323 (72%)	195 (84%)	36 (16%)	2 (1%)	14	45
3	O	233/323 (72%)	196 (84%)	34 (15%)	3 (1%)	9	38
All	All	3380/4310 (78%)	2978 (88%)	375 (11%)	27 (1%)	16	48

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	128	PRO
3	I	145	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	G	27	GLU
3	G	128	PRO
3	I	128	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/250 (77%)	187 (97%)	5 (3%)	40	60
1	D	192/250 (77%)	185 (96%)	7 (4%)	31	54
1	E	192/250 (77%)	188 (98%)	4 (2%)	47	64
1	J	192/250 (77%)	189 (98%)	3 (2%)	55	68
1	M	192/250 (77%)	182 (95%)	10 (5%)	21	46
2	B	188/202 (93%)	178 (95%)	10 (5%)	20	45
2	F	188/202 (93%)	177 (94%)	11 (6%)	18	43
2	H	188/202 (93%)	178 (95%)	10 (5%)	20	45
2	K	188/202 (93%)	178 (95%)	10 (5%)	20	45
2	N	188/202 (93%)	173 (92%)	15 (8%)	11	36
3	C	200/272 (74%)	193 (96%)	7 (4%)	32	54
3	G	200/272 (74%)	194 (97%)	6 (3%)	36	57
3	I	200/272 (74%)	191 (96%)	9 (4%)	24	48
3	L	200/272 (74%)	191 (96%)	9 (4%)	24	48
3	O	200/272 (74%)	193 (96%)	7 (4%)	32	54
All	All	2900/3620 (80%)	2777 (96%)	123 (4%)	26	50

5 of 123 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	46	LEU
3	L	76	LYS
2	K	132	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	L	72	GLU
3	O	126	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

Mol	Chain	Res	Type
2	K	162	GLN
3	C	119	GLN
2	N	48	GLN
2	N	221	GLN
3	C	161	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	225/297 (75%)	0.31	3 (1%)	75 53	75, 100, 167, 219	0
1	D	225/297 (75%)	0.33	6 (2%)	56 38	74, 96, 172, 219	0
1	E	225/297 (75%)	0.27	5 (2%)	62 44	77, 98, 163, 222	0
1	J	225/297 (75%)	0.34	6 (2%)	56 38	76, 99, 170, 212	0
1	M	225/297 (75%)	0.37	7 (3%)	51 35	75, 100, 172, 220	0
2	B	224/242 (92%)	0.36	5 (2%)	62 44	77, 101, 128, 163	0
2	F	224/242 (92%)	0.38	7 (3%)	51 35	78, 96, 124, 153	0
2	H	224/242 (92%)	0.41	8 (3%)	46 32	76, 96, 122, 156	0
2	K	224/242 (92%)	0.32	3 (1%)	75 53	78, 101, 127, 168	0
2	N	224/242 (92%)	0.31	7 (3%)	51 35	79, 103, 127, 161	0
3	C	235/323 (72%)	0.55	8 (3%)	48 33	76, 121, 195, 213	0
3	G	235/323 (72%)	0.46	8 (3%)	48 33	75, 116, 191, 212	0
3	I	235/323 (72%)	0.65	18 (7%)	19 18	74, 117, 200, 223	0
3	L	235/323 (72%)	0.49	11 (4%)	36 27	78, 122, 201, 226	0
3	O	235/323 (72%)	0.62	15 (6%)	25 21	79, 124, 193, 222	0
All	All	3420/4310 (79%)	0.41	117 (3%)	48 33	74, 102, 183, 226	0

The worst 5 of 117 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	I	247	GLY	5.2
3	O	16	LEU	5.0
3	O	247	GLY	5.0
3	L	246	ALA	4.4
2	N	235	ILE	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.