



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 04:09 AM UTC

PDB ID : 5YVQ / pdb_00005yvq
Title : Complex of Mu phage tail fiber and its chaperone
Authors : Takeda, S.; Sakai, K.; Iwazaki, T.; Yamashita, E.; Nakagawa, A.
Deposited on : 2017-11-27
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	504	9% 67% 29%
2	B	175	33% 54% 5% 41%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4099 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tail fiber protein S.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	358	2799	1737	511	539	12	0	3	0

- Molecule 2 is a protein called Tail fiber assembly protein U.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	103	855	546	140	166	3	0	1	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	388	Total	O	0	0
			388	388		
4	B	27	Total	O	0	0
			27	27		

4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	66.33Å 66.33Å 387.77Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	32.14 – 2.10 32.14 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (32.14-2.10) 99.6 (32.14-2.10)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.10Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.236 , 0.264 0.237 , 0.264	Depositor DCC
R_{free} test set	2973 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.604	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 57.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.039 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4099	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2860	0.57	0/3866
2	B	0.16	0/878	0.29	0/1190
All	All	0.34	0/3738	0.52	0/5056

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2799	0	2703	15	1
2	B	855	0	830	6	0
3	A	30	0	40	3	0
4	A	388	0	0	1	3
4	B	27	0	0	1	0
All	All	4099	0	3573	21	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:36:ASN:OD1	4:B:201:HOH:O	2.07	0.73
1:A:405[A]:ARG:NH2	4:A:705:HOH:O	2.32	0.62
2:B:6:ASN:ND2	2:B:32:GLU:OE1	2.37	0.57
1:A:471:HIS:HB3	1:A:484:LYS:HB2	1.87	0.56
1:A:177:ASN:OD1	1:A:180:ARG:NH2	2.41	0.54
1:A:309:MET:H	3:A:604:GOL:C3	2.21	0.52
1:A:138:LYS:HD3	1:A:141:LYS:HD3	1.93	0.51
1:A:451:LEU:HD11	1:A:486:TYR:CE1	2.46	0.50
1:A:253:MET:HE2	1:A:262:VAL:HG22	1.93	0.49
1:A:449:GLN:CD	1:A:486:TYR:HD1	2.22	0.48
2:B:95:MET:HE3	2:B:103:LYS:O	2.14	0.48
2:B:10:GLU:HB2	2:B:30[B]:TRP:CD1	2.50	0.47
2:B:52:VAL:HB	2:B:61:ALA:HB3	1.97	0.46
1:A:394:LYS:NZ	3:A:605:GOL:H2	2.30	0.46
1:A:451:LEU:HD11	1:A:486:TYR:CD1	2.51	0.45
1:A:416:MET:HA	1:A:421:ASN:O	2.17	0.44
1:A:394:LYS:HZ1	3:A:605:GOL:H2	1.84	0.43
2:B:30[B]:TRP:CZ3	2:B:36:ASN:HB2	2.54	0.42
1:A:454:GLY:H	1:A:484:LYS:HD3	1.85	0.42
1:A:449:GLN:HB2	1:A:488:ARG:CZ	2.51	0.41
1:A:454:GLY:N	1:A:484:LYS:HD3	2.36	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:ASP:OD1	4:A:708:HOH:O[5_666]	2.08	0.12
4:A:888:HOH:O	4:A:1044:HOH:O[5_566]	2.13	0.07
4:A:1011:HOH:O	4:A:1031:HOH:O[5_666]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/504 (71%)	348 (98%)	9 (2%)	0	100	100
2	B	102/175 (58%)	100 (98%)	2 (2%)	0	100	100
All	All	459/679 (68%)	448 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	295/412 (72%)	295 (100%)	0	100	100
2	B	95/157 (60%)	95 (100%)	0	100	100
All	All	390/569 (68%)	390 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	169	ASN
1	A	173	GLN
1	A	217	ASN
1	A	234	ASN
1	A	408	ASN
2	B	36	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	602	-	5,5,5	0.39	0	5,5,5	0.31	0
3	GOL	A	603	-	5,5,5	0.33	0	5,5,5	0.40	0
3	GOL	A	605	-	5,5,5	0.45	0	5,5,5	1.11	0
3	GOL	A	601	-	5,5,5	0.32	0	5,5,5	0.36	0
3	GOL	A	604	-	5,5,5	0.37	0	5,5,5	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	602	-	-	2/4/4/4	-
3	GOL	A	603	-	-	2/4/4/4	-
3	GOL	A	605	-	-	3/4/4/4	-
3	GOL	A	601	-	-	0/4/4/4	-
3	GOL	A	604	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	GOL	O1-C1-C2-C3
3	A	603	GOL	O1-C1-C2-C3
3	A	604	GOL	O1-C1-C2-C3
3	A	605	GOL	O1-C1-C2-C3
3	A	602	GOL	O1-C1-C2-O2
3	A	605	GOL	O1-C1-C2-O2
3	A	603	GOL	O1-C1-C2-O2
3	A	604	GOL	O1-C1-C2-O2
3	A	605	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	605	GOL	2	0
3	A	604	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	358/504 (71%)	0.72	43 (12%)	9 9	13, 28, 80, 161	3 (0%)
2	B	103/175 (58%)	2.55	58 (56%)	0 0	36, 75, 119, 176	1 (0%)
All	All	461/679 (67%)	1.13	101 (21%)	2 2	13, 33, 104, 176	4 (0%)

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	139	ALA	11.3
2	B	102	VAL	9.9
2	B	84	ALA	8.4
1	A	470	PHE	7.4
1	A	140	VAL	7.0
1	A	455	LEU	6.2
2	B	101	VAL	6.0
2	B	103	LYS	5.5
1	A	162	ASN	5.5
1	A	485	VAL	5.5
2	B	100	ALA	5.4
1	A	451	LEU	5.4
1	A	138	LYS	5.2
2	B	95	MET	5.2
1	A	484	LYS	5.1
2	B	80	PRO	4.8
2	B	83	THR	4.8
1	A	454	GLY	4.7
2	B	99	GLY	4.6
2	B	88	ALA	4.4
2	B	91	SER	4.4
1	A	486	TYR	4.3
2	B	44	PHE	4.3
1	A	159	ASP	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	97	LYS	4.0
1	A	472	THR	3.9
2	B	81	ASP	3.9
1	A	456	SER	3.9
2	B	1	MET	3.9
2	B	15	LYS	3.7
2	B	92	GLY	3.6
2	B	58	ILE	3.6
2	B	82	ILE	3.6
2	B	7	ILE	3.6
1	A	453	GLY	3.6
2	B	98	ASP	3.5
2	B	89	ASP	3.5
2	B	42	ASN	3.5
1	A	457	ARG	3.5
1	A	459	TYR	3.5
2	B	77	VAL	3.5
1	A	482	ASP	3.4
1	A	142	ILE	3.4
2	B	4	LEU	3.4
1	A	155	ARG	3.4
2	B	86	ARG	3.3
2	B	30[A]	TRP	3.3
2	B	17	GLN	3.2
2	B	51	ILE	3.2
2	B	3	HIS	3.0
2	B	79	ILE	3.0
1	A	504	ALA	3.0
2	B	18	TYR	3.0
1	A	141	LYS	3.0
2	B	87	ARG	2.9
2	B	53	TYR	2.9
2	B	13	LYS	2.9
1	A	163	LYS	2.8
2	B	12	PRO	2.7
1	A	483	ASP	2.7
1	A	161	PRO	2.7
2	B	90	ASP	2.7
1	A	466	VAL	2.6
1	A	149	ALA	2.6
1	A	168	GLN	2.6
2	B	41	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	489	PRO	2.6
1	A	151	LEU	2.6
1	A	461	ALA	2.5
2	B	2	MET	2.5
2	B	35	LYS	2.5
2	B	36	ASN	2.5
2	B	96	PHE	2.4
2	B	72	GLU	2.4
2	B	31	SER	2.4
1	A	197	GLU	2.4
1	A	150	ARG	2.4
2	B	76	VAL	2.4
2	B	75	SER	2.4
2	B	8	LYS	2.3
1	A	160	ILE	2.3
2	B	52	VAL	2.3
2	B	85	ASN	2.3
1	A	218	ASP	2.3
2	B	62	ILE	2.3
2	B	60	VAL	2.3
2	B	93	LYS	2.3
1	A	190	LEU	2.3
2	B	16	GLU	2.2
1	A	452	THR	2.2
2	B	55	GLU	2.2
2	B	45	GLN	2.2
1	A	469	GLY	2.2
1	A	450	SER	2.1
2	B	33	ASP	2.1
2	B	94	TRP	2.1
1	A	502	ALA	2.1
1	A	458	ASP	2.1
1	A	471	HIS	2.1
1	A	174	GLU	2.0
2	B	26	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	602	6/6	0.72	0.33	93,96,96,98	0
3	GOL	A	604	6/6	0.73	0.27	39,47,58,62	0
3	GOL	A	603	6/6	0.76	0.18	60,64,67,70	0
3	GOL	A	605	6/6	0.81	0.17	37,41,44,45	0
3	GOL	A	601	6/6	0.83	0.16	46,53,56,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.