



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:22 PM UTC

PDB ID : 6YVI / pdb_00006yvi
Title : EED in complex with a cyano-benzofuran
Authors : Read, J.A.
Deposited on : 2020-04-28
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	 % 83% 11% . .
1	B	366	 5% 80% 13% 6%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5865 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Polycomb protein EED.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	350	Total	C	N	O	S	0	0	0
			2793	1771	490	511	21			
1	B	344	Total	C	N	O	S	0	0	0
			2720	1725	470	504	21			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is N-(2,3-dihydro-1-benzofuran-4-ylmethyl)-8-(4-methylsulfonylphenyl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (CCD ID: L9W) (formula: C₂₁H₁₉N₅O₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			30	21	5	3	1		

- Molecule 4 is 5-fluoranyl-4-[[[8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-yl]amino]methyl]-2,3-dihydro-1-benzofuran-7-carbonitrile (CCD ID: PV5) (formula: C₂₁H₁₆FN₇O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	F	N	O	0	0
			30	21	1	7	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	175	Total	O	0	0
			175	175		
5	B	116	Total	O	0	0
			116	116		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Polycomb protein EED



• Molecule 1: Polycomb protein EED



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.89Å 86.53Å 97.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.36 – 2.26 64.36 – 2.27	Depositor EDS
% Data completeness (in resolution range)	67.2 (64.36-2.26) 67.6 (64.36-2.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 2.27Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.182 , 0.241 0.183 , 0.241	Depositor DCC
R_{free} test set	1194 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5865	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, L9W, PV5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/2864 (0.0%)	1.17	5/3887 (0.1%)
1	B	0.85	2/2787 (0.1%)	1.21	7/3785 (0.2%)
All	All	0.86	3/5651 (0.1%)	1.19	12/7672 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	290	ILE	CG1-CD1	-7.44	1.22	1.51
1	A	343	ILE	CA-CB	5.53	1.60	1.54
1	B	366	MET	SD-CE	-5.49	1.65	1.79

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	142	ASP	CA-CB-CG	6.64	119.24	112.60
1	A	342	LYS	CA-C-N	6.27	128.97	120.63
1	A	342	LYS	C-N-CA	6.27	128.97	120.63
1	B	381	ASN	CA-CB-CG	6.09	118.69	112.60
1	A	381	ASN	CA-CB-CG	6.07	118.67	112.60
1	B	146	ASN	CA-CB-CG	5.97	118.57	112.60
1	B	128	SER	CA-C-N	5.75	132.52	121.54
1	B	128	SER	C-N-CA	5.75	132.52	121.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	342	LYS	CA-C-N	5.21	128.38	120.77
1	B	342	LYS	C-N-CA	5.21	128.38	120.77
1	A	182	CYS	CA-C-N	5.02	126.98	120.56
1	A	182	CYS	C-N-CA	5.02	126.98	120.56

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	313	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2793	0	2669	19	0
1	B	2720	0	2568	19	0
2	A	1	0	0	0	0
3	A	30	0	0	0	0
4	B	30	0	0	0	0
5	A	175	0	0	0	0
5	B	116	0	0	0	0
All	All	5865	0	5237	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:186:TYR:OH	1:B:220:ILE:HG22	1.94	0.68
1:B:269:ARG:HE	1:B:292:GLN:HE21	1.44	0.66
1:B:120:ARG:HD3	1:B:139:VAL:HG22	1.80	0.64
1:A:215:LEU:HB2	1:A:229:PHE:HB2	1.82	0.62
1:A:215:LEU:HD21	1:A:253:SER:HB3	1.82	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:215:LEU:HB2	1:B:229:PHE:HB2	1.82	0.60
1:B:127:HIS:HB2	1:B:131:GLU:HB3	1.84	0.59
1:B:230:GLY:O	1:B:295:HIS:HA	2.03	0.57
1:A:281:ASN:HB3	1:A:284:LYS:HG2	1.87	0.57
1:B:215:LEU:HD21	1:B:253:SER:HB3	1.86	0.55
1:B:254:CYS:HB2	1:B:309:VAL:HB	1.89	0.55
1:A:254:CYS:HB2	1:A:309:VAL:HB	1.88	0.54
1:A:168:SER:HA	1:A:192:ALA:HB2	1.92	0.52
1:B:168:SER:HA	1:B:192:ALA:HB2	1.92	0.52
1:A:127:HIS:HB2	1:A:131:GLU:HB3	1.91	0.51
1:B:138:TYR:HB2	1:B:180:MET:HG3	1.94	0.50
1:A:343:ILE:HD12	1:A:347:GLU:HG2	1.95	0.49
1:B:168:SER:HA	1:B:192:ALA:CB	2.43	0.49
1:A:120:ARG:HD2	1:A:139:VAL:HG22	1.94	0.48
1:B:176:ASN:HB2	1:B:183:ILE:HD11	1.95	0.48
1:A:168:SER:HA	1:A:192:ALA:CB	2.44	0.48
1:A:360:GLN:O	1:A:381:ASN:HB2	2.14	0.47
1:B:269:ARG:CZ	1:B:294:ILE:HD13	2.45	0.47
1:B:416:THR:HA	1:B:426:ILE:O	2.15	0.46
1:A:169:ARG:HB3	1:A:171:ILE:HD12	1.98	0.45
1:B:343:ILE:HD12	1:B:347:GLU:HG2	1.99	0.45
1:A:338:ASP:HB3	1:A:343:ILE:HD11	1.99	0.45
1:A:250:LYS:HD2	1:A:262:LEU:HD11	1.97	0.45
1:A:232:VAL:HG22	1:A:295:HIS:HB3	1.98	0.44
1:A:110:PRO:HB3	1:A:125:GLU:OE1	2.17	0.44
1:B:360:GLN:O	1:B:381:ASN:HB2	2.17	0.43
1:A:106:LYS:O	1:A:109:ASP:HB2	2.18	0.43
1:A:416:THR:HA	1:A:426:ILE:O	2.19	0.42
1:B:106:LYS:O	1:B:109:ASP:HB2	2.20	0.42
1:A:262:LEU:HB3	1:A:299:PHE:HB3	2.01	0.42
1:B:262:LEU:HB3	1:B:299:PHE:HB3	2.01	0.41
1:A:242:ALA:HA	1:A:252:MET:O	2.20	0.41
1:B:188:GLY:O	1:B:218:TRP:HZ2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/366 (94%)	333 (96%)	13 (4%)	0	100	100
1	B	338/366 (92%)	325 (96%)	11 (3%)	2 (1%)	21	21
All	All	684/732 (93%)	658 (96%)	24 (4%)	2 (0%)	36	40

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	129	GLN
1	B	187	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/329 (92%)	291 (96%)	11 (4%)	31	39
1	B	291/329 (88%)	284 (98%)	7 (2%)	43	54
All	All	593/658 (90%)	575 (97%)	18 (3%)	36	45

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	88	LEU
1	A	120	ARG
1	A	137	SER
1	A	236	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	246	LEU
1	A	260	LEU
1	A	286	ASN
1	A	293	LYS
1	A	362	ASP
1	A	367	ARG
1	A	386	LEU
1	B	88	LEU
1	B	184	LYS
1	B	228	ILE
1	B	260	LEU
1	B	355	ARG
1	B	362	ASP
1	B	386	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	ASN
1	A	146	ASN
1	A	176	ASN
1	A	185	HIS
1	A	349	ASN
1	B	146	ASN
1	B	292	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PV5	B	501	-	34,34,34	0.49	0	41,49,49	1.29	3 (7%)
3	L9W	A	502	-	34,34,34	0.51	1 (2%)	43,50,50	1.27	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PV5	B	501	-	-	1/11/17/17	0/5/5/5
3	L9W	A	502	-	-	1/15/21/21	0/5/5/5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	L9W	C8-C7	2.23	1.41	1.37

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	L9W	N4-C9-N1	5.68	120.31	116.93
4	B	501	PV5	N5-C8-N2	5.61	120.26	116.93
4	B	501	PV5	C10-N2-C8	2.53	133.38	130.41
4	B	501	PV5	C7-C6-C9	-2.42	116.32	119.44
3	A	502	L9W	C19-C18-C17	2.39	110.18	107.82
3	A	502	L9W	C8-C7-C10	-2.28	116.50	119.44

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	502	L9W	N4-C12-C13-C14
4	B	501	PV5	N5-C11-C12-C17

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	350/366 (95%)	-0.25	3 (0%) 81 82	11, 27, 53, 66	0
1	B	344/366 (93%)	0.16	18 (5%) 33 31	15, 38, 68, 94	0
All	All	694/732 (94%)	-0.04	21 (3%) 52 53	11, 33, 63, 94	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	279	ASP	4.1
1	B	188	GLY	4.0
1	B	392	GLU	3.8
1	B	186	TYR	3.8
1	B	178	ILE	3.5
1	A	392	GLU	3.0
1	B	364	TRP	2.9
1	B	187	VAL	2.6
1	A	232	VAL	2.5
1	B	185	HIS	2.5
1	B	179	THR	2.5
1	B	302	ARG	2.4
1	B	373	TRP	2.4
1	B	108	GLY	2.3
1	B	401	CYS	2.3
1	B	339	ASP	2.3
1	B	143	ALA	2.2
1	B	129	GLN	2.1
1	B	181	GLN	2.1
1	A	401	CYS	2.0
1	B	160	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PV5	B	501	30/30	0.90	0.10	29,38,53,54	0
2	CA	A	501	1/1	0.94	0.13	58,58,58,58	0
3	L9W	A	502	30/30	0.95	0.07	12,24,37,41	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PV5 B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around L9W A 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.