



wwPDB EM Validation Summary Report ⓘ

Mar 29, 2026 – 11:52 PM UTC

PDB ID : 7YVS / pdb_00007yvs
EMDB ID : EMD-34137
Title : Complex structure of Clostridioides difficile binary toxin unfolded CDTa-bound CDTb-pore (short).
Authors : Yamada, T.; Kawamoto, A.; Yoshida, T.; Sato, Y.; Kato, T.; Tsuge, H.
Deposited on : 2022-08-19
Resolution : 2.80 Å(reported)
Based on initial model : 7VNJ

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	11806 (2.30 - 3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	675	 5% 63% 10% 22%
1	B	675	 2% 63% 11% 24%
1	C	675	 2% 62% 11% 25%
1	D	675	 2% 61% 12% 25%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	675	
1	F	675	
1	G	675	
2	H	428	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 30593 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribosylating binary toxin binding subunit CdtB.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	494	Total	C	N	O	S	11	0
			3935	2458	640	831	6		
1	B	494	Total	C	N	O	S	11	0
			3935	2458	640	831	6		
1	C	494	Total	C	N	O	S	1	0
			3867	2418	627	816	6		
1	D	497	Total	C	N	O	S	1	0
			3898	2438	631	823	6		
1	E	494	Total	C	N	O	S	11	0
			3935	2458	640	831	6		
1	F	494	Total	C	N	O	S	1	0
			3867	2418	627	816	6		
1	G	494	Total	C	N	O	S	21	0
			3935	2458	640	831	6		

- Molecule 2 is a protein called ADP-ribosylating binary toxin enzymatic subunit CdtA.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	H	395	Total	C	N	O	S	0	0
			3200	2038	534	623	5		

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	414	GLU	-	expression tag	UNP Q9KH42
H	415	ASN	-	expression tag	UNP Q9KH42
H	416	LEU	-	expression tag	UNP Q9KH42
H	417	TYR	-	expression tag	UNP Q9KH42
H	418	PHE	-	expression tag	UNP Q9KH42
H	419	GLN	-	expression tag	UNP Q9KH42
H	420	GLY	-	expression tag	UNP Q9KH42
H	421	LEU	-	expression tag	UNP Q9KH42
H	422	GLU	-	expression tag	UNP Q9KH42

Continued on next page...


Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
H	423	HIS	-	expression tag	UNP Q9KH42
H	424	HIS	-	expression tag	UNP Q9KH42
H	425	HIS	-	expression tag	UNP Q9KH42
H	426	HIS	-	expression tag	UNP Q9KH42
H	427	HIS	-	expression tag	UNP Q9KH42
H	428	HIS	-	expression tag	UNP Q9KH42

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		AltConf
3	A	3	Total 3	Ca 3	0
3	B	3	Total 3	Ca 3	0
3	C	3	Total 3	Ca 3	0
3	D	3	Total 3	Ca 3	0
3	E	3	Total 3	Ca 3	0
3	F	3	Total 3	Ca 3	0
3	G	3	Total 3	Ca 3	0

- Molecule 1: ADP-ribosylating binary toxin binding subunit CdtB






- Molecule 1: ADP-ribosylating binary toxin binding subunit CdtB



- Molecule 1: ADP-ribosylating binary toxin binding subunit CdtB



ASN TYR ASN ASN PHE PHE ASP PRO ASN ASN PRO LEU THR LYS MET THR ASP THR VAL ASN LEU ARG SER TYR PHE THR GLY GLY GLU ASN ILE MET THR TYR LYS LYS LEU ARG ILE TYR TYR ILE ALA THR THR PRO ASP ASP ARG ASP GLU LEU VAL LEU VAL SER VAL ASP

• Molecule 1: ADP-ribosylating binary toxin binding subunit CdtB

Chain G:  6% 62% 11% 27%

ASN ASN ASN PHE PHE ASP PRO ASN ASN PRO LEU THR LYS MET THR ASP THR VAL ASN LEU ARG SER TYR PHE THR GLY GLY GLU ASN ILE MET THR TYR LYS LYS LEU ARG ILE TYR TYR ILE ALA THR THR PRO ASP ASP ARG ASP GLU LEU VAL LEU VAL SER VAL ASP

ASN GLY PHE THR ALA VAL THR THR TYR TYR SER HIS THR THR ASP ASP S364 T365 A366 S370 T377 S380 A387 N392 V393 R394 D417 L444 S445 P446 L447 A448 I462 L470 L478 K490 M491 S492 S493 G494 Q495 S516 L545 K559 K560

D561 D568 D572 V576 F580 T584 L595 N602 E606 M609 N610 F619 F622 D623 D624 M632 V635 M636 T637 T638 D641 S646 A647 N648 K649 L650 N651 E652 T654 K655 I656 K657 I658 P659 M660 S661 E662 Y666 V670 D677 P678

L679 T680 S681 N682 S683 K687 E692 E693 K694 T695 D696 Y697 L698 V699 P700 E701 Q702 Y709 E710 T714 K716 E717 S718 S719 N720 N721 I722 E723 I724 L725 L726 G727 S728 G729 T730 T731 S737 N742 SER THR PRO ILE LEU ASP PRO ILE VAL LYS THR

ASP GLN ILE MET ASP HIS LYS ILE PHE LEU ASP PRO ASN PHE ASN PRO LYS THR VAL ASN THR ILE ASN MET TYR PHE GLY GLY GLU ASN ILE MET THR TYR LYS ALA THR ASP TYR ARG VAL ALA THR THR LEU LEU VAL TYR GLY PHE LYS ASP

ILE GLY THR LYS ASP LYS MET ARG ASN TYR LEU ASP PRO ASN PHE ASN PRO LYS THR VAL ASN THR ILE ARG SER TYR PHE GLY GLY GLU ASN ILE MET THR TYR LYS LEU LEU ILE THR PRO ASP ARG ASP GLU LEU LEU VAL LEU SER VAL ASP

• Molecule 2: ADP-ribosylating binary toxin enzymatic subunit CdtA

Chain H:  9% 73% 19% 8%

ALA PRO ILE ARG PRO ASP PHE LEU LYS ASP LYS GLU LYS LYS ALA LYS GLU W19 R21 R22 E23 A24 E25 R26 I27 E28 K29 Q29 K30 L31 E32 R33 E40 D45 E48 I49 S50 Y52 S53 Q54 T55 R56 M57 E70 D86 E109 E113 L116 E117 K118

F122 I126 D134 Y142 H157 L158 M166 T172 N173 N174 V175 S176 I179 R192 I193 V194 K198 H199 K202 V207 D215 N233 K234 L235 T236 P237 N238 E239 V243 N244 D245 Y246 M247 R248 G249 N255 N256 Y257 L258 I259 S260 N261 G262 P263

V264 N265 N266 P267 N268 P269 E270 L271 D272 S273 K274 I275 L278 E279 N280 R284 I287 R295 T305 S308 D312 N318 F322 W326 L331 I337 S338 T339 M346 F349 R352 L356 R357 I358 T359 I360 L369 S370 A371 I372 P373 G374 Y375

A376 G377 E378 F389 D395 T401 V407 P413 GLU ASN TYR PHE GLN GLY LEU HIS HIS HIS HIS HIS HIS

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	100912	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50.0	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.510	Depositor
Minimum map value	-0.716	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.037	Depositor
Recommended contour level	0.195	Depositor
Map size ($\text{\AA}$)	380.16, 380.16, 380.16	wwPDB
Map dimensions	432, 432, 432	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.88, 0.88, 0.88	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.19	0/4001	0.38	1/5423 (0.0%)
1	B	0.18	0/4001	0.34	1/5423 (0.0%)
1	C	0.19	0/3933	0.34	1/5332 (0.0%)
1	D	0.21	0/3966	0.37	1/5378 (0.0%)
1	E	0.19	0/4001	0.34	1/5423 (0.0%)
1	F	0.20	0/3933	0.36	1/5332 (0.0%)
1	G	0.19	0/4001	0.35	1/5423 (0.0%)
2	H	0.25	0/3269	0.41	0/4414
All	All	0.20	0/31105	0.36	7/42148 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	281	PHE	CA-CB-CG	6.55	120.35	113.80
1	G	281	PHE	CA-CB-CG	6.50	120.30	113.80
1	F	281	PHE	CA-CB-CG	6.34	120.14	113.80
1	C	281	PHE	CA-CB-CG	6.28	120.08	113.80
1	E	281	PHE	CA-CB-CG	6.11	119.91	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3935	0	3835	45	0
1	B	3935	0	3835	43	0
1	C	3867	0	3766	52	0
1	D	3898	0	3786	60	0
1	E	3935	0	3835	43	0
1	F	3867	0	3766	49	0
1	G	3935	0	3833	51	0
2	H	3200	0	3179	60	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
3	D	3	0	0	0	0
3	E	3	0	0	0	0
3	F	3	0	0	0	0
3	G	3	0	0	0	0
All	All	30593	0	29835	376	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 376 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:19:TRP:HE3	2:H:21:ARG:NH2	1.54	1.04
2:H:19:TRP:HZ2	2:H:142:TYR:HH	1.08	1.02
2:H:19:TRP:HE3	2:H:21:ARG:HH21	0.89	0.85
2:H:19:TRP:CE3	2:H:21:ARG:NH2	2.42	0.84
1:G:670:VAL:HG22	1:G:710:GLU:HG2	1.63	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	500/675 (74%)	470 (94%)	30 (6%)	0	100	100
1	B	500/675 (74%)	481 (96%)	19 (4%)	0	100	100
1	C	490/675 (73%)	462 (94%)	28 (6%)	0	100	100
1	D	493/675 (73%)	468 (95%)	25 (5%)	0	100	100
1	E	500/675 (74%)	474 (95%)	26 (5%)	0	100	100
1	F	490/675 (73%)	470 (96%)	20 (4%)	0	100	100
1	G	500/675 (74%)	472 (94%)	28 (6%)	0	100	100
2	H	393/428 (92%)	382 (97%)	11 (3%)	0	100	100
All	All	3866/5153 (75%)	3679 (95%)	187 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	446/601 (74%)	441 (99%)	5 (1%)	65	87
1	B	446/601 (74%)	445 (100%)	1 (0%)	87	96
1	C	438/601 (73%)	436 (100%)	2 (0%)	81	93
1	D	441/601 (73%)	440 (100%)	1 (0%)	87	96
1	E	446/601 (74%)	436 (98%)	10 (2%)	45	78
1	F	438/601 (73%)	435 (99%)	3 (1%)	76	91
1	G	446/601 (74%)	441 (99%)	5 (1%)	65	87
2	H	355/385 (92%)	344 (97%)	11 (3%)	35	70
All	All	3456/4592 (75%)	3418 (99%)	38 (1%)	68	87

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	22	LYS
2	H	173	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	25	GLU
2	H	30	LYS
2	H	176	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	567	ASN
2	H	229	ASN
1	E	376	ASN
1	G	634	ASN
1	E	329	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 21 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-34137. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 216




Y Index: 216




Z Index: 216


6.2.2 Raw map



X Index: 216



Y Index: 216




Z Index: 216


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 199




Y Index: 194




Z Index: 192


6.3.2 Raw map



X Index: 199



Y Index: 222




Z Index: 231


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.195. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 201 nm³; this corresponds to an approximate mass of 182 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.357 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.357 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.80	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.63	7.61	3.69

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.63 differs from the reported value 2.8 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-34137 and PDB model 7YVS. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.195 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.195).

9.4 Atom inclusion [i](#)



At the recommended contour level, 91% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.195) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8490	0.5490
A	0.8450	0.5540
B	0.8610	0.5590
C	0.8600	0.5590
D	0.8620	0.5600
E	0.8560	0.5540
F	0.8580	0.5540
G	0.8550	0.5570
H	0.7850	0.4860

1.0

0.0

<0.0