



wwPDB EM Validation Summary Report ⓘ

Mar 29, 2026 – 06:57 PM UTC

PDB ID : 8YVP / pdb_00008yvp
EMDB ID : EMD-39612
Title : canine immunoproteasome 20S subunit in complex with compound 1
Authors : Kashima, A.; Arai, Y.
Deposited on : 2024-03-29
Resolution : 2.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	7115 (2.00 - 3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	234	
1	E	234	
2	C	205	
2	D	205	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	P	234	
3	b	234	
4	J	255	
4	Q	255	
5	K	246	
5	R	246	
6	A	205	
6	F	205	
7	S	240	
7	X	240	
8	U	205	
8	Y	205	
9	W	264	
9	a	264	
10	H	241	
10	M	241	
11	T	201	
11	V	201	
12	I	248	
12	N	248	
13	G	263	
13	L	263	
14	O	261	
14	Z	261	

2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 48042 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	B	220	Total	C	N	O	S	0	0
			1656	1044	282	318	12		
1	E	220	Total	C	N	O	S	0	0
			1656	1044	282	318	12		

- Molecule 2 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	200	Total	C	N	O	S	0	0
			1551	980	270	292	9		
2	D	200	Total	C	N	O	S	0	0
			1551	980	270	292	9		

- Molecule 3 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	P	228	Total	C	N	O	S	0	0
			1781	1138	302	335	6		
3	b	228	Total	C	N	O	S	0	0
			1781	1138	302	335	6		

- Molecule 4 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	Q	235	Total	C	N	O	S	0	0
			1840	1168	315	346	11		
4	J	235	Total	C	N	O	S	0	0
			1840	1168	315	346	11		

- Molecule 5 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	R	236	Total	C	N	O	S	0	0
			1831	1163	304	351	13		
5	K	236	Total	C	N	O	S	0	0
			1831	1163	304	351	13		

- Molecule 6 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	202	Total	C	N	O	S	0	0
			1519	952	259	296	12		
6	A	202	Total	C	N	O	S	0	0
			1519	952	259	296	12		

- Molecule 7 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	S	212	Total	C	N	O	S	0	0
			1643	1041	280	312	10		
7	X	212	Total	C	N	O	S	0	0
			1643	1041	280	312	10		

- Molecule 8 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	Y	204	Total	C	N	O	S	0	0
			1592	1013	265	295	19		
8	U	204	Total	C	N	O	S	0	0
			1592	1013	265	295	19		

- Molecule 9 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	a	214	Total	C	N	O	S	0	0
			1671	1054	289	316	12		
9	W	214	Total	C	N	O	S	0	0
			1671	1054	289	316	12		

- Molecule 10 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	H	231	Total	C	N	O	S	0	0
			1761	1106	292	352	11		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
10	M	231	Total	C	N	O	S	0	0
			1761	1106	292	352	11		

- Molecule 11 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	V	196	Total	C	N	O	S	0	0
			1570	1006	267	288	9		
11	T	196	Total	C	N	O	S	0	0
			1570	1006	267	288	9		

- Molecule 12 is a protein called Proteasome subunit alpha type-7.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	I	232	Total	C	N	O	S	0	0
			1819	1146	322	346	5		
12	N	232	Total	C	N	O	S	0	0
			1819	1146	322	346	5		


- Molecule 13 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	L	234	Total	C	N	O	S	0	0
			1836	1150	328	347	11		
13	G	234	Total	C	N	O	S	0	0
			1836	1150	328	347	11		

- Molecule 14 is a protein called Proteasome subunit alpha type-4.


Mol	Chain	Residues	Atoms					AltConf	Trace
14	O	239	Total	C	N	O	S	0	0
			1881	1191	321	359	10		
14	Z	239	Total	C	N	O	S	0	0
			1881	1191	321	359	10		

- Molecule 15 is [(R)-cyclohexyl-[(1-cyclohexyl-1,2,3-triazol-4-yl)carbonylamino]methyl]boronic acid (CCD ID: A1L0D) (formula: C₁₆H₂₇BN₄O₃).



Mol	Chain	Residues	Atoms					AltConf
15	C	1	Total	B	C	N	O	0
			23	1	16	4	2	
15	C	1	Total	B	C	N	O	0
			23	1	16	4	2	
15	F	1	Total	B	C	N	O	0
			23	1	16	4	2	
15	D	1	Total	B	C	N	O	0
			23	1	16	4	2	
15	D	1	Total	B	C	N	O	0
			23	1	16	4	2	
15	A	1	Total	B	C	N	O	0
			23	1	16	4	2	

- Molecule 16 is THREONINE (CCD ID: THR) (formula: $C_4H_9NO_3$).




Mol	Chain	Residues	Atoms		AltConf
16	C	1	Total	O	0
			1	1	
16	D	1	Total	O	0
			1	1	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Proteasome subunit beta type-7




• Molecule 1: Proteasome subunit beta type-7




• Molecule 2: Proteasome subunit beta type-5




- Molecule 2: Proteasome subunit beta type-5




- Molecule 3: Proteasome subunit alpha type-2



- Molecule 3: Proteasome subunit alpha type-2




- Molecule 4: Proteasome subunit alpha type-3









• Molecule 4: Proteasome subunit alpha type-3




• Molecule 5: Proteasome subunit alpha type-6




• Molecule 5: Proteasome subunit alpha type-6




- Molecule 6: Proteasome subunit beta type-6




- Molecule 6: Proteasome subunit beta type-6



- Molecule 7: Proteasome subunit beta type-1




- Molecule 7: Proteasome subunit beta type-1




- Molecule 8: Proteasome subunit beta type-3

Chain Y:  56% 41% .




• Molecule 8: Proteasome subunit beta type-3

Chain U:  59% 38% .




• Molecule 9: Proteasome subunit beta type-4


Chain a:  47% 29% 5% 19%




• Molecule 9: Proteasome subunit beta type-4

Chain W:  48% 28% 5% 19%







• Molecule 10: Proteasome subunit alpha type-5




• Molecule 10: Proteasome subunit alpha type-5




• Molecule 11: Proteasome subunit beta type-2




- Molecule 11: Proteasome subunit beta type-2




- Molecule 12: Proteasome subunit alpha type-7



- Molecule 12: Proteasome subunit alpha type-7



- Molecule 13: Proteasome subunit alpha type-1



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	118136	Depositor
Resolution determination method	Not provided	
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	48.20	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1600	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.196	Depositor
Minimum map value	-0.107	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.008	Depositor
Recommended contour level	0.0282	Depositor
Map size (Å)	292.16095, 292.16095, 292.16095	wwPDB
Map dimensions	254, 254, 254	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.15024, 1.15024, 1.15024	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1L0D

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.37	0/1683	0.76	1/2276 (0.0%)
1	E	0.37	0/1683	0.76	1/2276 (0.0%)
2	C	0.39	0/1582	0.74	1/2137 (0.0%)
2	D	0.39	0/1582	0.74	1/2137 (0.0%)
3	P	0.33	0/1820	0.78	4/2466 (0.2%)
3	b	0.33	0/1820	0.78	4/2466 (0.2%)
4	J	0.32	0/1875	0.77	1/2524 (0.0%)
4	Q	0.32	0/1875	0.78	1/2524 (0.0%)
5	K	0.32	0/1863	0.78	1/2518 (0.0%)
5	R	0.32	0/1863	0.78	1/2518 (0.0%)
6	A	0.69	0/1546	1.00	2/2094 (0.1%)
6	F	0.69	0/1546	1.00	2/2094 (0.1%)
7	S	0.36	0/1674	0.79	3/2257 (0.1%)
7	X	0.36	0/1674	0.79	3/2257 (0.1%)
8	U	0.37	0/1621	0.75	0/2185
8	Y	0.37	0/1621	0.75	0/2185
9	W	0.37	0/1704	0.75	0/2306
9	a	0.37	0/1704	0.75	0/2306
10	H	0.32	0/1788	0.76	4/2415 (0.2%)
10	M	0.32	0/1788	0.76	4/2415 (0.2%)
11	T	0.37	0/1602	0.78	3/2167 (0.1%)
11	V	0.37	0/1602	0.78	3/2167 (0.1%)
12	I	0.33	0/1846	0.76	4/2495 (0.2%)
12	N	0.33	0/1846	0.76	4/2495 (0.2%)
13	G	0.32	0/1870	0.81	2/2528 (0.1%)
13	L	0.32	0/1870	0.81	2/2528 (0.1%)
14	O	0.36	0/1910	0.81	1/2573 (0.0%)
14	Z	0.37	0/1910	0.81	1/2573 (0.0%)
All	All	0.38	0/48768	0.79	54/65882 (0.1%)

There are no bond length outliers.

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	X	102	PHE	CA-C-N	8.72	129.25	119.92
7	X	102	PHE	C-N-CA	8.72	129.25	119.92
7	S	102	PHE	CA-C-N	8.68	129.21	119.92
7	S	102	PHE	C-N-CA	8.68	129.21	119.92
2	C	1	THR	O-C-N	8.59	136.75	123.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1656	0	1685	67	0
1	E	1656	0	1685	66	0
2	C	1551	0	1520	45	0
2	D	1551	0	1520	45	0
3	P	1781	0	1771	115	0
3	b	1781	0	1771	117	0
4	J	1840	0	1827	93	0
4	Q	1840	0	1827	94	0
5	K	1831	0	1841	111	0
5	R	1831	0	1841	104	0
6	A	1519	0	1487	56	0
6	F	1519	0	1487	55	0
7	S	1643	0	1638	46	0
7	X	1643	0	1638	44	0
8	U	1592	0	1612	57	0
8	Y	1592	0	1612	60	0
9	W	1671	0	1648	56	0
9	a	1671	0	1648	59	0
10	H	1761	0	1749	97	0
10	M	1761	0	1749	101	0
11	T	1570	0	1573	41	0
11	V	1570	0	1573	40	0
12	I	1819	0	1845	90	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
12	N	1819	0	1845	92	0
13	G	1836	0	1822	90	0
13	L	1836	0	1822	89	0
14	O	1881	0	1896	113	0
14	Z	1881	0	1896	105	0
15	A	23	0	0	0	0
15	C	46	0	0	1	0
15	D	46	0	0	1	0
15	F	23	0	0	0	0
16	C	1	0	0	1	0
16	D	1	0	0	1	0
All	All	48042	0	47828	1936	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 1936 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
14:Z:174:MET:HE1	14:Z:199:LYS:CB	1.54	1.37
14:O:174:MET:HE1	14:O:199:LYS:CB	1.54	1.36
12:I:41:VAL:HG11	12:I:134:VAL:CG1	1.57	1.32
12:N:41:VAL:HG11	12:N:134:VAL:CG1	1.57	1.30
10:H:31:ILE:HD11	10:H:158:PRO:HG3	1.24	1.17

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	218/234 (93%)	210 (96%)	7 (3%)	1 (0%)	24 43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	218/234 (93%)	210 (96%)	7 (3%)	1 (0%)	24	43
2	C	198/205 (97%)	190 (96%)	8 (4%)	0	100	100
2	D	198/205 (97%)	190 (96%)	8 (4%)	0	100	100
3	P	226/234 (97%)	210 (93%)	12 (5%)	4 (2%)	6	12
3	b	226/234 (97%)	210 (93%)	12 (5%)	4 (2%)	6	12
4	J	233/255 (91%)	220 (94%)	13 (6%)	0	100	100
4	Q	233/255 (91%)	219 (94%)	14 (6%)	0	100	100
5	K	232/246 (94%)	219 (94%)	12 (5%)	1 (0%)	30	49
5	R	232/246 (94%)	219 (94%)	12 (5%)	1 (0%)	30	49
6	A	200/205 (98%)	190 (95%)	9 (4%)	1 (0%)	24	43
6	F	200/205 (98%)	190 (95%)	9 (4%)	1 (0%)	24	43
7	S	210/240 (88%)	197 (94%)	11 (5%)	2 (1%)	12	24
7	X	210/240 (88%)	197 (94%)	11 (5%)	2 (1%)	12	24
8	U	202/205 (98%)	190 (94%)	11 (5%)	1 (0%)	24	43
8	Y	202/205 (98%)	190 (94%)	11 (5%)	1 (0%)	24	43
9	W	212/264 (80%)	201 (95%)	9 (4%)	2 (1%)	14	27
9	a	212/264 (80%)	201 (95%)	9 (4%)	2 (1%)	14	27
10	H	229/241 (95%)	214 (93%)	15 (7%)	0	100	100
10	M	229/241 (95%)	214 (93%)	15 (7%)	0	100	100
11	T	194/201 (96%)	189 (97%)	4 (2%)	1 (0%)	24	43
11	V	194/201 (96%)	189 (97%)	4 (2%)	1 (0%)	24	43
12	I	230/248 (93%)	215 (94%)	14 (6%)	1 (0%)	30	49
12	N	230/248 (93%)	215 (94%)	14 (6%)	1 (0%)	30	49
13	G	232/263 (88%)	217 (94%)	14 (6%)	1 (0%)	30	49
13	L	232/263 (88%)	217 (94%)	14 (6%)	1 (0%)	30	49
14	O	235/261 (90%)	228 (97%)	6 (3%)	1 (0%)	30	49
14	Z	235/261 (90%)	228 (97%)	6 (3%)	1 (0%)	30	49
All	All	6102/6604 (92%)	5779 (95%)	291 (5%)	32 (0%)	26	43

5 of 32 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
8	Y	78	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
14	O	59	VAL
14	Z	59	VAL
8	U	78	GLY
1	B	95	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	181/195 (93%)	145 (80%)	36 (20%)	1	2
1	E	181/195 (93%)	145 (80%)	36 (20%)	1	2
2	C	156/161 (97%)	128 (82%)	28 (18%)	2	3
2	D	156/161 (97%)	128 (82%)	28 (18%)	2	3
3	P	187/191 (98%)	148 (79%)	39 (21%)	1	2
3	b	187/191 (98%)	148 (79%)	39 (21%)	1	2
4	J	192/211 (91%)	153 (80%)	39 (20%)	1	2
4	Q	192/211 (91%)	153 (80%)	39 (20%)	1	2
5	K	201/210 (96%)	160 (80%)	41 (20%)	1	2
5	R	201/210 (96%)	160 (80%)	41 (20%)	1	2
6	A	159/162 (98%)	140 (88%)	19 (12%)	5	11
6	F	159/162 (98%)	140 (88%)	19 (12%)	5	11
7	S	177/198 (89%)	144 (81%)	33 (19%)	1	3
7	X	177/198 (89%)	144 (81%)	33 (19%)	1	3
8	U	174/175 (99%)	151 (87%)	23 (13%)	4	8
8	Y	174/175 (99%)	151 (87%)	23 (13%)	4	8
9	W	176/212 (83%)	144 (82%)	32 (18%)	2	3
9	a	176/212 (83%)	144 (82%)	32 (18%)	2	3
10	H	193/203 (95%)	154 (80%)	39 (20%)	1	2
10	M	193/203 (95%)	154 (80%)	39 (20%)	1	2

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
11	T	166/171 (97%)	149 (90%)	17 (10%)	7	15
11	V	166/171 (97%)	149 (90%)	17 (10%)	7	15
12	I	195/211 (92%)	146 (75%)	49 (25%)	0	1
12	N	195/211 (92%)	146 (75%)	49 (25%)	0	1
13	G	200/224 (89%)	158 (79%)	42 (21%)	1	2
13	L	200/224 (89%)	158 (79%)	42 (21%)	1	2
14	O	201/221 (91%)	153 (76%)	48 (24%)	1	1
14	Z	201/221 (91%)	153 (76%)	48 (24%)	1	1
All	All	5116/5490 (93%)	4146 (81%)	970 (19%)	3	3

5 of 970 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
10	H	231	LYS
12	N	100	ASP
11	T	145	ARG
12	N	41	VAL
9	W	51	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 119 such sidechains are listed below:

Mol	Chain	Res	Type
3	b	139	ASN
12	N	205	ASN
12	I	23	GLN
12	N	200	GLN
9	W	188	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are modelled with single atom - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
15	A1L0D	D	401	16	22,25,26	0.46	0	29,33,35	1.01	2 (6%)
15	A1L0D	A	401	6	22,25,26	0.40	0	29,33,35	0.81	1 (3%)
15	A1L0D	F	401	6	22,25,26	0.42	0	29,33,35	0.81	1 (3%)
15	A1L0D	C	401	2	22,25,26	0.45	0	29,33,35	0.86	1 (3%)
15	A1L0D	C	402	16	22,25,26	0.46	0	29,33,35	1.02	2 (6%)
15	A1L0D	D	403	2	22,25,26	0.45	0	29,33,35	0.86	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	A1L0D	D	401	16	-	8/13/34/36	0/3/3/3
15	A1L0D	A	401	6	-	4/13/34/36	0/3/3/3
15	A1L0D	F	401	6	-	4/13/34/36	0/3/3/3
15	A1L0D	C	401	2	-	4/13/34/36	0/3/3/3
15	A1L0D	C	402	16	-	8/13/34/36	0/3/3/3
15	A1L0D	D	403	2	-	4/13/34/36	0/3/3/3

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	F	401	A1L0D	C16-C4-C3	-2.70	109.50	112.16
15	A	401	A1L0D	C16-C4-C3	-2.69	109.51	112.16
15	C	402	A1L0D	C6-N4-N3	-2.49	117.56	120.94
15	D	401	A1L0D	C6-N4-N3	-2.49	117.56	120.94
15	D	401	A1L0D	C7-C6-N4	2.36	114.19	110.95

There are no chirality outliers.

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
15	C	401	A1L0D	C5-C1-C2-N1
15	C	401	A1L0D	N2-C1-C2-N1
15	C	401	A1L0D	C5-C1-C2-O3
15	C	401	A1L0D	N2-C1-C2-O3
15	C	402	A1L0D	N1-C3-C4-C12


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
15	D	401	A1L0D	1	0
15	C	402	A1L0D	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1L0D D 401




Bond lengths



Bond angles




Torsions




Rings


Ligand A1L0D A 401




Bond lengths




Bond angles




Torsions



Rings





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-39612. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 127




Y Index: 127




Z Index: 127


6.2.2 Raw map



X Index: 127



Y Index: 127




Z Index: 127


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 110




Y Index: 110




Z Index: 148


6.3.2 Raw map



X Index: 144



Y Index: 144




Z Index: 148


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0282. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_39612_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 254 nm³; this corresponds to an approximate mass of 230 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.400 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.400 Å⁻¹

8.2 Resolution estimates


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	-	-	-
Author-provided FSC curve	2.52	3.00	2.62
Unmasked-calculated*	2.97	3.54	3.05

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-39612 and PDB model 8YVP. Per-residue inclusion information can be found in [section 3](#) on [page 9](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0282 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0282).

9.4 Atom inclusion [i](#)



At the recommended contour level, 94% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0282) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8710	 0.6400
A	 0.9170	 0.6600
B	 0.9050	 0.6530
C	 0.9260	 0.6600
D	 0.9240	 0.6600
E	 0.9070	 0.6550
F	 0.9180	 0.6610
G	 0.8460	 0.6250
H	 0.8320	 0.6270
I	 0.8230	 0.6140
J	 0.8070	 0.6250
K	 0.7930	 0.6190
L	 0.8460	 0.6270
M	 0.8280	 0.6280
N	 0.8250	 0.6150
O	 0.8330	 0.6150
P	 0.8430	 0.6280
Q	 0.8070	 0.6220
R	 0.7910	 0.6150
S	 0.9210	 0.6570
T	 0.9290	 0.6680
U	 0.9360	 0.6690
V	 0.9270	 0.6670
W	 0.9330	 0.6630
X	 0.9210	 0.6560
Y	 0.9370	 0.6700
Z	 0.8320	 0.6170
a	 0.9330	 0.6620
b	 0.8410	 0.6300

