wwPDB X-ray Structure Validation Summary Report (i)

Nov 10, 2026 – 12:20 PM EST

PDB ID	:	9YVQ / pdb_00009yvq
Title	:	human Siderocalin bound to 343HOPO-Tm
Authors	:	Rupert, P.B.; Strong, R.K.
Deposited on	:	2025-10-23
Resolution	:	2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	6% 96%
1	B	178	6% 97%
1	C	178	7% 92% 5%
1	D	178	4% 91% 5%
1	E	178	10% 95%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	178	5% 94%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	A1DTM	A	301	X	-	-	-
2	A1DTM	B	301	X	-	-	-
2	A1DTM	C	301	X	-	-	-
2	A1DTM	D	301	X	-	-	-
2	A1DTM	E	301	X	-	-	-
2	A1DTM	F	301	X	-	-	-

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 9142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Neutrophil gelatinase-associated lipocalin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	173	Total	C	N	O	S	0	2	0
			1401	910	232	255	4			
1	B	175	Total	C	N	O	S	0	4	0
			1422	923	236	259	4			
1	C	173	Total	C	N	O	S	0	2	0
			1380	899	229	248	4			
1	D	171	Total	C	N	O	S	0	2	0
			1382	898	230	250	4			
1	E	175	Total	C	N	O	S	0	1	0
			1399	908	231	256	4			
1	F	174	Total	C	N	O	S	0	3	0
			1417	920	236	257	4			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	87	SER	CYS	engineered mutation	UNP P80188
B	87	SER	CYS	engineered mutation	UNP P80188
C	87	SER	CYS	engineered mutation	UNP P80188
D	87	SER	CYS	engineered mutation	UNP P80188
E	87	SER	CYS	engineered mutation	UNP P80188
F	87	SER	CYS	engineered mutation	UNP P80188

- Molecule 2 is [(1R)-6-[(R)-(3-{[(1S)-1-(oxido-kappaO)-6-(oxo-kappaO)-1,6-dihydropyridine-2-carbonyl]amino}propyl){4-[(S)-[(1R)-1-(oxido-kappaO)-6-oxo-1,6-dihydropyridine-2-carbonyl](3-{[(1S)-1-(oxido-kappaO)-6-(oxo-kappaO)-1,6-dihydropyridine-2-carbonyl]amino}propyl)amino]butyl}carbamoyl]-2-(oxo-kappaO)pyridin-1(2H)-olato-kappaO]thulate(1-) (CCD ID: A1DTM) (formula: $C_{34}H_{34}N_8O_{12}Tm$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	Tm	0	0
			55	34	8	12	1		
2	B	1	Total	C	N	O	Tm	0	0
			55	34	8	12	1		
2	C	1	Total	C	N	O	Tm	0	0
			55	34	8	12	1		
2	D	1	Total	C	N	O	Tm	0	0
			55	34	8	12	1		
2	E	1	Total	C	N	O	Tm	0	0
			55	34	8	12	1		
2	F	1	Total	C	N	O	Tm	0	0
			55	34	8	12	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	1	Total O S 5 4 1	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0
5	B	1	Total Cl 1 1	0	0
5	C	1	Total Cl 1 1	0	0
5	D	1	Total Cl 1 1	0	0
5	E	1	Total Cl 1 1	0	0
5	F	1	Total Cl 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	65	Total O 65 65	0	0
6	B	67	Total O 67 67	0	0
6	C	47	Total O 47 47	0	0
6	D	56	Total O 56 56	0	0
6	E	33	Total O 33 33	0	0
6	F	50	Total O 50 50	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Neutrophil gelatinase-associated lipocalin


Chain A:






• Molecule 1: Neutrophil gelatinase-associated lipocalin

Chain B:






• Molecule 1: Neutrophil gelatinase-associated lipocalin


Chain C:





• Molecule 1: Neutrophil gelatinase-associated lipocalin


Chain D:






• Molecule 1: Neutrophil gelatinase-associated lipocalin


Chain E:





• Molecule 1: Neutrophil gelatinase-associated lipocalin

Chain F:


5%
94%
GLN
ASP
SER
T4
S5
S14
Q26
W31
Y32
V33
D45
K46
D47
P48
Y52
I80
Y138
L148
N151
D177
GLY

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	108.16Å 115.01Å 121.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.79 – 2.10 41.79 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.6 (41.79-2.10) 97.8 (41.79-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	4.59 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.233 , 0.265 0.237 , 0.268	Depositor DCC
R_{free} test set	4438 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å^2)	22.9	Xtriage
Anisotropy	0.125	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 33.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	9142	wwPDB-VP
Average B, all atoms (Å^2)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.12% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4, GOL, A1DTM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.48	0/1443	0.78	0/1956
1	B	0.47	0/1471	0.78	0/1997
1	C	0.49	0/1423	0.77	0/1935
1	D	0.48	0/1421	0.77	0/1929
1	E	0.47	0/1436	0.77	0/1952
1	F	0.47	0/1460	0.78	0/1981
All	All	0.48	0/8654	0.78	0/11750

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1401	0	1388	1	0
1	B	1422	0	1406	1	0
1	C	1380	0	1352	6	0
1	D	1382	0	1358	4	0
1	E	1399	0	1365	5	0
1	F	1417	0	1403	4	0
2	A	55	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	55	0	0	0	0
2	C	55	0	0	0	0
2	D	55	0	0	0	0
2	E	55	0	0	1	0
2	F	55	0	0	0	0
3	A	5	0	0	0	0
3	B	15	0	0	0	0
3	C	20	0	0	0	0
3	D	20	0	0	0	0
3	E	15	0	0	0	0
4	A	12	0	16	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
6	A	65	0	0	0	0
6	B	67	0	0	0	0
6	C	47	0	0	0	0
6	D	56	0	0	0	0
6	E	33	0	0	1	0
6	F	50	0	0	1	0
All	All	9142	0	8288	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 21 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:26:GLN:NE2	6:F:401:HOH:O	2.35	0.59
1:B:59:LYS:HE3	1:B:63:SER:OG	2.02	0.59
1:C:144:LEU:HD13	1:C:148:LEU:HD23	1.91	0.52
1:C:87:SER:O	1:E:105:SER:HB2	2.11	0.51
1:F:33:VAL:HG21	1:F:52:TYR:CE2	2.48	0.48

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	171/178 (96%)	167 (98%)	4 (2%)	0	100	100
1	B	177/178 (99%)	174 (98%)	3 (2%)	0	100	100
1	C	173/178 (97%)	170 (98%)	3 (2%)	0	100	100
1	D	169/178 (95%)	166 (98%)	3 (2%)	0	100	100
1	E	174/178 (98%)	170 (98%)	4 (2%)	0	100	100
1	F	175/178 (98%)	171 (98%)	4 (2%)	0	100	100
All	All	1039/1068 (97%)	1018 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/163 (96%)	156 (100%)	0	100	100
1	B	158/163 (97%)	157 (99%)	1 (1%)	78	86
1	C	150/163 (92%)	148 (99%)	2 (1%)	61	69
1	D	152/163 (93%)	151 (99%)	1 (1%)	76	83
1	E	153/163 (94%)	153 (100%)	0	100	100
1	F	157/163 (96%)	157 (100%)	0	100	100
All	All	926/978 (95%)	922 (100%)	4 (0%)	84	89

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	97	ILE
1	C	20	GLN
1	C	97	ILE
1	D	49	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	88	GLN
1	E	49	GLN
1	C	20	GLN
1	C	49	GLN
1	C	88	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 29 ligands modelled in this entry, 6 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
3	SO4	B	303	-	4,4,4	0.34	0	6,6,6	0.08	0
4	GOL	A	303	-	5,5,5	0.12	0	5,5,5	0.42	0
2	A1DTM	B	301	-	50,64,64	1.29	8 (16%)	68,106,106	2.36	13 (19%)
3	SO4	E	303	-	4,4,4	0.33	0	6,6,6	0.08	0
3	SO4	D	302	-	4,4,4	0.35	0	6,6,6	0.05	0
2	A1DTM	F	301	-	50,64,64	1.24	8 (16%)	68,106,106	2.38	12 (17%)
2	A1DTM	E	301	-	50,64,64	1.28	8 (16%)	68,106,106	2.43	13 (19%)
2	A1DTM	A	301	-	50,64,64	1.28	8 (16%)	68,106,106	2.44	12 (17%)
3	SO4	D	305	-	4,4,4	0.34	0	6,6,6	0.08	0
3	SO4	D	304	-	4,4,4	0.34	0	6,6,6	0.07	0
3	SO4	E	304	-	4,4,4	0.34	0	6,6,6	0.08	0
3	SO4	E	302	-	4,4,4	0.34	0	6,6,6	0.07	0
3	SO4	B	302	-	4,4,4	0.34	0	6,6,6	0.08	0
3	SO4	C	305	-	4,4,4	0.34	0	6,6,6	0.08	0
3	SO4	D	303	-	4,4,4	0.34	0	6,6,6	0.07	0
4	GOL	A	304	-	5,5,5	0.10	0	5,5,5	0.29	0
3	SO4	A	302	-	4,4,4	0.33	0	6,6,6	0.08	0
3	SO4	C	304	-	4,4,4	0.34	0	6,6,6	0.08	0
3	SO4	B	304	-	4,4,4	0.34	0	6,6,6	0.07	0
3	SO4	C	302	-	4,4,4	0.35	0	6,6,6	0.07	0
2	A1DTM	D	301	-	50,64,64	1.29	8 (16%)	68,106,106	2.42	15 (22%)
3	SO4	C	303	-	4,4,4	0.34	0	6,6,6	0.08	0
2	A1DTM	C	301	-	50,64,64	1.31	8 (16%)	68,106,106	2.42	12 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1DTM	F	301	-	5/5/7/9	18/45/101/101	-
2	A1DTM	E	301	-	5/5/7/9	13/45/101/101	-
2	A1DTM	D	301	-	5/5/7/9	4/45/101/101	-
4	GOL	A	303	-	-	4/4/4/4	-
2	A1DTM	B	301	-	5/5/7/9	13/45/101/101	-
2	A1DTM	A	301	-	5/5/7/9	10/45/101/101	-
4	GOL	A	304	-	-	1/4/4/4	-
2	A1DTM	C	301	-	5/5/7/9	19/45/101/101	-

The worst 5 of 48 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	301	A1DTM	C36-N35	3.60	1.41	1.39
2	B	301	A1DTM	C36-N35	3.54	1.41	1.39
2	D	301	A1DTM	C36-N35	3.41	1.41	1.39
2	F	301	A1DTM	C26-N27	3.39	1.42	1.38
2	A	301	A1DTM	C26-N27	3.34	1.41	1.38

The worst 5 of 77 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	C	301	A1DTM	O49-C26-C25	-9.32	119.61	126.29
2	E	301	A1DTM	O9-C4-C5	-8.96	119.86	126.29
2	B	301	A1DTM	O9-C4-C5	-8.87	119.93	126.29
2	A	301	A1DTM	O9-C4-C5	-8.85	119.94	126.29
2	D	301	A1DTM	O9-C4-C5	-8.72	120.03	126.29

5 of 30 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	301	A1DTM	N27
2	A	301	A1DTM	N15
2	A	301	A1DTM	N21
2	A	301	A1DTM	N45
2	A	301	A1DTM	N3

5 of 82 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	A1DTM	O54-C16-N15-C14
2	A	301	A1DTM	O54-C16-N15-C17
2	A	301	A1DTM	C22-C16-N15-C14
2	A	301	A1DTM	C22-C16-N15-C17
2	B	301	A1DTM	O54-C16-N15-C14

There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	301	A1DTM	1	0
2	A	301	A1DTM	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1DTM B 301




Ligand A1DTM F 301




Bond lengths



Bond angles




Torsions




Rings


Ligand A1DTM E 301




Bond lengths



Bond angles




Torsions




Rings


Ligand A1DTM A 301




Bond lengths



Bond angles




Torsions




Rings


Ligand A1DTM D 301




Bond lengths



Bond angles




Torsions



Rings

Ligand A1DTM C 301



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	173/178 (97%)	0.47	11 (6%)	25	27	16, 29, 44, 62	2 (1%)
1	B	175/178 (98%)	0.52	11 (6%)	26	27	16, 29, 48, 73	4 (2%)
1	C	173/178 (97%)	0.67	12 (6%)	23	24	18, 33, 50, 83	2 (1%)
1	D	171/178 (96%)	0.49	8 (4%)	36	38	16, 31, 51, 62	2 (1%)
1	E	175/178 (98%)	0.98	17 (9%)	13	14	24, 40, 59, 75	2 (1%)
1	F	174/178 (97%)	0.73	9 (5%)	33	35	20, 34, 51, 76	4 (2%)
All	All	1041/1068 (97%)	0.64	68 (6%)	25	26	16, 33, 53, 83	16 (1%)

The worst 5 of 68 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	45	ASP	10.9
1	F	45	ASP	7.8
1	A	44	GLU	4.7
1	F	4	THR	4.4
1	E	4	THR	4.3

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	SO4	B	303	5/5	0.59	0.19	102,103,105,105	0
3	SO4	B	304	5/5	0.60	0.14	82,84,85,86	0
3	SO4	E	302	5/5	0.63	0.15	106,107,108,108	0
3	SO4	C	304	5/5	0.65	0.16	87,87,87,88	0
3	SO4	E	303	5/5	0.67	0.14	82,83,87,87	0
3	SO4	D	305	5/5	0.69	0.12	81,82,84,85	0
3	SO4	D	303	5/5	0.69	0.13	85,85,87,88	0
3	SO4	D	304	5/5	0.69	0.14	86,87,88,88	0
3	SO4	C	305	5/5	0.70	0.13	88,88,89,93	0
3	SO4	E	304	5/5	0.70	0.15	92,94,94,95	0
3	SO4	C	302	5/5	0.71	0.14	67,69,70,70	0
4	GOL	A	303	6/6	0.71	0.19	52,52,52,53	0
3	SO4	A	302	5/5	0.74	0.13	75,76,77,79	0
4	GOL	A	304	6/6	0.75	0.19	50,52,53,55	0
3	SO4	B	302	5/5	0.85	0.10	61,62,64,64	0
3	SO4	C	303	5/5	0.86	0.10	64,65,66,66	0
3	SO4	D	302	5/5	0.88	0.14	57,59,59,61	0
2	A1DTM	C	301	55/55	0.95	0.17	46,69,80,81	1
5	CL	C	306	1/1	0.96	0.08	39,39,39,39	0
2	A1DTM	A	301	55/55	0.97	0.12	36,49,59,63	1
2	A1DTM	B	301	55/55	0.98	0.09	28,37,41,41	1
2	A1DTM	D	301	55/55	0.98	0.10	30,38,45,46	1
5	CL	A	305	1/1	0.98	0.07	33,33,33,33	0
5	CL	B	305	1/1	0.98	0.05	29,29,29,29	0
2	A1DTM	E	301	55/55	0.98	0.11	35,46,54,58	1
5	CL	E	305	1/1	0.98	0.07	36,36,36,36	0
5	CL	D	306	1/1	0.99	0.03	30,30,30,30	0
2	A1DTM	F	301	55/55	0.99	0.09	30,42,46,49	1
5	CL	F	302	1/1	0.99	0.05	32,32,32,32	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1DTM C 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around A1DTM A 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around A1DTM B 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)










Electron density around A1DTM D 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around A1DTM E 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around A1DTM F 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers

There are no such residues in this entry.