



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:28 AM UTC

PDB ID : 5YWS / pdb_00005yws
Title : Crystal structure of TREX1 in complex with a Y structured DNA
Authors : Hsiao, Y.Y.
Deposited on : 2017-11-30
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	6% 79% 5% 16%
1	B	276	5% 82% • 15%
2	C	22	14% 77% 23%
2	D	22	9% 91% 9%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8718 atoms, of which 3984 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Three-prime repair exonuclease 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	231	Total	C	H	N	O	S	0	0	0
			3549	1119	1774	314	333	9			
1	B	234	Total	C	H	N	O	S	0	0	0
			3584	1129	1790	318	338	9			

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-33	MET	-	expression tag	UNP Q91XB0
A	-32	GLY	-	expression tag	UNP Q91XB0
A	-31	SER	-	expression tag	UNP Q91XB0
A	-30	SER	-	expression tag	UNP Q91XB0
A	-29	HIS	-	expression tag	UNP Q91XB0
A	-28	HIS	-	expression tag	UNP Q91XB0
A	-27	HIS	-	expression tag	UNP Q91XB0
A	-26	HIS	-	expression tag	UNP Q91XB0
A	-25	HIS	-	expression tag	UNP Q91XB0
A	-24	HIS	-	expression tag	UNP Q91XB0
A	-23	SER	-	expression tag	UNP Q91XB0
A	-22	SER	-	expression tag	UNP Q91XB0
A	-21	GLY	-	expression tag	UNP Q91XB0
A	-20	LEU	-	expression tag	UNP Q91XB0
A	-19	VAL	-	expression tag	UNP Q91XB0
A	-18	PRO	-	expression tag	UNP Q91XB0
A	-17	ARG	-	expression tag	UNP Q91XB0
A	-16	GLY	-	expression tag	UNP Q91XB0
A	-15	SER	-	expression tag	UNP Q91XB0
A	-14	HIS	-	expression tag	UNP Q91XB0
A	-13	MET	-	expression tag	UNP Q91XB0
A	-12	ALA	-	expression tag	UNP Q91XB0
A	-11	SER	-	expression tag	UNP Q91XB0
A	-10	MET	-	expression tag	UNP Q91XB0
A	-9	THR	-	expression tag	UNP Q91XB0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	GLY	-	expression tag	UNP Q91XB0
A	-7	GLY	-	expression tag	UNP Q91XB0
A	-6	GLN	-	expression tag	UNP Q91XB0
A	-5	GLN	-	expression tag	UNP Q91XB0
A	-4	MET	-	expression tag	UNP Q91XB0
A	-3	GLY	-	expression tag	UNP Q91XB0
A	-2	ARG	-	expression tag	UNP Q91XB0
A	-1	GLY	-	expression tag	UNP Q91XB0
A	0	SER	-	expression tag	UNP Q91XB0
B	-33	MET	-	expression tag	UNP Q91XB0
B	-32	GLY	-	expression tag	UNP Q91XB0
B	-31	SER	-	expression tag	UNP Q91XB0
B	-30	SER	-	expression tag	UNP Q91XB0
B	-29	HIS	-	expression tag	UNP Q91XB0
B	-28	HIS	-	expression tag	UNP Q91XB0
B	-27	HIS	-	expression tag	UNP Q91XB0
B	-26	HIS	-	expression tag	UNP Q91XB0
B	-25	HIS	-	expression tag	UNP Q91XB0
B	-24	HIS	-	expression tag	UNP Q91XB0
B	-23	SER	-	expression tag	UNP Q91XB0
B	-22	SER	-	expression tag	UNP Q91XB0
B	-21	GLY	-	expression tag	UNP Q91XB0
B	-20	LEU	-	expression tag	UNP Q91XB0
B	-19	VAL	-	expression tag	UNP Q91XB0
B	-18	PRO	-	expression tag	UNP Q91XB0
B	-17	ARG	-	expression tag	UNP Q91XB0
B	-16	GLY	-	expression tag	UNP Q91XB0
B	-15	SER	-	expression tag	UNP Q91XB0
B	-14	HIS	-	expression tag	UNP Q91XB0
B	-13	MET	-	expression tag	UNP Q91XB0
B	-12	ALA	-	expression tag	UNP Q91XB0
B	-11	SER	-	expression tag	UNP Q91XB0
B	-10	MET	-	expression tag	UNP Q91XB0
B	-9	THR	-	expression tag	UNP Q91XB0
B	-8	GLY	-	expression tag	UNP Q91XB0
B	-7	GLY	-	expression tag	UNP Q91XB0
B	-6	GLN	-	expression tag	UNP Q91XB0
B	-5	GLN	-	expression tag	UNP Q91XB0
B	-4	MET	-	expression tag	UNP Q91XB0
B	-3	GLY	-	expression tag	UNP Q91XB0
B	-2	ARG	-	expression tag	UNP Q91XB0
B	-1	GLY	-	expression tag	UNP Q91XB0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	SER	-	expression tag	UNP Q91XB0

- Molecule 2 is a DNA chain called A STEM LOOP DNA WITH Y-STRUCTURAL TERMINAL.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	17	Total	C	H	N	O	P	0	0	0
			537	164	193	61	103	16			
2	D	20	Total	C	H	N	O	P	0	0	0
			631	193	227	71	121	19			

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		
3	B	2	Total	Mg	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	161	Total	O	0	0
			161	161		
4	B	179	Total	O	0	0
			179	179		
4	C	26	Total	O	0	0
			26	26		
4	D	47	Total	O	0	0
			47	47		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.53Å 96.60Å 101.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.33 – 2.00 29.33 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.1 (29.33-2.00) 97.0 (29.33-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.34 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.189 , 0.226 0.191 , 0.226	Depositor DCC
R_{free} test set	3013 reflections (7.51%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.433	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 39.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.015 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8718	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/1819	0.64	0/2482
1	B	0.32	0/1838	0.65	0/2507
2	C	0.21	0/383	0.44	0/587
2	D	0.21	0/450	0.45	0/690
All	All	0.30	0/4490	0.61	0/6266

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1775	1774	1774	8	0
1	B	1794	1790	1790	4	0
2	C	344	193	194	0	0
2	D	404	227	228	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	161	0	0	4	0
4	B	179	0	0	2	0
4	C	26	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	47	0	0	0	0
All	All	4734	3984	3986	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:LYS:NZ	4:A:402:HOH:O	2.24	0.68
1:A:41:ARG:NH1	1:A:45:GLU:OE2	2.34	0.61
1:A:140:ARG:NH1	4:A:405:HOH:O	2.36	0.57
1:B:213:GLN:NE2	4:B:405:HOH:O	2.37	0.56
1:A:41:ARG:NH2	1:A:220:ASP:OD2	2.45	0.50
1:B:131:PHE:HB2	1:B:132:PRO:HD3	1.94	0.49
1:B:137:GLU:OE1	1:B:140:ARG:NH2	2.46	0.49
1:A:172:GLY:N	4:A:413:HOH:O	2.46	0.49
1:A:169:SER:HB3	1:A:170:GLY:HA2	1.94	0.48
1:A:131:PHE:HB2	1:A:132:PRO:HD3	1.96	0.46
1:B:51:GLN:NE2	4:B:410:HOH:O	2.49	0.46
1:A:224:ARG:NH2	4:A:411:HOH:O	2.48	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/276 (83%)	225 (98%)	4 (2%)	0	100	100
1	B	232/276 (84%)	231 (100%)	1 (0%)	0	100	100
All	All	461/552 (84%)	456 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/232 (85%)	194 (98%)	4 (2%)	48	54
1	B	200/232 (86%)	199 (100%)	1 (0%)	81	87
All	All	398/464 (86%)	393 (99%)	5 (1%)	61	68

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	LEU
1	A	171	ASN
1	A	175	LYS
1	A	235	THR
1	B	215	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	HIS
1	A	195	HIS
1	B	164	GLN
1	B	171	ASN
1	B	195	HIS
1	B	213	GLN
1	B	222	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/276 (83%)	0.26	16 (6%) 23 21	19, 31, 72, 122	0
1	B	234/276 (84%)	0.16	15 (6%) 25 24	17, 28, 71, 125	0
2	C	17/22 (77%)	0.97	3 (17%) 4 3	27, 55, 100, 115	0
2	D	20/22 (90%)	0.58	2 (10%) 12 11	24, 45, 81, 96	0
All	All	502/596 (84%)	0.25	36 (7%) 21 20	17, 31, 78, 125	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	170	GLY	5.9
1	A	168	PRO	5.3
1	A	235	THR	4.4
1	B	168	PRO	4.4
1	A	49	ILE	4.2
1	B	50	SER	4.1
1	A	48	SER	3.8
1	A	169	SER	3.8
1	B	167	SER	3.7
1	B	169	SER	3.7
1	B	235	THR	3.4
1	B	53	HIS	3.2
1	B	170	GLY	3.1
1	A	167	SER	3.0
1	A	165	ALA	3.0
1	A	166	SER	2.9
2	D	14	DA	2.8
1	B	49	ILE	2.8
1	B	166	SER	2.8
1	B	171	ASN	2.7
1	B	51	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	8	DC	2.7
2	D	13	DT	2.7
1	B	172	GLY	2.5
1	A	83	GLU	2.5
2	C	9	DT	2.5
1	A	5	THR	2.5
1	B	128	ARG	2.2
1	A	148	ASP	2.2
1	A	50	SER	2.1
1	A	171	ASN	2.1
1	A	52	GLY	2.1
2	C	6	DC	2.1
1	B	234	GLY	2.1
1	A	46	ASN	2.1
1	B	140	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	B	302	1/1	0.91	0.33	29,29,29,29	0
3	MG	A	301	1/1	0.92	0.06	26,26,26,26	0
3	MG	A	302	1/1	0.94	0.16	31,31,31,31	0
3	MG	B	301	1/1	0.98	0.03	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.