



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:40 PM UTC

PDB ID : 7YWS / pdb_00007yws
Title : Modified oligopeptidase B from *S. proteomaculans* in intermediate conformation with 3 spermine molecules at 1.7 Å resolution
Authors : Petrenko, D.E.; Boyko, K.M.; Nikolaeva, A.Y.; Vlaskina, A.V.; Mikhailova, A.G.; Timofeev, V.I.; Rakitina, T.V.
Deposited on : 2022-02-14
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	677	5% 86% 12%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6128 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Oligopeptidase B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	677	5568	3535	936	1084	13	0	3	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	HIS	-	expression tag	UNP B3VI58
A	71	GLU	ILE	conflict	UNP B3VI58
A	72	ASN	PRO	conflict	UNP B3VI58
A	73	LEU	GLN	conflict	UNP B3VI58
A	74	TYR	GLN	conflict	UNP B3VI58
A	75	PHE	GLU	conflict	UNP B3VI58
A	76	GLN	HIS	conflict	UNP B3VI58

- Molecule 2 is SPERMINE (CCD ID: SPM) (formula: $C_{10}H_{26}N_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			14	10	4		
2	A	1	Total	C	N	0	0
			14	10	4		
2	A	1	Total	C	N	0	0
			14	10	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	518	Total	O	0	0
			518	518		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Oligopeptidase B



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.40Å 100.45Å 108.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.40 – 1.70 29.40 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.6 (29.40-1.70) 98.6 (29.40-1.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.40 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.185 , 0.218 0.193 , 0.227	Depositor DCC
R_{free} test set	4484 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	22.2	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 28.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6128	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SPM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	0/5714	1.22	3/7768 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	154	ASP	CA-CB-CG	5.23	117.83	112.60
1	A	57	GLN	CA-C-O	-5.18	115.35	120.90
1	A	355	PHE	CA-CB-CG	-5.02	108.78	113.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	617	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5568	0	5273	68	0
2	A	42	0	78	6	0
3	A	518	0	0	10	0
All	All	6128	0	5351	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:LEU:CD2	1:A:141:LEU:HD11	1.75	1.17
1:A:132:LEU:HD21	1:A:141:LEU:HD11	1.32	1.04
1:A:367:ASN:HD21	1:A:376:ARG:H	1.21	0.88
1:A:621:GLN:HA	3:A:1141:HOH:O	1.74	0.88
1:A:245:THR:HG23	1:A:246:THR:OG1	1.74	0.88
1:A:64:TYR:HH	1:A:659:PHE:HD1	1.23	0.84
1:A:377:TYR:OH	1:A:398:ARG:HD2	1.76	0.82
1:A:132:LEU:HD23	1:A:141:LEU:HD11	1.63	0.78
1:A:300:GLN:NE2	1:A:305:GLN:HG2	2.00	0.77
1:A:300:GLN:HE21	1:A:305:GLN:HG2	1.51	0.75
1:A:337:LEU:HD12	3:A:1033:HOH:O	1.87	0.73
1:A:271:HIS:HE1	1:A:286:SER:OG	1.74	0.71
1:A:441:ALA:H	1:A:445:ASN:HD21	1.41	0.69
1:A:442:ARG:HH22	2:A:701:SPM:H31	1.60	0.67
1:A:64:TYR:OH	1:A:659:PHE:HD1	1.79	0.66
1:A:115:ILE:HD11	1:A:120:ARG:HD2	1.78	0.65
1:A:516:ILE:HD11	1:A:526:VAL:HG21	1.79	0.64
2:A:701:SPM:H61	3:A:1168:HOH:O	1.97	0.63
1:A:621:GLN:HB3	1:A:623:TRP:CZ3	2.34	0.63
1:A:86:ARG:HH11	1:A:103:GLN:HE21	1.46	0.62
1:A:120:ARG:HD3	1:A:145:GLU:OE1	2.01	0.61
1:A:279:HIS:HE1	3:A:959:HOH:O	1.83	0.61
1:A:652:HIS:CE1	3:A:1146:HOH:O	2.55	0.59
1:A:300:GLN:HE21	1:A:305:GLN:CG	2.16	0.58
1:A:314:ILE:HG13	3:A:1193:HOH:O	2.04	0.57
1:A:120:ARG:CD	1:A:145:GLU:OE1	2.52	0.57
1:A:442:ARG:HD2	2:A:701:SPM:H111	1.88	0.55
1:A:571:PRO:O	1:A:572:LEU:HB2	2.06	0.54
1:A:447:LEU:HD23	1:A:476:VAL:HB	1.89	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:LEU:CD2	1:A:141:LEU:CD1	2.68	0.54
1:A:78:VAL:CG2	1:A:402:LYS:HB2	2.38	0.53
1:A:337:LEU:CD1	3:A:1033:HOH:O	2.53	0.53
1:A:344:HIS:HD2	1:A:346:GLN:H	1.56	0.53
1:A:132:LEU:HD21	1:A:141:LEU:CD1	2.22	0.52
1:A:245:THR:HG21	1:A:269:LYS:NZ	2.25	0.51
1:A:73:LEU:H	1:A:73:LEU:HD23	1.76	0.51
1:A:271:HIS:HD2	3:A:1068:HOH:O	1.92	0.51
1:A:271:HIS:CE1	1:A:286:SER:OG	2.61	0.50
1:A:73:LEU:HD12	1:A:74:TYR:CE1	2.47	0.50
1:A:203:ARG:HD2	1:A:214:GLU:OE1	2.12	0.49
1:A:613:THR:OG1	1:A:622:TYR:HA	2.13	0.49
1:A:83:HIS:HD2	3:A:1030:HOH:O	1.95	0.48
1:A:654:GLY:O	1:A:655:LYS:HB2	2.13	0.48
1:A:279:HIS:HD2	3:A:859:HOH:O	1.96	0.48
1:A:314:ILE:HD12	1:A:315:GLU:N	2.30	0.47
1:A:152:GLN:OE1	1:A:172[A]:ASN:ND2	2.47	0.47
1:A:344:HIS:CD2	1:A:346:GLN:H	2.33	0.46
1:A:649:ASP:HB3	1:A:651:GLY:O	2.14	0.46
1:A:624:GLU:HB2	1:A:625:PRO:HD3	1.97	0.46
1:A:340:LEU:HB2	1:A:353:ILE:CG1	2.46	0.46
1:A:143:VAL:O	1:A:155:ILE:HA	2.16	0.46
1:A:314:ILE:HD12	1:A:315:GLU:CD	2.41	0.45
1:A:459:MET:HE1	2:A:703:SPM:H92	1.99	0.45
1:A:86:ARG:HD2	1:A:103:GLN:HE21	1.82	0.44
1:A:613:THR:HB	1:A:625:PRO:HG2	2.00	0.44
1:A:86:ARG:HH11	1:A:103:GLN:NE2	2.15	0.44
1:A:340:LEU:HB2	1:A:353:ILE:HG12	2.00	0.43
1:A:344:HIS:CD2	1:A:347:SER:H	2.36	0.43
1:A:17:HIS:HE1	1:A:583:PRO:O	2.01	0.43
1:A:621:GLN:CB	1:A:623:TRP:CH2	3.02	0.43
1:A:557:PRO:O	1:A:620:VAL:HG11	2.19	0.42
1:A:226:VAL:HG22	1:A:241:LEU:HD12	2.01	0.42
1:A:442:ARG:HH12	2:A:701:SPM:C3	2.32	0.42
1:A:120:ARG:HD2	1:A:145:GLU:OE1	2.20	0.41
1:A:79:PRO:HA	1:A:87:TYR:O	2.20	0.41
1:A:314:ILE:HG13	1:A:314:ILE:H	1.73	0.41
1:A:124:ARG:HA	1:A:124:ARG:HE	1.86	0.41
1:A:280:GLN:HE21	1:A:280:GLN:HA	1.86	0.40
2:A:702:SPM:H111	2:A:702:SPM:H81	1.95	0.40
1:A:64:TYR:OH	1:A:659:PHE:CD1	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	678/677 (100%)	642 (95%)	29 (4%)	7 (1%)	12 3

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	618	SER
1	A	619	GLN
1	A	653	GLY
1	A	572	LEU
1	A	617	ASP
1	A	655	LYS
1	A	654	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	594/591 (100%)	565 (95%)	29 (5%)	22 8

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	16	THR
1	A	37	GLN
1	A	54	LYS
1	A	73	LEU
1	A	92	GLU
1	A	96	GLU
1	A	109	GLU
1	A	120	ARG
1	A	124	ARG
1	A	132	LEU
1	A	150	ARG
1	A	191	LYS
1	A	241	LEU
1	A	244	THR
1	A	249	ILE
1	A	250	LEU
1	A	252	LEU
1	A	280	GLN
1	A	305	GLN
1	A	309	LEU
1	A	315	GLU
1	A	337	LEU
1	A	373	GLU
1	A	515	LEU
1	A	548	LEU
1	A	636	LYS
1	A	652	HIS
1	A	655	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1	HIS
1	A	17	HIS
1	A	37	GLN
1	A	72	ASN
1	A	83	HIS
1	A	103	GLN
1	A	119	GLN
1	A	181	ASN
1	A	271	HIS
1	A	279	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	280	GLN
1	A	297	GLN
1	A	300	GLN
1	A	339	GLN
1	A	344	HIS
1	A	346	GLN
1	A	367	ASN
1	A	445	ASN
1	A	502	GLN
1	A	518	GLN
1	A	544	GLN
1	A	585	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SPM	A	702	-	13,13,13	0.23	0	12,12,12	0.58	0
2	SPM	A	701	-	13,13,13	0.36	0	12,12,12	0.90	1 (8%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SPM	A	703	-	13,13,13	0.17	0	12,12,12	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SPM	A	702	-	-	6/11/11/11	-
2	SPM	A	701	-	-	8/11/11/11	-
2	SPM	A	703	-	-	7/11/11/11	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	SPM	C7-C6-N5	2.36	118.41	112.07

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	SPM	C8-C9-N10-C11
2	A	703	SPM	C7-C8-C9-N10
2	A	701	SPM	N10-C11-C12-C13
2	A	702	SPM	C2-C3-C4-N5
2	A	703	SPM	N5-C6-C7-C8
2	A	701	SPM	C12-C11-N10-C9
2	A	701	SPM	N5-C6-C7-C8
2	A	702	SPM	C7-C6-N5-C4
2	A	703	SPM	C6-C7-C8-C9
2	A	702	SPM	C7-C8-C9-N10
2	A	701	SPM	C6-C7-C8-C9
2	A	703	SPM	N10-C11-C12-C13
2	A	702	SPM	C6-C7-C8-C9
2	A	702	SPM	C3-C4-N5-C6
2	A	703	SPM	C2-C3-C4-N5
2	A	701	SPM	N1-C2-C3-C4
2	A	701	SPM	C11-C12-C13-N14
2	A	703	SPM	C3-C4-N5-C6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	703	SPM	C8-C9-N10-C11
2	A	702	SPM	C8-C9-N10-C11
2	A	701	SPM	C3-C4-N5-C6

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	702	SPM	1	0
2	A	701	SPM	4	0
2	A	703	SPM	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	677/677 (100%)	0.14	32 (4%)	36 40	12, 23, 45, 134	3 (0%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	655	LYS	5.9
1	A	653	GLY	5.5
1	A	656	SER	5.4
1	A	657	GLY	5.3
1	A	652	HIS	5.2
1	A	73	LEU	5.0
1	A	659	PHE	4.8
1	A	620	VAL	4.6
1	A	654	GLY	4.5
1	A	651	GLY	4.4
1	A	74	TYR	4.0
1	A	650	SER	3.9
1	A	94	GLY	3.8
1	A	257	ALA	3.7
1	A	1	HIS	3.6
1	A	616	HIS	3.5
1	A	356	ASP	3.1
1	A	658	ARG	3.0
1	A	618	SER	3.0
1	A	354	ALA	3.0
1	A	355	PHE	3.0
1	A	75	PHE	2.9
1	A	619	GLN	2.7
1	A	193	ALA	2.6
1	A	353	ILE	2.6
1	A	148	LEU	2.5
1	A	150	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	639	ASP	2.2
1	A	132	LEU	2.1
1	A	96	GLU	2.1
1	A	93	PRO	2.1
1	A	571	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SPM	A	703	14/14	0.75	0.19	47,50,66,69	0
2	SPM	A	702	14/14	0.82	0.18	40,47,60,61	0
2	SPM	A	701	14/14	0.82	0.14	30,35,38,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.