



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:09 PM UTC

PDB ID : 7YYZ / pdb_00007yyz
Title : Molecular snapshots of drug release from tubulin: 10 microseconds after photoactivation.
Authors : Wranik, M.; Weinert, T.; Standfuss, J.
Deposited on : 2022-02-18
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	7% 87% 10% .
2	B	445	16% 82% 13% .
3	F	169	9% 83% 8% 8%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 8228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	437	Total	C	N	O	S	0	2	0
			3415	2163	579	650	23			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	426	Total	C	N	O	S	0	2	0
			3338	2099	566	646	27			

- Molecule 3 is a protein called Designed Ankyrin Repeat Protein (DARPIN) D1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	155	Total	C	N	O	S	0	0	0
			1154	727	198	226	3			

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		


- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 8 is Azo-Combretastatin A4 (trans) (CCD ID: VYT) (formula: C₁₆H₁₈N₂O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	C	N	O	0	0
			23	16	2	5		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	129	Total	O	0	0
			129	129		
9	B	69	Total	O	0	0
			69	69		
9	F	38	Total	O	0	0
			38	38		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain




- Molecule 2: Tubulin beta-2B chain



- Molecule 3: Designed Ankyrin Repeat Protein (DARPIN) D1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.53Å 92.58Å 83.99Å 90.00° 96.71° 90.00°	Depositor
Resolution (Å)	9.49 – 2.20 9.49 – 2.20	Depositor EDS
% Data completeness (in resolution range)	93.6 (9.49-2.20) 92.4 (9.49-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.78 (at 2.21Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.299 , 0.355 (Not available) , 0.396	Depositor DCC
R_{free} test set	1859 reflections (3.50%)	wwPDB-VP
Wilson B-factor (Å ²)	30.6	Xtriage
Anisotropy	0.221	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 64.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	8228	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VYT, CA, GDP, MG, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.07	0/3499	0.22	0/4752
2	B	0.07	0/3417	0.21	0/4629
3	F	0.06	0/1170	0.17	0/1590
All	All	0.07	0/8086	0.21	0/10971

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3415	0	3319	27	0
2	B	3338	0	3213	41	0
3	F	1154	0	1156	11	0
4	A	32	0	12	1	0
5	A	1	0	0	0	0
6	A	1	0	0	0	0
7	B	28	0	12	1	0
8	B	23	0	0	1	0
9	A	129	0	0	2	0
9	B	69	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	F	38	0	0	1	0
All	All	8228	0	7712	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:39:ASP:N	2:B:39:ASP:OD1	2.16	0.78
2:B:268:PRO:HG2	2:B:300:MET:HB2	1.74	0.70
1:A:100:ALA:O	9:A:601:HOH:O	2.08	0.70
1:A:172:TYR:HB3	1:A:205:ASP:HA	1.74	0.69
2:B:354:CYS:SG	9:B:655:HOH:O	2.50	0.69
1:A:304:LYS:O	1:A:390:ARG:NH2	2.28	0.66
2:B:113:VAL:HG23	2:B:151:LEU:HD22	1.78	0.65
2:B:293:MET:HG2	2:B:367:PHE:HB2	1.80	0.64
2:B:213:ARG:HH12	3:F:159:GLU:HB2	1.64	0.62
2:B:247:ASN:HD22	2:B:249:ASP:HB2	1.64	0.62
3:F:79:MET:HE3	3:F:111:THR:HG21	1.84	0.60
2:B:396:HIS:HE2	3:F:57:TYR:HH	1.49	0.59
2:B:249:ASP:HB3	2:B:252:LYS:HB2	1.84	0.59
1:A:399:TYR:O	1:A:402:ARG:NH1	2.35	0.59
2:B:213:ARG:NH2	3:F:160:ASP:OD1	2.37	0.57
2:B:121:ARG:NH2	2:B:158:GLU:OE2	2.37	0.57
2:B:378:PHE:HD1	2:B:415:MET:HE3	1.71	0.56
3:F:121:ALA:HB1	3:F:161:LEU:HD21	1.88	0.55
1:A:26:LEU:HD12	1:A:363:VAL:HG12	1.90	0.54
2:B:396:HIS:NE2	3:F:57:TYR:OH	2.33	0.54
3:F:23:ARG:NH2	9:F:204:HOH:O	2.40	0.53
1:A:263:PRO:O	1:A:266:HIS:ND1	2.35	0.52
2:B:73:MET:HG3	2:B:77:ARG:HE	1.73	0.52
2:B:117:LEU:HD11	2:B:154:LYS:HB3	1.92	0.52
1:A:93:ILE:HG22	1:A:114:ILE:HD11	1.92	0.51
1:A:401:LYS:NZ	2:B:429:THR:O	2.31	0.51
2:B:42:LEU:HD22	2:B:243:PRO:HG2	1.91	0.51
2:B:54:ALA:HB3	2:B:58:LYS:HB3	1.93	0.51
1:A:15:GLN:NE2	4:A:501:GTP:O6	2.44	0.50
2:B:12:CYS:HB2	7:B:501:GDP:C8	2.45	0.50
2:B:183:TYR:O	2:B:187:LEU:HG	2.12	0.50
3:F:31:ARG:O	3:F:31:ARG:NH1	2.45	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:236:VAL:HG23	2:B:237:THR:HG23	1.94	0.49
2:B:337:ASN:HB3	2:B:340:TYR:HD2	1.79	0.48
1:A:270:ALA:HB3	1:A:302[B]:MET:HE2	1.97	0.47
2:B:213:ARG:HH22	3:F:159:GLU:HB2	1.80	0.47
2:B:61:PRO:HD3	2:B:84:ILE:HG13	1.97	0.46
1:A:328:VAL:HG11	1:A:353:VAL:HG11	1.97	0.46
2:B:253:LEU:HD13	8:B:502:VYT:C6	2.46	0.45
2:B:28:HIS:CE1	2:B:241:ARG:HB3	2.51	0.45
1:A:90:GLU:O	1:A:121:ARG:NH1	2.46	0.45
1:A:296:PHE:HB2	1:A:339:ARG:NH1	2.32	0.44
1:A:200:CYS:HA	1:A:266:HIS:HB2	1.98	0.44
1:A:304:LYS:HA	1:A:304:LYS:HE2	1.99	0.44
2:B:113:VAL:HG21	2:B:154:LYS:HG3	2.00	0.44
1:A:62:VAL:HG11	1:A:88:HIS:CD2	2.53	0.44
2:B:68:LEU:HD23	2:B:112:LEU:HD22	1.99	0.44
1:A:98:ASP:OD2	2:B:252:LYS:NZ	2.48	0.44
3:F:60:LEU:HD11	3:F:98:VAL:HG21	1.99	0.44
2:B:407:GLU:HG2	9:B:609:HOH:O	2.17	0.44
1:A:205:ASP:HB2	1:A:303:VAL:HA	2.01	0.43
2:B:45:GLU:HB2	2:B:243:PRO:HG3	2.01	0.43
2:B:202:ILE:HD13	2:B:229:VAL:HG13	1.99	0.43
3:F:40:VAL:HG13	3:F:66:LEU:HD22	1.99	0.43
1:A:21:TRP:CZ3	1:A:63:PRO:HB3	2.54	0.43
2:B:104:GLY:O	2:B:109:GLY:HA3	2.19	0.42
2:B:23:VAL:HG21	2:B:230:SER:HB2	2.00	0.42
1:A:70:LEU:HB2	1:A:98:ASP:HA	2.01	0.42
1:A:343:PHE:CD1	1:A:349:THR:HG23	2.55	0.42
9:A:601:HOH:O	2:B:255:VAL:HG11	2.19	0.42
2:B:211:CYS:HB3	2:B:217:LEU:HD12	2.02	0.42
2:B:9:ALA:HA	2:B:66:VAL:O	2.20	0.42
1:A:188:ILE:HD12	1:A:425:MET:HG3	2.02	0.42
1:A:140:SER:HA	1:A:171:ILE:HB	2.01	0.41
1:A:67:PHE:HB2	1:A:92:LEU:HD23	2.03	0.41
1:A:409:VAL:HA	1:A:413:MET:O	2.20	0.41
2:B:11:GLN:O	2:B:15:GLN:HB2	2.21	0.41
2:B:178:THR:HG22	2:B:180:VAL:HG22	2.03	0.40
2:B:384:GLN:NE2	9:B:620:HOH:O	2.53	0.40
1:A:188:ILE:HG23	1:A:425:MET:HG3	2.03	0.40
1:A:176:GLN:OE1	1:A:176:GLN:N	2.52	0.40
2:B:28:HIS:NE2	2:B:241:ARG:HB3	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	437/451 (97%)	421 (96%)	14 (3%)	2 (0%)	24	27
2	B	424/445 (95%)	414 (98%)	10 (2%)	0	100	100
3	F	153/169 (90%)	151 (99%)	2 (1%)	0	100	100
All	All	1014/1065 (95%)	986 (97%)	26 (3%)	2 (0%)	43	51

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	349	THR
1	A	322	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/379 (97%)	366 (100%)	1 (0%)	86	93
2	B	366/383 (96%)	361 (99%)	5 (1%)	59	75
3	F	120/132 (91%)	119 (99%)	1 (1%)	73	85
All	All	853/894 (95%)	846 (99%)	7 (1%)	73	85

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	39	ASP
2	B	151	LEU
2	B	221	THR
2	B	246	LEU
2	B	353	VAL
3	F	144	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	GLN
1	A	128	GLN
1	A	133	GLN
1	A	233	GLN
1	A	393	HIS
2	B	8	GLN
2	B	134	GLN
2	B	227	HIS
2	B	247	ASN
2	B	334	GLN
2	B	384	GLN
3	F	59	HIS
3	F	166	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	VYT	B	502	-	24,24,24	0.21	0	32,32,32	0.19	0
7	GDP	B	501	-	29,30,30	1.17	3 (10%)	45,47,47	1.79	7 (15%)
4	GTP	A	501	5	33,34,34	1.00	2 (6%)	50,54,54	1.57	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	VYT	B	502	-	-	0/13/13/13	0/2/2/2
7	GDP	B	501	-	-	5/16/32/32	0/3/3/3
4	GTP	A	501	5	-	9/22/38/38	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	501	GDP	C5-C4	3.15	1.47	1.38
7	B	501	GDP	C6-N1	-2.38	1.34	1.38
4	A	501	GTP	PA-O3A	2.28	1.62	1.59
4	A	501	GTP	C2-N3	2.22	1.38	1.33
7	B	501	GDP	C5-N7	-2.02	1.35	1.39

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	501	GDP	C5-C4-N3	-6.13	118.63	128.39
7	B	501	GDP	C2-N3-C4	5.05	120.99	112.30
4	A	501	GTP	C5-C4-N3	-4.87	120.63	128.39
4	A	501	GTP	C2-N3-C4	4.61	120.25	112.30
7	B	501	GDP	N9-C4-N3	4.51	134.97	125.95
7	B	501	GDP	C6-C5-N7	3.21	136.14	130.29
4	A	501	GTP	N9-C4-N3	3.09	132.13	125.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	501	GTP	C2-N1-C6	-2.84	119.96	125.11
4	A	501	GTP	N9-C8-N7	-2.71	108.37	113.40
7	B	501	GDP	C4-C5-N7	-2.57	106.60	110.67
4	A	501	GTP	C8-N7-C5	2.52	108.75	104.26
4	A	501	GTP	C5-C6-N1	2.51	119.64	113.25
4	A	501	GTP	O6-C6-C5	-2.39	120.23	126.53
7	B	501	GDP	O6-C6-C5	-2.11	120.95	126.53
7	B	501	GDP	C3'-C2'-C1'	2.05	105.35	101.46

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	501	GTP	PB-O3B-PG-O2G
4	A	501	GTP	C5'-O5'-PA-O3A
4	A	501	GTP	C5'-O5'-PA-O2A
7	B	501	GDP	C5'-O5'-PA-O3A
7	B	501	GDP	C5'-O5'-PA-O1A
7	B	501	GDP	C5'-O5'-PA-O2A
4	A	501	GTP	C3'-C4'-C5'-O5'
4	A	501	GTP	C5'-O5'-PA-O1A
4	A	501	GTP	O4'-C4'-C5'-O5'
4	A	501	GTP	C4'-C5'-O5'-PA
7	B	501	GDP	PB-O3A-PA-O2A
4	A	501	GTP	PB-O3A-PA-O1A
4	A	501	GTP	PB-O3A-PA-O2A
7	B	501	GDP	PB-O3A-PA-O1A


There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	502	VYT	1	0
7	B	501	GDP	1	0
4	A	501	GTP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	437/451 (96%)	0.79	30 (6%) 23 20	14, 28, 48, 72	2 (0%)
2	B	426/445 (95%)	1.26	73 (17%) 4 3	15, 38, 66, 78	2 (0%)
3	F	155/169 (91%)	0.74	15 (9%) 13 10	21, 29, 44, 60	0
All	All	1018/1065 (95%)	0.98	118 (11%) 9 7	14, 32, 61, 78	4 (0%)

All (118) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	79	GLY	6.8
2	B	73	MET	6.3
1	A	335	ILE	5.6
2	B	288	GLU	4.6
2	B	277	GLY	4.5
1	A	339	ARG	4.2
1	A	338	LYS	4.1
2	B	356	ILE	4.1
2	B	57	ASN	4.0
1	A	59	GLY	4.0
2	B	335	ASN	4.0
2	B	38	GLY	3.8
2	B	333	VAL	3.7
2	B	80	PRO	3.7
2	B	29	GLY	3.6
1	A	337	THR	3.5
3	F	31	ARG	3.5
2	B	119	VAL	3.5
2	B	36	TYR	3.5
2	B	22	GLU	3.4
2	B	247	ASN	3.4
1	A	334	THR	3.4
2	B	58	LYS	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	361	LEU	3.3
2	B	248	ALA	3.3
2	B	217	LEU	3.2
2	B	322	SER	3.2
1	A	400	ALA	3.2
2	B	39	ASP	3.1
1	A	426	ALA	3.1
2	B	76	VAL	3.1
2	B	84	ILE	3.1
2	B	128	ASP	3.1
2	B	40	SER	3.0
2	B	278	SER	3.0
1	A	347	CYS	2.9
2	B	326	VAL	2.9
2	B	91	VAL	2.9
1	A	80	THR	2.8
1	A	416	GLY	2.8
2	B	78	SER	2.8
2	B	308	GLY	2.8
1	A	437	VAL	2.7
2	B	55	THR	2.7
2	B	93	GLY	2.7
3	F	13	ASP	2.7
1	A	297	GLU	2.7
2	B	359	ARG	2.7
3	F	150	PHE	2.6
1	A	349	THR	2.6
2	B	56	GLY	2.6
2	B	5	VAL	2.6
2	B	92	PHE	2.6
1	A	44	GLY	2.6
2	B	82	GLY	2.6
2	B	30	ILE	2.6
1	A	88	HIS	2.6
2	B	357	PRO	2.6
2	B	35	SER	2.6
1	A	18	ASN	2.5
1	A	146	GLY	2.5
3	F	75	ALA	2.5
2	B	346	PRO	2.5
2	B	89	ASN	2.5
2	B	431	ASP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	41	THR	2.4
1	A	283	HIS	2.4
1	A	17	GLY	2.4
2	B	17	GLY	2.4
2	B	221	THR	2.4
2	B	165	ASN	2.4
1	A	333	ALA	2.4
2	B	238	THR	2.4
3	F	16	LYS	2.4
2	B	75	SER	2.4
2	B	275	SER	2.4
2	B	118	ASP	2.3
2	B	274	THR	2.3
3	F	25	GLY	2.3
1	A	343	PHE	2.3
2	B	430	ALA	2.3
2	B	54	ALA	2.3
1	A	159	VAL	2.3
3	F	44	ASP	2.2
2	B	45	GLU	2.2
2	B	229	VAL	2.2
2	B	360	GLY	2.2
3	F	37	GLY	2.2
2	B	242	PHE	2.2
3	F	145	PHE	2.2
2	B	397	TRP	2.2
1	A	42	ILE	2.2
2	B	336	LYS	2.2
2	B	218	THR	2.2
2	B	276	ARG	2.1
3	F	93	LEU	2.1
2	B	31	ASP	2.1
2	B	157	GLU	2.1
1	A	178	SER	2.1
1	A	232	SER	2.1
2	B	123	GLU	2.1
3	F	151	ASP	2.1
2	B	220	PRO	2.1
2	B	421	GLU	2.1
3	F	164	ILE	2.1
1	A	372	GLN	2.1
1	A	123	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	409	THR	2.1
2	B	49	VAL	2.1
2	B	81	PHE	2.1
2	B	85	PHE	2.1
2	B	321	MET	2.0
2	B	410	GLU	2.0
3	F	104	ALA	2.0
3	F	160	ASP	2.0
2	B	328	GLU	2.0
1	A	306	ASP	2.0
3	F	167	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	VYT	B	502	23/23	0.78	0.12	23,38,46,47	0
6	CA	A	503	1/1	0.88	0.08	40,40,40,40	0
7	GDP	B	501	28/28	0.89	0.11	27,35,39,46	0
4	GTP	A	501	32/32	0.90	0.11	15,21,25,28	0
5	MG	A	502	1/1	0.96	0.19	11,11,11,11	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GTP A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.