



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 09:56 AM UTC

PDB ID : 4YZF / pdb_00004yzf
Title : Crystal structure of the anion exchanger domain of human erythrocyte Band 3
Authors : Alguel, Y.; Arakawa, T.; Yugiri, T.K.; Iwanari, H.; Hatae, H.; Iwata, M.; Abe, Y.; Hino, T.; Suno, C.I.; Kuma, H.; Kang, D.; Murata, T.; Hamakubo, T.; Cameron, A.D.; Kobayashi, T.; Hamasaki, N.; Iwata, S.
Deposited on : 2015-03-25
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1085 (3.54-3.46)
Clashscore	190562	1140 (3.54-3.46)
Ramachandran outliers	187476	1113 (3.54-3.46)
Sidechain outliers	187428	1114 (3.54-3.46)
RSRZ outliers	180081	1084 (3.54-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	911	3% 30% 20% . 48%
1	B	911	4% 31% 19% . 48%
1	C	911	4% 30% 20% . 48%
1	D	911	6% 31% 20% . 48%
2	E	223	7% 72% 23% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	223	
2	I	223	
2	K	223	
3	F	218	
3	H	218	
3	J	218	
3	L	218	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 28724 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Band 3 anion transport protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	475	Total	C	N	O	S	0	0	0
			3769	2535	598	619	17			
1	B	475	Total	C	N	O	S	0	0	0
			3769	2535	598	619	17			
1	C	475	Total	C	N	O	S	0	0	0
			3769	2535	598	619	17			
1	D	475	Total	C	N	O	S	0	0	0
			3769	2535	598	619	17			


- Molecule 2 is a protein called FAB fragment of Immunoglobulin (IgG) molecule.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	223	Total	C	N	O	S	0	0	0
			1690	1077	274	334	5			
2	G	223	Total	C	N	O	S	0	0	0
			1690	1077	274	334	5			
2	I	223	Total	C	N	O	S	0	0	0
			1690	1077	274	334	5			
2	K	223	Total	C	N	O	S	0	0	0
			1690	1077	274	334	5			

- Molecule 3 is a protein called FAB fragment of Immunoglobulin (IgG) molecule.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	218	Total	C	N	O	S	0	0	0
			1694	1055	285	346	8			
3	H	218	Total	C	N	O	S	0	0	0
			1694	1055	285	346	8			
3	J	218	Total	C	N	O	S	0	0	0
			1694	1055	285	346	8			
3	L	218	Total	C	N	O	S	0	0	0
			1694	1055	285	346	8			


- Molecule 4 is 2,2'-ethane-1,2-diylbis{5-[(sulfanylmethyl)amino]benzenesulfonic acid} (CCD ID: 4KU) (formula: C₁₆H₂₀N₂O₆S₄).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			28	16	2	6	4		
4	B	1	Total	C	N	O	S	0	0
			28	16	2	6	4		
4	C	1	Total	C	N	O	S	0	0
			28	16	2	6	4		
4	D	1	Total	C	N	O	S	0	0
			28	16	2	6	4		

- Molecule 1: Band 3 anion transport protein







• Molecule 1: Band 3 anion transport protein




• Molecule 1: Band 3 anion transport protein







• Molecule 2: FAB fragment of Immunoglobulin (IgG) molecule




• Molecule 2: FAB fragment of Immunoglobulin (IgG) molecule









- Molecule 2: FAB fragment of Immunoglobulin (IgG) molecule




- Molecule 2: FAB fragment of Immunoglobulin (IgG) molecule




- Molecule 3: FAB fragment of Immunoglobulin (IgG) molecule




- Molecule 3: FAB fragment of Immunoglobulin (IgG) molecule







- Molecule 3: FAB fragment of Immunoglobulin (IgG) molecule



- Molecule 3: FAB fragment of Immunoglobulin (IgG) molecule



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	152.82Å 171.96Å 271.70Å 90.00° 101.16° 90.00°	Depositor
Resolution (Å)	37.72 – 3.50 37.72 – 3.50	Depositor EDS
% Data completeness (in resolution range)	94.6 (37.72-3.50) 94.8 (37.72-3.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 3.48Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1951)	Depositor
R, R_{free}	0.274 , 0.290 0.277 , 0.295	Depositor DCC
R_{free} test set	4135 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	115.1	Xtriage
Anisotropy	0.214	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 86.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	28724	wwPDB-VP
Average B, all atoms (Å ²)	145.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.75 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.3662e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 4KU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/3862	1.00	22/5246 (0.4%)
1	B	0.49	0/3862	0.97	14/5246 (0.3%)
1	C	0.53	0/3862	0.98	18/5246 (0.3%)
1	D	0.49	0/3862	0.94	14/5246 (0.3%)
2	E	0.59	0/1737	1.32	18/2377 (0.8%)
2	G	0.48	0/1737	0.93	3/2377 (0.1%)
2	I	0.55	0/1737	1.20	15/2377 (0.6%)
2	K	0.47	0/1737	0.92	3/2377 (0.1%)
3	F	0.55	0/1736	0.93	2/2360 (0.1%)
3	H	0.47	1/1736 (0.1%)	0.84	1/2360 (0.0%)
3	J	0.48	0/1736	0.92	3/2360 (0.1%)
3	L	0.44	0/1736	0.82	1/2360 (0.0%)
All	All	0.51	1/29340 (0.0%)	0.99	114/39932 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	H	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	112	ARG	CA-C	5.01	1.59	1.52

The worst 5 of 114 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	867	VAL	CA-C-N	-9.20	110.80	119.82
1	A	867	VAL	C-N-CA	-9.20	110.80	119.82
2	E	96	TYR	N-CA-C	8.99	123.26	108.96
2	E	97	CYS	N-CA-C	-8.58	96.64	110.20
1	C	659	PHE	CA-C-N	8.56	128.63	119.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	H	111	LYS	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3769	0	3990	154	0
1	B	3769	0	3990	144	0
1	C	3769	0	3988	141	0
1	D	3769	0	3989	144	0
2	E	1690	0	1645	40	0
2	G	1690	0	1645	28	0
2	I	1690	0	1645	46	0
2	K	1690	0	1645	32	0
3	F	1694	0	1609	34	0
3	H	1694	0	1609	29	0
3	J	1694	0	1609	40	0
3	L	1694	0	1609	27	0
4	A	28	0	15	1	0
4	B	28	0	14	2	0
4	C	28	0	15	3	0
4	D	28	0	15	2	0
All	All	28724	0	29032	811	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 811 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:828:VAL:HG23	1:A:829:LYS:HG3	1.48	0.94
1:D:851:LYS:HE3	1:D:859:LEU:HD22	1.51	0.92
1:C:828:VAL:HG23	1:C:829:LYS:HG3	1.49	0.92
1:D:828:VAL:HG23	1:D:829:LYS:HG3	1.54	0.90
1:D:737:ALA:HB3	1:D:738:LEU:HB2	1.55	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	467/911 (51%)	435 (93%)	31 (7%)	1 (0%)	43	74
1	B	467/911 (51%)	436 (93%)	30 (6%)	1 (0%)	43	74
1	C	467/911 (51%)	436 (93%)	30 (6%)	1 (0%)	43	74
1	D	467/911 (51%)	436 (93%)	30 (6%)	1 (0%)	43	74
2	E	221/223 (99%)	211 (96%)	7 (3%)	3 (1%)	9	38
2	G	221/223 (99%)	214 (97%)	7 (3%)	0	100	100
2	I	221/223 (99%)	212 (96%)	6 (3%)	3 (1%)	9	38
2	K	221/223 (99%)	213 (96%)	8 (4%)	0	100	100
3	F	216/218 (99%)	208 (96%)	8 (4%)	0	100	100
3	H	216/218 (99%)	209 (97%)	7 (3%)	0	100	100
3	J	216/218 (99%)	209 (97%)	7 (3%)	0	100	100
3	L	216/218 (99%)	211 (98%)	5 (2%)	0	100	100
All	All	3616/5408 (67%)	3430 (95%)	176 (5%)	10 (0%)	36	67

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	166	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	I	166	SER
2	E	159	PRO
2	I	159	PRO
2	E	210	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	417/786 (53%)	396 (95%)	21 (5%)	22	48
1	B	417/786 (53%)	399 (96%)	18 (4%)	26	51
1	C	417/786 (53%)	398 (95%)	19 (5%)	24	50
1	D	417/786 (53%)	399 (96%)	18 (4%)	26	51
2	E	190/190 (100%)	184 (97%)	6 (3%)	34	59
2	G	190/190 (100%)	185 (97%)	5 (3%)	40	62
2	I	190/190 (100%)	184 (97%)	6 (3%)	34	59
2	K	190/190 (100%)	185 (97%)	5 (3%)	40	62
3	F	193/193 (100%)	188 (97%)	5 (3%)	40	62
3	H	193/193 (100%)	189 (98%)	4 (2%)	47	66
3	J	193/193 (100%)	188 (97%)	5 (3%)	40	62
3	L	193/193 (100%)	189 (98%)	4 (2%)	47	66
All	All	3200/4676 (68%)	3084 (96%)	116 (4%)	31	57

5 of 116 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	877	ILE
2	K	160	VAL
1	D	769	LEU
2	K	63	THR
2	I	160	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 40 such sidechains are listed below:

Mol	Chain	Res	Type
2	I	79	GLN
3	L	41	GLN
2	I	174	HIS
2	K	16	GLN
3	L	80	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	4KU	D	1000	1	29,29,29	1.74	3 (10%)	36,42,42	2.58	14 (38%)
4	4KU	B	1000	1	29,29,29	1.72	4 (13%)	36,42,42	2.78	14 (38%)
4	4KU	C	1000	1	29,29,29	1.71	4 (13%)	36,42,42	2.77	14 (38%)
4	4KU	A	1000	1	29,29,29	1.75	4 (13%)	36,42,42	3.24	16 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	4KU	D	1000	1	-	6/21/23/23	0/2/2/2
4	4KU	B	1000	1	-	7/21/23/23	0/2/2/2
4	4KU	C	1000	1	-	6/21/23/23	0/2/2/2
4	4KU	A	1000	1	-	2/21/23/23	0/2/2/2

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1000	4KU	CAI-SAG	-6.35	1.66	1.81
4	D	1000	4KU	CAI-SAG	-6.18	1.66	1.81
4	C	1000	4KU	CAI-SAG	-6.06	1.66	1.81
4	B	1000	4KU	CAI-SAG	-5.97	1.67	1.81
4	D	1000	4KU	CAJ-SAH	-5.95	1.67	1.81

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1000	4KU	CAR-CAX-CAZ	-9.97	115.58	125.37
4	A	1000	4KU	CAR-CAQ-CAW	-7.99	99.98	112.76
4	B	1000	4KU	CAR-CAX-CAZ	-7.82	117.69	125.37
4	C	1000	4KU	CAR-CAX-CAZ	-7.65	117.86	125.37
4	D	1000	4KU	CAR-CAX-CAZ	-7.48	118.03	125.37

There are no chirality outliers.

5 of 21 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	1000	4KU	CAW-CAQ-CAR-CAX
4	B	1000	4KU	CAW-CAQ-CAR-CAX
4	A	1000	4KU	CAR-CAQ-CAW-CAY
4	B	1000	4KU	CAQ-CAR-CAX-CAZ
4	B	1000	4KU	CAL-CAV-NAT-CAJ


There are no ring outliers.


4 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1000	4KU	2	0
4	B	1000	4KU	2	0
4	C	1000	4KU	3	0
4	A	1000	4KU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	475/911 (52%)	0.29	31 (6%)	25 15	71, 103, 169, 208	0
1	B	475/911 (52%)	0.50	39 (8%)	17 11	90, 142, 202, 249	0
1	C	475/911 (52%)	0.27	32 (6%)	24 15	79, 114, 184, 226	0
1	D	475/911 (52%)	0.66	57 (12%)	9 6	103, 151, 219, 259	0
2	E	223/223 (100%)	0.49	16 (7%)	21 14	79, 105, 150, 177	0
2	G	223/223 (100%)	0.46	18 (8%)	18 12	124, 152, 211, 250	0
2	I	223/223 (100%)	1.06	56 (25%)	1 2	79, 112, 310, 355	0
2	K	223/223 (100%)	0.51	18 (8%)	18 12	126, 152, 209, 248	0
3	F	218/218 (100%)	0.32	6 (2%)	55 32	81, 111, 141, 179	0
3	H	218/218 (100%)	0.55	17 (7%)	19 12	125, 181, 255, 268	0
3	J	218/218 (100%)	0.84	28 (12%)	7 6	90, 164, 292, 310	0
3	L	218/218 (100%)	0.44	17 (7%)	19 12	127, 169, 242, 253	0
All	All	3664/5408 (67%)	0.50	335 (9%)	15 10	71, 139, 238, 355	0

The worst 5 of 335 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	802	GLY	8.9
2	K	141	THR	7.8
2	K	140	ASP	6.5
2	G	148	LEU	6.4
3	F	194	ASN	6.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	4KU	B	1000	28/28	0.68	0.13	143,146,152,155	0
4	4KU	D	1000	28/28	0.77	0.11	148,152,159,163	0
4	4KU	A	1000	28/28	0.80	0.15	105,108,112,114	0
4	4KU	C	1000	28/28	0.84	0.12	113,118,123,125	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 4KU D 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 4KU A 1000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.