



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:10 AM UTC

PDB ID : 4YZR / pdb_00004yzr
Title : Bacillus subtilis 168 Bacillaene Polyketide Synthase (PKS) Cytochrome P450 PksS
Authors : Race, P.R.; Zulkepli, A.Z.; Anderson, R.J.L.
Deposited on : 2015-03-25
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1216 (1.36-1.36)
Clashscore	190562	1232 (1.36-1.36)
Ramachandran outliers	187476	1220 (1.36-1.36)
Sidechain outliers	187428	1220 (1.36-1.36)
RSRZ outliers	180081	1214 (1.36-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	405	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3511 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyketide biosynthesis cytochrome P450 PksS.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3155	2013	550	576	16	0	4	0

- Molecule 2 is HISTIDINE (CCD ID: HIS) (formula: $C_6H_{10}N_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	N		
2	A	1	5	3	2	0	0

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	308	Total O 308 308	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Polyketide biosynthesis cytochrome P450 PksS



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.81Å 66.69Å 74.54Å 90.00° 107.68° 90.00°	Depositor
Resolution (Å)	46.50 – 1.35 46.50 – 1.35	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.50-1.35) 99.8 (46.50-1.35)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.183 , 0.211 0.185 , 0.213	Depositor DCC
R_{free} test set	4920 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	14.0	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 29.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3511	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.77	44/3232 (1.4%)	1.41	14/4360 (0.3%)

All (44) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	31	HIS	CA-C	-8.68	1.42	1.52
1	A	391	VAL	CA-CB	8.54	1.64	1.54
1	A	241	ALA	N-CA	7.46	1.55	1.46
1	A	107	HIS	C-N	-7.38	1.24	1.34
1	A	241	ALA	CA-C	-6.82	1.44	1.52
1	A	214	SER	C-O	6.79	1.31	1.24
1	A	25	ARG	C-O	-6.76	1.16	1.24
1	A	370	LEU	C-O	6.75	1.32	1.24
1	A	393	LEU	C-O	6.72	1.31	1.24
1	A	349	ILE	CA-CB	6.60	1.62	1.54
1	A	240	VAL	CA-CB	-6.38	1.46	1.54
1	A	95	ASP	N-CA	-6.29	1.38	1.45
1	A	278	VAL	N-CA	6.27	1.53	1.46
1	A	176	VAL	N-CA	6.26	1.54	1.46
1	A	299	ASP	N-CA	-6.12	1.38	1.46
1	A	192	GLU	CA-C	6.09	1.60	1.52
1	A	259	GLN	CD-NE2	-6.08	1.20	1.33
1	A	109	ALA	N-CA	-6.04	1.38	1.46
1	A	131	ASP	CA-C	-5.97	1.45	1.52
1	A	68	ASN	N-CA	-5.96	1.38	1.46
1	A	350	HIS	CA-CB	-5.95	1.44	1.53
1	A	154	ILE	C-N	-5.90	1.26	1.33
1	A	64	ARG	C-O	-5.77	1.16	1.23
1	A	15	HIS	C-O	-5.77	1.17	1.24
1	A	308	LYS	CA-C	-5.76	1.46	1.52
1	A	119	ARG	CZ-NH1	-5.75	1.24	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	10	GLY	C-O	5.72	1.30	1.23
1	A	133	MET	SD-CE	-5.66	1.65	1.79
1	A	62	ILE	C-O	5.64	1.30	1.24
1	A	196	TYR	N-CA	-5.58	1.39	1.46
1	A	225	LYS	N-CA	5.47	1.52	1.45
1	A	125	ILE	CA-C	-5.47	1.46	1.52
1	A	167	GLN	N-CA	-5.35	1.39	1.46
1	A	7	HIS	N-CA	-5.33	1.40	1.46
1	A	164	ASP	CA-C	5.26	1.59	1.52
1	A	292	ALA	C-O	-5.23	1.17	1.24
1	A	305	HIS	CA-C	-5.22	1.46	1.52
1	A	68	ASN	CA-C	-5.19	1.45	1.52
1	A	69	VAL	C-N	-5.19	1.25	1.33
1	A	372	ARG	C-O	5.15	1.30	1.24
1	A	312	MET	C-O	-5.13	1.17	1.23
1	A	197	ILE	CB-CG1	-5.10	1.43	1.53
1	A	131	ASP	N-CA	5.03	1.52	1.46
1	A	401	VAL	N-CA	-5.01	1.40	1.46

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	67	LYS	N-CA-C	7.96	120.87	111.71
1	A	244	GLU	N-CA-C	-7.56	101.94	112.45
1	A	119	ARG	CD-NE-CZ	6.91	134.08	124.40
1	A	404	SER	O-C-N	-6.65	116.28	123.26
1	A	181	GLY	CA-C-N	-5.91	113.22	122.73
1	A	181	GLY	C-N-CA	-5.91	113.22	122.73
1	A	294	ARG	NE-CZ-NH1	-5.51	115.99	121.50
1	A	399	LEU	O-C-N	5.38	125.31	121.12
1	A	131	ASP	O-C-N	-5.30	116.50	122.12
1	A	239	VAL	CA-C-O	-5.24	115.61	121.17
1	A	180	GLU	N-CA-C	5.23	117.66	111.33
1	A	55	ALA	O-C-N	-5.05	116.87	122.07
1	A	165	ARG	NE-CZ-NH2	-5.02	114.68	119.20
1	A	164	ASP	O-C-N	5.01	128.98	122.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3155	0	3158	66	0
2	A	5	0	3	2	0
3	A	43	0	30	4	0
4	A	308	0	0	11	0
All	All	3511	0	3191	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:293:ASN:N	1:A:312:MET:CE	1.80	1.43
1:A:293:ASN:N	1:A:312:MET:HE1	1.02	1.34
1:A:21:LEU:HD21	1:A:290:MET:HE2	1.19	1.19
1:A:292:ALA:HB1	1:A:312:MET:CE	1.71	1.19
1:A:293:ASN:H	1:A:312:MET:CE	1.50	1.16
1:A:6:PHE:HZ	1:A:290:MET:HE1	1.04	1.14
1:A:6:PHE:CZ	1:A:290:MET:HE1	1.81	1.13
1:A:292:ALA:C	1:A:312:MET:HE1	1.78	1.08
1:A:292:ALA:HB1	1:A:312:MET:HE2	1.38	1.02
1:A:293:ASN:CA	1:A:312:MET:HE1	1.89	1.01
1:A:292:ALA:HB1	1:A:312:MET:HE3	1.43	0.99
1:A:293:ASN:H	1:A:312:MET:HE2	1.28	0.95
1:A:292:ALA:CB	1:A:312:MET:CE	2.44	0.95
1:A:292:ALA:C	1:A:312:MET:CE	2.40	0.88
1:A:294:ARG:O	1:A:312:MET:SD	2.34	0.85
1:A:243:LEU:O	1:A:247:VAL:HG23	1.76	0.85
1:A:292:ALA:CB	1:A:312:MET:HE2	2.09	0.81
1:A:21:LEU:HD21	1:A:290:MET:CE	2.07	0.80
1:A:13:PHE:CZ	1:A:290:MET:HE3	2.17	0.79
1:A:294:ARG:H	1:A:312:MET:CE	1.97	0.77
1:A:88:SER:OG	4:A:601:HOH:O	1.98	0.77
1:A:71:ASN:O	1:A:72:GLN:CB	2.33	0.76
1:A:363:HIS:HD2	4:A:863:HOH:O	1.68	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:ASN:HB3	4:A:605:HOH:O	1.86	0.73
1:A:241:ALA:O	1:A:245[B]:THR:HG22	1.89	0.73
1:A:292:ALA:CB	1:A:312:MET:HE3	2.12	0.73
1:A:71:ASN:O	1:A:72:GLN:HB2	1.91	0.70
1:A:13:PHE:CE1	1:A:290:MET:HE3	2.28	0.69
1:A:177:ASP:OD1	1:A:179:SER:HB2	1.93	0.68
1:A:292:ALA:CA	1:A:312:MET:CE	2.73	0.66
1:A:292:ALA:CA	1:A:312:MET:HE3	2.27	0.63
1:A:245[A]:THR:HG22	4:A:726:HOH:O	2.02	0.60
1:A:294:ARG:N	1:A:312:MET:HE1	2.17	0.58
1:A:71:ASN:O	1:A:72:GLN:CG	2.52	0.57
1:A:294:ARG:N	1:A:312:MET:SD	2.72	0.56
1:A:194:LYS:HE3	1:A:232:TYR:CE2	2.40	0.56
1:A:243:LEU:O	1:A:247:VAL:CG2	2.51	0.55
2:A:501:HIS:CG	4:A:844:HOH:O	2.59	0.55
1:A:15:HIS:HD2	4:A:874:HOH:O	1.90	0.55
1:A:69:VAL:O	1:A:71:ASN:HB2	2.06	0.54
1:A:245[B]:THR:HG23	1:A:246:THR:N	2.22	0.54
1:A:68:ASN:CB	4:A:605:HOH:O	2.52	0.54
1:A:54:MET:HE1	1:A:347:PHE:CE2	2.44	0.52
1:A:54:MET:HE1	1:A:347:PHE:CD2	2.46	0.51
1:A:13:PHE:HZ	1:A:290:MET:HE3	1.72	0.51
1:A:218:HIS:HD2	4:A:872:HOH:O	1.93	0.51
1:A:245[B]:THR:CG2	1:A:246:THR:N	2.74	0.50
1:A:265:GLU:HG3	1:A:269:GLN:HE21	1.77	0.50
1:A:241:ALA:HB1	3:A:502:HEM:C4C	2.47	0.49
1:A:294:ARG:H	1:A:312:MET:HE1	1.64	0.49
1:A:245[A]:THR:HG21	4:A:844:HOH:O	2.14	0.48
1:A:96:THR:HG23	1:A:97:PRO:HA	1.95	0.48
1:A:91:MET:CE	1:A:103:ARG:HD2	2.44	0.47
1:A:96:THR:CG2	1:A:97:PRO:HA	2.44	0.47
1:A:242:GLY:HA2	3:A:502:HEM:HMC3	1.97	0.46
1:A:293:ASN:C	1:A:312:MET:HE1	2.40	0.46
1:A:21:LEU:CD2	1:A:290:MET:HE2	2.14	0.45
1:A:91:MET:HE1	1:A:103:ARG:HD2	1.98	0.44
1:A:119:ARG:HH12	1:A:363:HIS:CE1	2.36	0.43
1:A:354:GLY:HA3	3:A:502:HEM:C3C	2.53	0.43
1:A:54:MET:HE1	1:A:347:PHE:CZ	2.55	0.42
1:A:149:LEU:HB3	1:A:150:PRO:HD3	2.02	0.42
1:A:116:GLU:OE2	1:A:119:ARG:NH2	2.47	0.42
1:A:143:LYS:HE3	4:A:626:HOH:O	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:285:THR:O	1:A:286:SER:C	2.64	0.41
3:A:502:HEM:CMB	3:A:502:HEM:HBB2	2.51	0.41
1:A:245[B]:THR:HG21	2:A:501:HIS:HD2	1.86	0.40
1:A:207:LYS:HG3	4:A:797:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/405 (95%)	374 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	347/361 (96%)	347 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	HIS
1	A	169	GLN
1	A	187	ASN
1	A	218	HIS
1	A	269	GLN
1	A	293	ASN
1	A	305	HIS
1	A	363	HIS
1	A	386	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HIS	A	501	3	5,5,11	0.56	0	5,5,14	0.79	0
3	HEM	A	502	2,1	50,50,50	1.65	8 (16%)	67,82,82	1.68	13 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	502	2,1	-	1/14/54/54	-
2	HIS	A	501	3	-	-	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	HEM	O2A-CGA	-3.91	1.17	1.30
3	A	502	HEM	CBA-CGA	3.80	1.59	1.50
3	A	502	HEM	FE-NC	3.70	2.07	1.95
3	A	502	HEM	FE-NA	3.49	2.06	1.95
3	A	502	HEM	FE-NB	3.36	2.05	1.94
3	A	502	HEM	C2A-C3A	-2.89	1.31	1.38
3	A	502	HEM	CBD-CGD	2.54	1.56	1.50
3	A	502	HEM	CMB-C2B	-2.16	1.46	1.50

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	HEM	C3B-C2B-C1B	5.57	110.60	106.41
3	A	502	HEM	O2A-CGA-O1A	3.90	133.36	123.33
3	A	502	HEM	O1A-CGA-CBA	-3.34	112.50	123.09
3	A	502	HEM	C2A-C1A-NA	-3.24	106.56	110.15
3	A	502	HEM	CHB-C4A-NA	3.10	129.49	123.86
3	A	502	HEM	C1B-NB-C4B	3.05	108.81	105.21
3	A	502	HEM	C4B-C3B-C2B	-3.02	104.50	107.28
3	A	502	HEM	C2B-C1B-NB	-2.64	106.81	109.84
3	A	502	HEM	O1D-CGD-CBD	-2.29	115.83	123.09
3	A	502	HEM	CMB-C2B-C1B	-2.19	121.61	125.03
3	A	502	HEM	C4A-NA-C1A	2.14	109.31	105.82
3	A	502	HEM	C2D-C1D-ND	2.09	112.32	109.90
3	A	502	HEM	CHC-C4B-NB	2.01	126.58	124.42

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	502	HEM	CAA-CBA-CGA-O2A

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	HIS	2	0
3	A	502	HEM	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/405 (95%)	0.62	36 (9%) 14 20	6, 16, 38, 67	4 (1%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	40	ALA	6.9
1	A	181	GLY	5.5
1	A	180	GLU	4.9
1	A	312	MET	4.8
1	A	84	ILE	4.6
1	A	206	ILE	4.5
1	A	71	ASN	4.5
1	A	178	THR	4.2
1	A	405	LYS	3.7
1	A	5	MET	3.7
1	A	179	SER	3.7
1	A	66	VAL	3.6
1	A	243	LEU	3.6
1	A	112	PRO	3.5
1	A	36	LYS	3.4
1	A	210	ASP	3.4
1	A	42	TYR	3.3
1	A	183	ARG	3.3
1	A	72	GLN	3.2
1	A	185	LEU	3.1
1	A	41	THR	2.9
1	A	35	LEU	2.9
1	A	83	ASP	2.9
1	A	207	LYS	2.8
1	A	349	ILE	2.7
1	A	68	ASN	2.6
1	A	85	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	182	ASN	2.6
1	A	391	VAL	2.5
1	A	339	PRO	2.5
1	A	184	GLU	2.4
1	A	70	MET	2.4
1	A	110	PHE	2.4
1	A	269	GLN	2.2
1	A	162	LYS	2.2
1	A	177	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	HIS	A	501	5/11	0.91	0.10	15,15,18,21	0
3	HEM	A	502	43/43	0.99	0.05	6,8,11,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.