



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 07:37 AM UTC

PDB ID : 6YZD / pdb_00006yzd
Title : Crystal structure of the M295A variant of Ssl1
Authors : Mielenbrink, S.; Olbrich, A.; Urlacher, V.; Span, I.
Deposited on : 2020-05-06
Resolution : 1.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4041 (1.44-1.40)
Clashscore	190562	4154 (1.44-1.40)
Ramachandran outliers	187476	4083 (1.44-1.40)
Sidechain outliers	187428	4082 (1.44-1.40)
RSRZ outliers	180081	4039 (1.44-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	325	3% 78% 5% 17%
1	BBB	325	5% 78% 5% 17%
1	CCC	325	5% 77% 7% 17%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12574 atoms, of which 6001 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Copper oxidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	AAA	270	Total	C	H	N	O	S	106	0	0
			4087	1307	1994	383	394	9			
1	BBB	270	Total	C	H	N	O	S	106	1	0
			4107	1312	2006	386	394	9			
1	CCC	271	Total	C	H	N	O	S	105	0	0
			4101	1312	2001	384	395	9			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	1	MET	-	initiating methionine	UNP B5HSR1
AAA	2	HIS	-	expression tag	UNP B5HSR1
AAA	3	HIS	-	expression tag	UNP B5HSR1
AAA	4	HIS	-	expression tag	UNP B5HSR1
AAA	5	HIS	-	expression tag	UNP B5HSR1
AAA	6	HIS	-	expression tag	UNP B5HSR1
AAA	7	HIS	-	expression tag	UNP B5HSR1
AAA	295	ALA	MET	engineered mutation	UNP B5HSR1
BBB	1	MET	-	initiating methionine	UNP B5HSR1
BBB	2	HIS	-	expression tag	UNP B5HSR1
BBB	3	HIS	-	expression tag	UNP B5HSR1
BBB	4	HIS	-	expression tag	UNP B5HSR1
BBB	5	HIS	-	expression tag	UNP B5HSR1
BBB	6	HIS	-	expression tag	UNP B5HSR1
BBB	7	HIS	-	expression tag	UNP B5HSR1
BBB	295	ALA	MET	engineered mutation	UNP B5HSR1
CCC	1	MET	-	initiating methionine	UNP B5HSR1
CCC	2	HIS	-	expression tag	UNP B5HSR1
CCC	3	HIS	-	expression tag	UNP B5HSR1
CCC	4	HIS	-	expression tag	UNP B5HSR1
CCC	5	HIS	-	expression tag	UNP B5HSR1
CCC	6	HIS	-	expression tag	UNP B5HSR1
CCC	7	HIS	-	expression tag	UNP B5HSR1

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
CCC	295	ALA	MET	engineered mutation	UNP B5HSR1

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	AAA	5	Total Cu 5 5	0	0
2	BBB	5	Total Cu 5 5	0	0
2	CCC	4	Total Cu 4 4	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	1	Total O S 5 4 1	0	0
3	CCC	1	Total O S 5 4 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AAA	103	Total O 103 103	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	BBB	75	Total 75	O 75	0	0
4	CCC	77	Total 77	O 77	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Copper oxidase



- Molecule 1: Copper oxidase



- Molecule 1: Copper oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.44Å 104.10Å 163.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.36 – 1.41 30.36 – 1.41	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.36-1.41) 99.8 (30.36-1.41)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 1.41Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.200 , 0.225 0.207 , 0.231	Depositor DCC
R_{free} test set	8343 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.237	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 23.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12574	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.05	1/2151 (0.0%)	1.12	3/2919 (0.1%)
1	BBB	1.08	0/2162	1.16	1/2933 (0.0%)
1	CCC	1.08	1/2159 (0.0%)	1.18	3/2931 (0.1%)
All	All	1.07	2/6472 (0.0%)	1.15	7/8783 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	CCC	311	TYR	C-O	5.27	1.30	1.23
1	AAA	254	VAL	C-O	5.00	1.30	1.24

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	161	HIS	CA-CB-CG	-6.68	107.12	113.80
1	BBB	161	HIS	CA-CB-CG	-5.58	108.22	113.80
1	CCC	139	ASP	CA-CB-CG	5.43	118.03	112.60
1	AAA	250	ASP	CB-CA-C	5.16	115.28	110.17
1	CCC	57	ASP	N-CA-C	-5.15	106.95	113.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2093	1994	1974	6	0
1	BBB	2101	2006	1987	7	0
1	CCC	2100	2001	1981	11	0
2	AAA	5	0	0	0	0
2	BBB	5	0	0	0	0
2	CCC	4	0	0	0	0
3	AAA	5	0	0	0	0
3	CCC	5	0	0	0	0
4	AAA	103	0	0	0	0
4	BBB	75	0	0	0	0
4	CCC	77	0	0	1	0
All	All	6573	6001	5942	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 23 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:CCC:59:GLN:HE22	1:CCC:199:ASN:HB3	1.44	0.82
1:AAA:59:GLN:HE22	1:AAA:199:ASN:HB3	1.55	0.71
1:CCC:44:GLU:OE1	1:CCC:46:ARG:NH2	2.30	0.64
1:CCC:244:MET:HE2	4:CCC:508:HOH:O	1.99	0.61
1:AAA:53:GLU:HG2	1:AAA:91:THR:OG1	2.02	0.60

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	268/325 (82%)	264 (98%)	4 (2%)	0	100	100
1	BBB	269/325 (83%)	263 (98%)	6 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	CCC	269/325 (83%)	264 (98%)	5 (2%)	0	100	100
All	All	806/975 (83%)	791 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	216/255 (85%)	214 (99%)	2 (1%)	70	43
1	BBB	217/255 (85%)	216 (100%)	1 (0%)	81	62
1	CCC	217/255 (85%)	215 (99%)	2 (1%)	70	43
All	All	650/765 (85%)	645 (99%)	5 (1%)	73	47

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	220	MET
1	AAA	236	ARG
1	BBB	220	MET
1	CCC	220	MET
1	CCC	236	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 14 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	CCC	405	-	4,4,4	0.32	0	6,6,6	0.16	0
3	SO4	AAA	406	-	4,4,4	0.20	0	6,6,6	0.29	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	AAA	270/325 (83%)	-0.13	9 (3%)	49	52	11, 18, 32, 64	14 (5%)
1	BBB	270/325 (83%)	0.31	17 (6%)	26	28	13, 22, 46, 76	15 (5%)
1	CCC	271/325 (83%)	0.31	15 (5%)	30	33	13, 22, 44, 60	14 (5%)
All	All	811/975 (83%)	0.16	41 (5%)	33	36	11, 20, 42, 76	43 (5%)

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	CCC	56	ALA	6.9
1	CCC	57	ASP	5.5
1	CCC	313	PRO	4.4
1	CCC	203	HIS	4.2
1	AAA	312	GLU	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CU	BBB	405	1/1	0.74	0.18	92,92,92,92	0
3	SO4	CCC	405	5/5	0.92	0.09	41,44,46,57	0
3	SO4	AAA	406	5/5	0.93	0.10	28,30,37,38	0
2	CU	AAA	405	1/1	0.93	0.19	60,60,60,60	0
2	CU	AAA	404	1/1	0.97	0.05	22,22,22,22	0
2	CU	AAA	402	1/1	0.97	0.05	22,22,22,22	0
2	CU	BBB	402	1/1	0.97	0.06	22,22,22,22	0
2	CU	CCC	403	1/1	0.98	0.04	21,21,21,21	0
2	CU	BBB	403	1/1	0.98	0.04	27,27,27,27	0
2	CU	CCC	402	1/1	0.98	0.06	24,24,24,24	0
2	CU	BBB	404	1/1	0.99	0.20	34,34,34,34	0
2	CU	AAA	403	1/1	0.99	0.22	30,30,30,30	0
2	CU	BBB	401	1/1	0.99	0.03	23,23,23,23	0
2	CU	CCC	404	1/1	1.00	0.17	28,28,28,28	0
2	CU	AAA	401	1/1	1.00	0.02	19,19,19,19	0
2	CU	CCC	401	1/1	1.00	0.02	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CU BBB 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU AAA 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU AAA 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU AAA 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU BBB 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU CCC 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU BBB 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU CCC 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU BBB 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU AAA 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU BBB 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around CU CCC 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CU AAA 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CU CCC 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.