



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:31 AM UTC

PDB ID : 7YZ0 / pdb_00007yz0
Title : Molecular snapshots of drug release from tubulin: 100 microseconds after photoactivation.
Authors : Wranik, M.; Weinert, T.; Standfuss, J.
Deposited on : 2022-02-18
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	
2	B	445	
3	F	169	

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 8262 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	437	Total	C	N	O	S	0	2	0
			3412	2161	579	648	24			


- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	426	Total	C	N	O	S	0	2	0
			3346	2101	570	647	28			

- Molecule 3 is a protein called Designed Ankyrin Repeat Protein (DARPIN) D1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	155	Total	C	N	O	S	0	0	0
			1154	727	198	226	3			

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		


- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 8 is Azo-Combretastatin A4 (trans) (CCD ID: VYT) (formula: C₁₆H₁₈N₂O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	C	N	O	0	0
			23	16	2	5		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	127	Total	O	0	0
			127	127		
9	B	94	Total	O	0	0
			94	94		
9	F	44	Total	O	0	0
			44	44		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Tubulin alpha-1B chain




• Molecule 2: Tubulin beta-2B chain



• Molecule 3: Designed Ankyrin Repeat Protein (DARPIN) D1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.53Å 92.58Å 83.99Å 90.00° 96.71° 90.00°	Depositor
Resolution (Å)	9.49 – 2.20 9.49 – 2.20	Depositor EDS
% Data completeness (in resolution range)	92.5 (9.49-2.20) 91.3 (9.49-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.66 (at 2.21Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.305 , 0.363 0.305 , 0.363	Depositor DCC
R_{free} test set	1841 reflections (3.50%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.253	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 77.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.55$, $\langle L^2 \rangle = 0.39$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	8262	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GDP, GTP, VYT, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.07	0/3496	0.23	0/4748
2	B	0.07	0/3428	0.23	0/4641
3	F	0.06	0/1170	0.18	0/1590
All	All	0.07	0/8094	0.22	0/10979

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3412	0	3318	29	0
2	B	3346	0	3216	40	0
3	F	1154	0	1156	13	0
4	A	32	0	12	1	0
5	A	1	0	0	0	0
6	A	1	0	0	0	0
7	B	28	0	12	0	0
8	B	23	0	0	1	0
9	A	127	0	0	4	0
9	B	94	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	F	44	0	0	0	0
All	All	8262	0	7714	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:268:PRO:HG2	2:B:300:MET:HB2	1.69	0.73
2:B:161:ASP:O	2:B:251:ARG:NH2	2.24	0.70
2:B:293:MET:HG2	2:B:367:PHE:HB2	1.74	0.69
2:B:48:ASN:OD1	2:B:48:ASN:N	2.26	0.68
1:A:172:TYR:HB3	1:A:205:ASP:HA	1.75	0.67
2:B:396:HIS:NE2	3:F:57:TYR:OH	2.28	0.67
2:B:52:ASN:HD22	2:B:86:ARG:HH12	1.45	0.64
1:A:107:HIS:NE2	9:A:601:HOH:O	2.29	0.63
2:B:2:ARG:NH2	2:B:128:ASP:OD2	2.32	0.62
2:B:102:ALA:HB2	2:B:403:MET:HE2	1.80	0.62
2:B:145:SER:OG	2:B:188:SER:OG	2.17	0.62
1:A:2:ARG:O	1:A:133:GLN:NE2	2.36	0.58
2:B:165:ASN:ND2	9:B:617:HOH:O	2.37	0.57
3:F:79:MET:HE3	3:F:111:THR:HG21	1.88	0.56
3:F:121:ALA:HB1	3:F:161:LEU:HD21	1.89	0.55
1:A:26:LEU:HD12	1:A:363:VAL:HG12	1.89	0.55
2:B:398:TYR:HB3	2:B:403:MET:HE3	1.88	0.55
1:A:123:ARG:NH1	9:A:608:HOH:O	2.35	0.54
2:B:36:TYR:OH	2:B:40:SER:O	2.25	0.54
3:F:27:ASP:N	3:F:27:ASP:OD1	2.39	0.53
2:B:121:ARG:NE	2:B:158:GLU:OE1	2.42	0.52
1:A:293:ASN:O	1:A:339:ARG:NH1	2.43	0.52
2:B:251:ARG:NH1	9:B:623:HOH:O	2.43	0.52
1:A:370:LYS:NZ	9:A:610:HOH:O	2.36	0.51
2:B:31:ASP:OD1	2:B:35:SER:N	2.42	0.51
2:B:1:MET:HA	2:B:131:GLN:HB3	1.94	0.50
2:B:382:SER:HB2	2:B:415:MET:HE3	1.93	0.50
3:F:60:LEU:HD11	3:F:98:VAL:HG21	1.92	0.50
1:A:62:VAL:HG11	1:A:88:HIS:CD2	2.47	0.50
1:A:433:GLU:HA	1:A:437:VAL:HG12	1.94	0.49
3:F:133:LEU:HD11	3:F:165:LEU:HD23	1.94	0.49
1:A:215:ARG:NH1	9:A:617:HOH:O	2.45	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:ARG:NH2	1:A:299:ALA:O	2.44	0.49
1:A:93:ILE:HG22	1:A:114:ILE:HD11	1.94	0.49
2:B:45:GLU:O	2:B:46[B]:ARG:HG2	2.13	0.49
2:B:211:CYS:HB3	2:B:217:LEU:HD12	1.95	0.48
3:F:23:ARG:NH1	3:F:57:TYR:OH	2.46	0.48
1:A:15:GLN:NE2	4:A:501:GTP:O6	2.46	0.48
2:B:253:LEU:HD13	8:B:502:VYT:C6	2.42	0.48
2:B:45:GLU:O	2:B:46[C]:ARG:HG2	2.13	0.48
2:B:323:MET:HE1	2:B:353:VAL:HG11	1.96	0.48
2:B:143:THR:HG23	2:B:147:MET:HE2	1.96	0.47
2:B:28:HIS:CE1	2:B:241:ARG:HB3	2.50	0.47
2:B:61:PRO:HD2	2:B:84:ILE:O	2.15	0.47
3:F:18:LEU:HD11	3:F:30:VAL:HG13	1.98	0.46
1:A:413:MET:HE3	1:A:417:GLU:HB3	1.97	0.46
1:A:188:ILE:HG23	1:A:425:MET:HG3	1.97	0.46
1:A:397:LEU:HD22	2:B:430:ALA:HB1	1.99	0.45
3:F:40:VAL:HG13	3:F:66:LEU:HD22	1.98	0.45
2:B:375:GLN:HB2	2:B:419:VAL:HG13	1.98	0.44
2:B:396:HIS:CE1	3:F:57:TYR:HH	2.36	0.44
2:B:290:THR:HA	2:B:293:MET:HE3	2.00	0.44
1:A:21:TRP:CZ3	1:A:63:PRO:HB3	2.53	0.43
1:A:292:THR:HG22	1:A:335:ILE:HD12	2.01	0.43
1:A:167:LEU:HG	1:A:200:CYS:HB3	2.00	0.43
2:B:301:ALA:O	2:B:303:CYS:N	2.51	0.43
3:F:164:ILE:O	3:F:167:LYS:HG3	2.17	0.43
1:A:221:ARG:CD	2:B:323:MET:HB3	2.49	0.43
1:A:215:ARG:NH1	1:A:216:ASN:OD1	2.51	0.43
1:A:139:HIS:CD2	1:A:150:THR:HG21	2.52	0.43
1:A:147:SER:HB2	1:A:190:THR:HB	2.00	0.43
2:B:202:ILE:HD13	2:B:229:VAL:HG13	2.02	0.42
1:A:167:LEU:HD13	1:A:252:LEU:HD22	2.02	0.42
2:B:239:CYS:O	2:B:248:ALA:HB2	2.20	0.41
1:A:75:ILE:HD12	1:A:94:THR:HG22	2.02	0.41
1:A:141:PHE:HD1	1:A:141:PHE:HA	1.76	0.41
2:B:286:VAL:HB	2:B:287:PRO:HD3	2.02	0.41
2:B:293:MET:HE2	2:B:365:ALA:HB1	2.02	0.41
1:A:200:CYS:HA	1:A:266:HIS:HB2	2.01	0.41
2:B:15:GLN:NE2	9:B:618:HOH:O	2.37	0.41
2:B:206:ALA:HB2	2:B:302:ALA:N	2.36	0.41
1:A:221:ARG:HD2	2:B:323:MET:HB3	2.03	0.41
1:A:296:PHE:HB2	1:A:339:ARG:NH1	2.36	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:293:MET:HE1	2:B:317:PHE:CZ	2.57	0.40
2:B:178:THR:HB	2:B:181:GLU:HG3	2.02	0.40
3:F:43:THR:HG22	3:F:49:THR:HG22	2.03	0.40
3:F:23:ARG:HG3	3:F:57:TYR:CE1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	437/451 (97%)	420 (96%)	15 (3%)	2 (0%)	24	27
2	B	424/445 (95%)	409 (96%)	12 (3%)	3 (1%)	18	19
3	F	153/169 (90%)	151 (99%)	2 (1%)	0	100	100
All	All	1014/1065 (95%)	980 (97%)	29 (3%)	5 (0%)	24	27

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	243	PRO
2	B	275	SER
1	A	322	ASP
1	A	349	THR
2	B	248	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/379 (97%)	366 (100%)	1 (0%)	86	93
2	B	368/383 (96%)	359 (98%)	9 (2%)	43	58
3	F	120/132 (91%)	118 (98%)	2 (2%)	53	69
All	All	855/894 (96%)	843 (99%)	12 (1%)	61	75

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	GLU
2	B	42	LEU
2	B	46[B]	ARG
2	B	46[C]	ARG
2	B	48	ASN
2	B	111	GLU
2	B	121	ARG
2	B	221	THR
2	B	246	LEU
2	B	288	GLU
3	F	27	ASP
3	F	144	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	ASN
2	B	11	GLN
2	B	52	ASN
2	B	99	ASN
2	B	329	GLN
2	B	335	ASN
2	B	337	ASN
2	B	347	ASN
2	B	375	GLN
2	B	426	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	GDP	B	501	-	29,30,30	1.15	3 (10%)	45,47,47	1.78	6 (13%)
8	VYT	B	502	-	24,24,24	0.22	0	32,32,32	0.21	0
4	GTP	A	501	5	33,34,34	1.03	4 (12%)	50,54,54	1.57	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GDP	B	501	-	-	4/16/32/32	0/3/3/3
8	VYT	B	502	-	-	0/13/13/13	0/2/2/2
4	GTP	A	501	5	-	7/22/38/38	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	501	GDP	C5-C4	3.15	1.47	1.38
7	B	501	GDP	C6-N1	-2.38	1.34	1.38
4	A	501	GTP	PA-O3A	2.38	1.62	1.59
4	A	501	GTP	C2-N3	2.19	1.38	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	GTP	PB-O3A	2.14	1.61	1.59
4	A	501	GTP	PB-O3B	2.05	1.61	1.59
7	B	501	GDP	C5-N7	-2.04	1.35	1.39

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	501	GDP	C5-C4-N3	-6.07	118.72	128.39
7	B	501	GDP	C2-N3-C4	5.03	120.96	112.30
4	A	501	GTP	C5-C4-N3	-4.94	120.52	128.39
4	A	501	GTP	C2-N3-C4	4.60	120.23	112.30
7	B	501	GDP	N9-C4-N3	4.56	135.06	125.95
7	B	501	GDP	C6-C5-N7	3.23	136.16	130.29
4	A	501	GTP	N9-C4-N3	3.15	132.25	125.95
4	A	501	GTP	C2-N1-C6	-2.88	119.89	125.11
4	A	501	GTP	N9-C8-N7	-2.70	108.39	113.40
4	A	501	GTP	C8-N7-C5	2.55	108.81	104.26
4	A	501	GTP	C5-C6-N1	2.51	119.63	113.25
7	B	501	GDP	C4-C5-N7	-2.51	106.70	110.67
4	A	501	GTP	O6-C6-C5	-2.40	120.19	126.53
7	B	501	GDP	O6-C6-C5	-2.13	120.91	126.53

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	501	GTP	PB-O3B-PG-O2G
4	A	501	GTP	C5'-O5'-PA-O3A
4	A	501	GTP	C5'-O5'-PA-O1A
4	A	501	GTP	C5'-O5'-PA-O2A
7	B	501	GDP	C5'-O5'-PA-O3A
7	B	501	GDP	C5'-O5'-PA-O1A
7	B	501	GDP	C5'-O5'-PA-O2A
4	A	501	GTP	PB-O3A-PA-O1A
4	A	501	GTP	C4'-C5'-O5'-PA
7	B	501	GDP	PB-O3A-PA-O2A
4	A	501	GTP	PB-O3A-PA-O2A


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	502	VYT	1	0
4	A	501	GTP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	437/451 (96%)	0.76	33 (7%)	20 17	14, 28, 48, 77	2 (0%)
2	B	426/445 (95%)	1.28	82 (19%)	3 2	20, 39, 66, 77	2 (0%)
3	F	155/169 (91%)	0.76	12 (7%)	19 17	20, 30, 47, 66	0
All	All	1018/1065 (95%)	0.98	127 (12%)	8 6	14, 32, 61, 77	4 (0%)

All (127) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	73	MET	6.4
1	A	339	ARG	5.7
2	B	127	CYS	5.3
2	B	248	ALA	5.1
2	B	319	GLY	5.1
2	B	29	GLY	4.5
2	B	32	PRO	4.5
1	A	437	VAL	4.1
2	B	79	GLY	4.1
1	A	335	ILE	4.1
1	A	341	ILE	4.0
2	B	93	GLY	3.9
2	B	322	SER	3.8
2	B	126	SER	3.7
2	B	95	SER	3.7
3	F	167	LYS	3.7
2	B	82	GLY	3.6
2	B	54	ALA	3.6
2	B	244	GLY	3.6
2	B	114	ASP	3.6
2	B	397	TRP	3.6
2	B	36	TYR	3.6
2	B	80	PRO	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	110	ALA	3.6
3	F	162	ALA	3.4
2	B	75	SER	3.4
1	A	159	VAL	3.4
2	B	335	ASN	3.3
1	A	236	SER	3.3
2	B	46[B]	ARG	3.3
2	B	92	PHE	3.3
2	B	274	THR	3.2
2	B	284	LEU	3.2
1	A	337	THR	3.2
1	A	43	GLY	3.1
3	F	65	VAL	3.1
2	B	409	THR	3.1
2	B	277	GLY	3.1
1	A	2	ARG	3.0
2	B	41	ASP	3.0
1	A	37	PRO	3.0
2	B	84	ILE	2.9
2	B	323	MET	2.9
2	B	58	LYS	2.8
2	B	243	PRO	2.8
1	A	338	LYS	2.8
2	B	57	ASN	2.8
1	A	62	VAL	2.8
1	A	41	THR	2.8
1	A	42	ILE	2.8
1	A	85	GLN	2.8
1	A	336	LYS	2.8
2	B	130	LEU	2.7
2	B	365	ALA	2.7
1	A	430	LYS	2.6
3	F	32	ILE	2.6
1	A	309	HIS	2.6
2	B	118	ASP	2.6
1	A	347	CYS	2.6
2	B	14	ASN	2.6
3	F	13	ASP	2.6
2	B	246	LEU	2.5
3	F	28	ASP	2.5
2	B	40	SER	2.5
2	B	286	VAL	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	396	HIS	2.5
2	B	74	ASP	2.5
2	B	34	GLY	2.5
2	B	238	THR	2.5
2	B	212	PHE	2.5
1	A	282	TYR	2.5
1	A	334	THR	2.5
2	B	219	THR	2.5
2	B	218	THR	2.4
2	B	91	VAL	2.4
2	B	33	THR	2.4
2	B	72	THR	2.4
1	A	344	VAL	2.4
2	B	389	PHE	2.4
1	A	82	THR	2.4
2	B	429	THR	2.4
2	B	112	LEU	2.4
1	A	308	ARG	2.4
2	B	12	CYS	2.3
3	F	31	ARG	2.3
2	B	111	GLU	2.3
2	B	298	ASN	2.3
2	B	38	GLY	2.3
1	A	84	ARG	2.3
2	B	83	GLN	2.3
2	B	331	LEU	2.3
2	B	271	ALA	2.3
1	A	188	ILE	2.3
2	B	253	LEU	2.2
1	A	331	ALA	2.2
1	A	409	VAL	2.2
2	B	217	LEU	2.2
3	F	145	PHE	2.2
2	B	89	ASN	2.2
2	B	414	ASN	2.2
2	B	431	ASP	2.2
1	A	278	ALA	2.2
2	B	247	ASN	2.2
2	B	87	PRO	2.2
1	A	256	GLN	2.2
2	B	291	GLN	2.2
2	B	242	PHE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	285	THR	2.2
1	A	221	ARG	2.1
2	B	101	TRP	2.1
2	B	333	VAL	2.1
2	B	353	VAL	2.1
2	B	239	CYS	2.1
2	B	278	SER	2.1
2	B	222	TYR	2.1
3	F	166	GLN	2.1
2	B	357	PRO	2.1
1	A	17	GLY	2.1
1	A	436	GLY	2.1
2	B	276	ARG	2.1
2	B	78	SER	2.1
3	F	156	ASN	2.1
2	B	326	VAL	2.1
2	B	35	SER	2.0
3	F	146	GLY	2.0
3	F	150	PHE	2.0
2	B	417	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	VYT	B	502	23/23	0.73	0.16	36,45,59,61	0
7	GDP	B	501	28/28	0.90	0.10	23,31,35,37	0
4	GTP	A	501	32/32	0.90	0.11	13,20,26,33	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	CA	A	503	1/1	0.92	0.05	32,32,32,32	0
5	MG	A	502	1/1	0.96	0.20	5,5,5,5	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.