



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 02:56 AM UTC

PDB ID : 1Z0E / pdb_00001z0e
Title : Crystal Structure of A. fulgidus Lon proteolytic domain
Authors : Botos, I.; Melnikov, E.E.; Cherry, S.; Kozlov, S.; Makhovskaya, O.V.; Tropea, J.E.; Gustchina, A.; Rotanova, T.V.; Wlodawer, A.
Deposited on : 2005-03-01
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	
1	B	205	
1	C	205	
1	D	205	
1	E	205	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	205	% 51%26%12%6%5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9285 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative protease La homolog type.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	195	Total	C	N	O	S	0	0	0
			1457	921	247	284	5			
1	B	195	Total	C	N	O	S	0	0	0
			1457	921	247	284	5			
1	C	194	Total	C	N	O	S	0	0	0
			1448	915	245	283	5			
1	D	195	Total	C	N	O	S	0	0	0
			1457	921	247	284	5			
1	E	195	Total	C	N	O	S	0	0	0
			1457	921	247	284	5			
1	F	195	Total	C	N	O	S	0	0	0
			1457	921	247	284	5			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	89	Total	O	0	0
			89	89		
2	B	65	Total	O	0	0
			65	65		
2	C	121	Total	O	0	0
			121	121		
2	D	106	Total	O	0	0
			106	106		
2	E	67	Total	O	0	0
			67	67		
2	F	104	Total	O	0	0
			104	104		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative protease La homolog type



- Molecule 1: Putative protease La homolog type




- Molecule 1: Putative protease La homolog type







- Molecule 1: Putative protease La homolog type



- Molecule 1: Putative protease La homolog type



- Molecule 1: Putative protease La homolog type



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.68Å 86.11Å 135.61Å 90.00° 94.70° 90.00°	Depositor
Resolution (Å)	15.00 – 2.05 15.00 – 2.05	Depositor EDS
% Data completeness (in resolution range)	100.0 (15.00-2.05) 96.5 (15.00-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.213 , 0.299 0.211 , 0.300	Depositor DCC
R_{free} test set	1009 reflections (1.49%)	wwPDB-VP
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9285	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.70	12/1470 (0.8%)	1.59	21/1983 (1.1%)
1	B	1.48	9/1470 (0.6%)	1.42	15/1983 (0.8%)
1	C	1.83	25/1461 (1.7%)	1.61	20/1972 (1.0%)
1	D	1.77	22/1470 (1.5%)	1.57	23/1983 (1.2%)
1	E	1.50	7/1470 (0.5%)	1.45	19/1983 (1.0%)
1	F	1.71	19/1470 (1.3%)	1.63	25/1983 (1.3%)
All	All	1.67	94/8811 (1.1%)	1.55	123/11887 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	F	0	2
All	All	0	3

The worst 5 of 94 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	496	ILE	CA-CB	10.61	1.67	1.54
1	F	556	ALA	CA-CB	9.08	1.67	1.53
1	C	566	ILE	C-O	-8.64	1.17	1.24
1	F	595	VAL	CA-CB	8.41	1.63	1.54
1	D	542	GLU	CB-CG	8.23	1.77	1.52

The worst 5 of 123 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	446	ILE	CB-CA-C	-11.02	93.86	110.82
1	F	441	ILE	CB-CA-C	-10.51	96.58	111.40
1	D	452	SER	N-CA-C	8.09	122.42	112.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	483	LYS	N-CA-C	8.00	127.83	110.80
1	F	446	ILE	CB-CA-C	-7.93	98.81	110.62

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	438	SER	Peptide
1	F	579	GLU	Peptide
1	F	580	HIS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1457	0	1523	78	0
1	B	1457	0	1523	91	0
1	C	1448	0	1510	128	1
1	D	1457	0	1523	72	2
1	E	1457	0	1523	55	0
1	F	1457	0	1523	96	1
2	A	89	0	0	25	0
2	B	65	0	0	17	0
2	C	121	0	0	35	1
2	D	106	0	0	23	0
2	E	67	0	0	17	0
2	F	104	0	0	23	0
All	All	9285	0	9125	499	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 499 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:542:GLU:CG	1:D:542:GLU:CB	1.77	1.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:480:ILE:HG21	2:A:691:HOH:O	1.11	1.27
1:B:480:ILE:HG21	2:B:682:HOH:O	1.37	1.24
1:C:418:LEU:N	2:C:725:HOH:O	1.62	1.23
1:C:502:TYR:HB2	2:C:726:HOH:O	1.12	1.22

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:486:GLY:O	1:F:578:ALA:CB[1_455]	1.82	0.38
1:C:585:GLU:OE2	1:D:597:GLU:OE1[2_546]	2.13	0.07
1:D:568:LYS:CD	2:C:738:HOH:O[2_556]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	191/205 (93%)	183 (96%)	8 (4%)	0	100	100
1	B	191/205 (93%)	181 (95%)	5 (3%)	5 (3%)	4	0
1	C	190/205 (93%)	179 (94%)	5 (3%)	6 (3%)	3	0
1	D	191/205 (93%)	183 (96%)	6 (3%)	2 (1%)	12	6
1	E	191/205 (93%)	181 (95%)	9 (5%)	1 (0%)	24	16
1	F	191/205 (93%)	177 (93%)	6 (3%)	8 (4%)	2	0
All	All	1145/1230 (93%)	1084 (95%)	39 (3%)	22 (2%)	6	2

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	436	GLY
1	B	452	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	579	GLU
1	F	476	ASN
1	F	477	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/168 (95%)	138 (86%)	22 (14%)	3	1
1	B	160/168 (95%)	138 (86%)	22 (14%)	3	1
1	C	159/168 (95%)	130 (82%)	29 (18%)	2	0
1	D	160/168 (95%)	139 (87%)	21 (13%)	4	1
1	E	160/168 (95%)	133 (83%)	27 (17%)	2	0
1	F	160/168 (95%)	135 (84%)	25 (16%)	2	0
All	All	959/1008 (95%)	813 (85%)	146 (15%)	3	0

5 of 146 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	575	LEU
1	F	608	ARG
1	E	604	LYS
1	F	485	THR
1	C	437	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	491	ASN
1	A	551	GLN
1	B	593	ASN
1	E	607	ASN
1	F	558	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	195/205 (95%)	-0.12	1 (0%) 87 89	26, 33, 52, 60	0
1	B	195/205 (95%)	0.28	7 (3%) 46 46	29, 39, 60, 78	0
1	C	194/205 (94%)	-0.21	5 (2%) 57 59	23, 30, 54, 65	0
1	D	195/205 (95%)	-0.24	1 (0%) 87 89	24, 31, 48, 60	0
1	E	195/205 (95%)	0.40	9 (4%) 37 37	30, 38, 57, 77	0
1	F	195/205 (95%)	-0.10	3 (1%) 72 73	26, 33, 56, 60	0
All	All	1169/1230 (95%)	0.00	26 (2%) 62 64	23, 35, 56, 78	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	502	TYR	4.8
1	B	418	LEU	4.2
1	E	418	LEU	4.1
1	E	419	PHE	3.8
1	B	420	ILE	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.