



wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 12:46 PM UTC

PDB ID : 7Z10 / pdb_00007z10
EMDB ID : EMD-14436
Title : Monomeric respiratory complex IV isolated from *S. cerevisiae*
Authors : Marechal, A.; Hartley, A.; Ing, G.; Pinotsis, N.
Deposited on : 2022-02-24
Resolution : 3.87 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	8831 (3.37 - 4.37)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	a	534	 16% 74% 23% .
2	b	236	 14% 71% 27% .
3	c	269	 14% 80% 19% .
4	d	121	 16% 75% 23% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	e	133	
6	f	104	
7	g	59	
8	h	47	
9	i	55	

2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 12574 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c oxidase subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	a	534	Total	C	N	O	S	0	0
			4162	2778	649	713	22		

- Molecule 2 is a protein called Cytochrome c oxidase subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	b	236	Total	C	N	O	S	0	0
			1889	1242	286	351	10		

- Molecule 3 is a protein called CYTOCHROME C OXIDASE SUBUNIT 3; SYNONYM: CYTOCHROME C OXIDASE POLYPEPTIDE III, COX3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	c	269	Total	C	N	O	S	0	0
			2146	1430	344	357	15		

- Molecule 4 is a protein called Cytochrome c oxidase subunit 4, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	d	121	Total	C	N	O	S	0	0
			913	576	151	181	5		

- Molecule 5 is a protein called Cytochrome c oxidase polypeptide 5A, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	e	133	Total	C	N	O	S	0	0
			1049	663	184	198	4		

- Molecule 6 is a protein called Cytochrome c oxidase subunit 6, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	f	102	Total	C	N	O	S	0	0
			851	545	137	168	1		

- Molecule 7 is a protein called CYTOCHROME C OXIDASE SUBUNIT 7; SYNONYM: CYTOCHROME C OXIDASE POLYPEPTIDE VII, COX7.

Mol	Chain	Residues	Atoms				AltConf	Trace
7	g	59	Total	C	N	O	0	0
			484	328	83	73		

- Molecule 8 is a protein called Cytochrome c oxidase polypeptide VIII, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	h	47	Total	C	N	O	S	0	0
			383	261	62	59	1		


- Molecule 9 is a protein called CYTOCHROME C OXIDASE SUBUNIT 7A; SYNONYM: CYTOCHROME C OXIDASE POLYPEPTIDE VIIA, COX9.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	i	55	Total	C	N	O	S	0	0
			456	300	79	74	3		

- Molecule 10 is COPPER (II) ION (CCD ID: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
10	a	1	Total	Cu	0
			1	1	

- Molecule 11 is HEME-A (CCD ID: HEA) (formula: C₄₉H₅₆FeN₄O₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
11	a	1	Total 60	C 49	Fe 1	N 4	O 6	0
11	a	1	Total 60	C 49	Fe 1	N 4	O 6	0

- Molecule 12 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	AltConf
12	a	1	Total Mg 1 1	0

- Molecule 13 is DI-PALMITOYL-3-SN-PHOSPHATIDYLETHANOLAMINE (CCD ID: PEF) (formula: $C_{37}H_{74}NO_8P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
13	a	1	Total	C	N	O	P	0
			33	23	1	8	1	
13	a	1	Total	C	N	O	P	0
			47	37	1	8	1	
13	c	1	Total	C	N	O	P	0
			36	26	1	8	1	

- Molecule 14 is DINUCLEAR COPPER ION (CCD ID: CUA) (formula: Cu₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		AltConf
14	b	1	Total 2	Cu 2	0


- Molecule 15 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
15	d	1	Total 1	Zn 1	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cytochrome c oxidase subunit 1




• Molecule 2: Cytochrome c oxidase subunit 2




- Molecule 3: CYTOCHROME C OXIDASE SUBUNIT 3; SYNONYM: CYTOCHROME C OXIDASE POLYPEPTIDE III, COX3




- Molecule 4: Cytochrome c oxidase subunit 4, mitochondrial




- Molecule 5: Cytochrome c oxidase polypeptide 5A, mitochondrial



- Molecule 6: Cytochrome c oxidase subunit 6, mitochondrial




- Molecule 7: CYTOCHROME C OXIDASE SUBUNIT 7; SYNONYM: CYTOCHROME C OXIDASE POLYPEPTIDE VII, COX7



- Molecule 8: Cytochrome c oxidase polypeptide VIII, mitochondrial





- Molecule 9: CYTOCHROME C OXIDASE SUBUNIT 7A; SYNONYM: CYTOCHROME C OXIDASE POLYPEPTIDE VIIA, COX9

Chain i: 11% 78% 18% .



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	72409	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	48	Depositor
Minimum defocus (nm)	1400	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	23.830	Depositor
Minimum map value	-14.394	Depositor
Average map value	0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	3.6	Depositor
Map size (Å)	208.64, 208.64, 208.64	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.815, 0.815, 0.815	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU, MG, CUA, PEF, HEA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	a	0.36	0/4290	0.58	2/5857 (0.0%)
2	b	0.25	0/1941	0.48	0/2653
3	c	0.26	0/2218	0.55	2/3036 (0.1%)
4	d	0.29	0/932	0.57	0/1269
5	e	0.25	0/1074	0.50	0/1451
6	f	0.27	0/868	0.45	0/1174
7	g	0.29	0/500	0.64	0/681
8	h	0.27	0/397	0.42	0/533
9	i	0.21	0/468	0.45	0/626
All	All	0.30	0/12688	0.54	4/17280 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	a	0	1
3	c	0	2
All	All	0	3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	c	135	THR	CA-C-N	5.63	132.30	121.54
3	c	135	THR	C-N-CA	5.63	132.30	121.54
1	a	119	VAL	CA-C-N	5.59	129.21	120.82
1	a	119	VAL	C-N-CA	5.59	129.21	120.82

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	a	520	SER	Peptide
3	c	124	TRP	Peptide
3	c	125	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	a	4162	0	4192	73	0
2	b	1889	0	1866	42	0
3	c	2146	0	2137	22	0
4	d	913	0	910	17	0
5	e	1049	0	1030	13	0
6	f	851	0	822	13	0
7	g	484	0	517	6	0
8	h	383	0	386	0	0
9	i	456	0	469	7	0
10	a	1	0	0	0	0
11	a	120	0	108	7	0
12	a	1	0	0	0	0
13	a	80	0	112	2	0
13	c	36	0	45	2	0
14	b	2	0	0	0	0
15	d	1	0	0	0	0
All	All	12574	0	12594	172	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 172 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:a:68:PHE:HB3	1:a:154:THR:HG21	1.73	0.70
4:d:115:PRO:HD2	4:d:118:SER:HB2	1.77	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:a:378:HIS:O	1:a:382:SER:HB3	1.95	0.66
2:b:223:GLU:O	2:b:229:HIS:ND1	2.29	0.66
1:a:532:VAL:HB	4:d:107:ARG:HH21	1.61	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	a	532/534 (100%)	499 (94%)	33 (6%)	0	100	100
2	b	234/236 (99%)	217 (93%)	17 (7%)	0	100	100
3	c	267/269 (99%)	251 (94%)	16 (6%)	0	100	100
4	d	119/121 (98%)	95 (80%)	24 (20%)	0	100	100
5	e	131/133 (98%)	120 (92%)	11 (8%)	0	100	100
6	f	100/104 (96%)	98 (98%)	2 (2%)	0	100	100
7	g	57/59 (97%)	50 (88%)	7 (12%)	0	100	100
8	h	45/47 (96%)	42 (93%)	3 (7%)	0	100	100
9	i	53/55 (96%)	51 (96%)	2 (4%)	0	100	100
All	All	1538/1558 (99%)	1423 (92%)	115 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	a	447/447 (100%)	399 (89%)	48 (11%)	6	24
2	b	209/209 (100%)	196 (94%)	13 (6%)	16	43
3	c	228/228 (100%)	206 (90%)	22 (10%)	8	29
4	d	102/102 (100%)	91 (89%)	11 (11%)	6	24
5	e	110/110 (100%)	105 (96%)	5 (4%)	24	49
6	f	91/93 (98%)	84 (92%)	7 (8%)	12	37
7	g	50/50 (100%)	43 (86%)	7 (14%)	3	17
8	h	39/39 (100%)	36 (92%)	3 (8%)	12	37
9	i	46/46 (100%)	42 (91%)	4 (9%)	9	33
All	All	1322/1324 (100%)	1202 (91%)	120 (9%)	11	31

5 of 120 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	b	205	ASN
7	g	42	VAL
3	c	145	LEU
7	g	41	VAL
9	i	20	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
3	c	251	HIS
6	f	70	GLN
4	d	144	ASN
6	f	120	GLN
1	a	475	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
13	PEF	c	301	-	35,35,46	1.07	2 (5%)	38,40,51	1.15	3 (7%)
13	PEF	a	605	-	32,32,46	1.13	2 (6%)	35,37,51	1.28	5 (14%)
13	PEF	a	606	-	46,46,46	0.97	2 (4%)	49,51,51	1.10	3 (6%)
14	CUA	b	301	2	0,1,1	-	-	-	-	-
11	HEA	a	603	1	67,67,67	2.29	25 (37%)	81,103,103	2.60	35 (43%)
11	HEA	a	602	1	67,67,67	2.35	27 (40%)	81,103,103	2.65	35 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
13	PEF	c	301	-	-	9/39/39/50	-
13	PEF	a	605	-	-	11/36/36/50	-
13	PEF	a	606	-	-	19/50/50/50	-
11	HEA	a	603	1	-	11/36/76/76	-
11	HEA	a	602	1	-	15/36/76/76	-

The worst 5 of 58 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	a	603	HEA	FE-NB	5.15	2.10	1.94
11	a	602	HEA	C3B-C2B	5.09	1.46	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	a	603	HEA	FE-ND	5.09	2.10	1.94
11	a	602	HEA	FE-NB	5.06	2.10	1.94
11	a	603	HEA	C3B-C2B	4.96	1.46	1.34

The worst 5 of 81 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	a	603	HEA	C3D-C4D-ND	7.23	117.34	110.35
11	a	602	HEA	C2B-C1B-NB	6.18	117.04	109.90
11	a	603	HEA	C3C-C2C-C1C	-6.09	99.99	107.17
11	a	603	HEA	C2A-C1A-NA	6.09	116.20	110.32
11	a	602	HEA	C3C-C4C-NC	6.00	114.85	109.80

There are no chirality outliers.

5 of 65 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	a	602	HEA	C2A-C3A-CMA-OMA
11	a	602	HEA	C4A-C3A-CMA-OMA
11	a	602	HEA	C11-C12-C13-C14
11	a	603	HEA	C2A-C3A-CMA-OMA
11	a	603	HEA	C4A-C3A-CMA-OMA


There are no ring outliers.


4 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
13	c	301	PEF	2	0
13	a	605	PEF	2	0
11	a	603	HEA	3	0
11	a	602	HEA	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-14436. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 117



Y Index: 113




Z Index: 133


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 3.6. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 86 nm^3 ; this corresponds to an approximate mass of 78 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.258 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.258 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.87	-	-
Author-provided FSC curve	3.65	4.26	3.77
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-14436 and PDB model 7Z10. Per-residue inclusion information can be found in section [3](#) on page [9](#).

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 3.6 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 85% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (3.6) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7350	0.4330
a	0.7700	0.4500
b	0.6560	0.4180
c	0.6600	0.4080
d	0.8020	0.4520
e	0.7590	0.4320
f	0.8200	0.4270
g	0.7580	0.4460
h	0.7880	0.4540
i	0.6750	0.3980

1.0

0.0

<0.0