



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:44 PM UTC

PDB ID : 2Z2T / pdb_00002z2t
Title : Crystal structure of the complex between gp41 fragment N36 and fusion inhibitor SC34EK
Authors : Nakamura, S.; Ohkubo, T.; Kobayashi, Y.
Deposited on : 2007-05-28
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	38	11% 87%13%
1	B	38	11% 68%24%8%
1	C	38	3% 87%11%
2	D	36	11% 81%19%
2	E	36	22% 78%19%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	36	 A horizontal bar chart showing the quality of chain F. The bar is divided into three segments: a red segment on the left labeled '6%', a green segment in the middle labeled '86%', and a yellow segment on the right labeled '14%'.

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2010 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called gp41 fragment N36.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	38	Total	C	N	O	0	0	1
			298	188	57	53			
1	B	38	Total	C	N	O	0	0	1
			298	188	57	53			
1	C	38	Total	C	N	O	0	0	1
			298	188	57	53			

- Molecule 2 is a protein called Fusion inhibitor peptide SC34EK.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	36	Total	C	N	O	0	0	1
			312	198	52	62			
2	E	36	Total	C	N	O	0	0	1
			312	198	52	62			
2	F	36	Total	C	N	O	0	0	1
			312	198	52	62			

- Molecule 3 is ACETIC ACID (CCD ID: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	37	Total 37	O 37	0	0
5	B	17	Total 17	O 17	0	0
5	C	29	Total 29	O 29	0	0
5	D	30	Total 30	O 30	0	0
5	E	27	Total 27	O 27	0	0
5	F	31	Total 31	O 31	0	0


- Molecule 1: gp41 fragment N36




- Molecule 1: gp41 fragment N36



- Molecule 1: gp41 fragment N36




- Molecule 2: Fusion inhibitor peptide SC34EK





- Molecule 2: Fusion inhibitor peptide SC34EK





- Molecule 2: Fusion inhibitor peptide SC34EK



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	105.01Å 105.01Å 78.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.85 – 2.10 19.85 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.85-2.10) 99.6 (19.85-2.10)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.54 (at 2.09Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.213 , 0.238 0.211 , 0.238	Depositor DCC
R_{free} test set	1494 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 43.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2010	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACY, NLE, ACE, NH2, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/297	0.92	0/402
1	B	0.59	0/297	1.06	1/402 (0.2%)
1	C	0.62	0/297	0.95	1/402 (0.2%)
2	D	0.56	0/304	1.10	2/401 (0.5%)
2	E	0.52	0/304	0.96	0/401
2	F	0.53	0/304	1.10	2/401 (0.5%)
All	All	0.57	0/1803	1.02	6/2409 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
2	E	0	1
All	All	0	2

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2580	ILE	O-C-N	-11.39	108.91	122.12
2	D	1660	LEU	CA-C-N	9.01	137.92	121.70
2	D	1660	LEU	C-N-CA	9.01	137.92	121.70
2	F	3660	LEU	CA-C-N	6.49	133.38	121.70
2	F	3660	LEU	C-N-CA	6.49	133.38	121.70
1	C	3580	ILE	O-C-N	-5.71	114.24	122.12

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	2580	ILE	Mainchain
2	E	2660	LEU	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	298	0	314	5	0
1	B	298	0	314	10	0
1	C	298	0	314	1	0
2	D	312	0	314	5	0
2	E	312	0	314	6	0
2	F	312	0	314	2	0
3	C	4	0	3	0	0
4	E	5	0	0	0	0
5	A	37	0	0	0	0
5	B	17	0	0	1	0
5	C	29	0	0	0	0
5	D	30	0	0	1	0
5	E	27	0	0	0	0
5	F	31	0	0	1	0
All	All	2010	0	1887	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2562:GLN:HG2	2:E:2642:ILE:CD1	1.98	0.94
1:B:2562:GLN:HG2	2:E:2642:ILE:HD12	1.54	0.87
1:A:1552:GLN:HE21	1:B:2552:GLN:HE21	1.22	0.83
2:F:3643:GLU:HG2	5:F:146:HOH:O	1.90	0.70
1:A:1551:GLN:HE21	2:D:1653:GLN:HE21	1.40	0.68
1:B:2554:ASN:ND2	5:B:162:HOH:O	2.25	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2551:GLN:OE1	1:B:2551:GLN:HA	2.02	0.60
1:B:2562:GLN:HG2	2:E:2642:ILE:HD11	1.84	0.58
2:F:3633:ARG:NH2	2:F:3637:GLU:OE1	2.37	0.57
1:B:2565:LEU:HD13	2:E:2638:TYR:HB2	1.88	0.56
2:E:2629:NLE:HE3	2:E:2633:ARG:HH21	1.77	0.50
2:D:1633:ARG:NH2	5:D:79:HOH:O	2.39	0.49
1:A:1566:LEU:HG	1:B:2566:LEU:HD21	1.94	0.49
1:A:1571:TRP:CH2	1:A:1575:GLN:HG3	2.52	0.45
1:A:1551:GLN:NE2	2:D:1653:GLN:HE21	2.12	0.43
1:C:3577:GLN:HG3	2:E:2628:TRP:CE3	2.52	0.43
2:D:1658:LYS:HA	2:D:1661:LYS:HE2	2.01	0.43
1:B:2551:GLN:O	1:B:2555:LEU:HG	2.19	0.41
1:B:2577:GLN:HG3	2:D:1628:TRP:CE3	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	36/38 (95%)	36 (100%)	0	0	100	100
1	B	36/38 (95%)	36 (100%)	0	0	100	100
1	C	36/38 (95%)	35 (97%)	1 (3%)	0	100	100
2	D	33/36 (92%)	33 (100%)	0	0	100	100
2	E	33/36 (92%)	33 (100%)	0	0	100	100
2	F	33/36 (92%)	33 (100%)	0	0	100	100
All	All	207/222 (93%)	206 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	32/32 (100%)	32 (100%)	0	100	100
1	B	32/32 (100%)	27 (84%)	5 (16%)	2	1
1	C	32/32 (100%)	28 (88%)	4 (12%)	4	2
2	D	33/33 (100%)	32 (97%)	1 (3%)	36	41
2	E	33/33 (100%)	30 (91%)	3 (9%)	9	6
2	F	33/33 (100%)	32 (97%)	1 (3%)	36	41
All	All	195/195 (100%)	181 (93%)	14 (7%)	13	11

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2546	SER
1	B	2550	GLN
1	B	2551	GLN
1	B	2565	LEU
1	B	2581	LEU
1	C	3565	LEU
1	C	3566	LEU
1	C	3577	GLN
1	C	3581	LEU
2	D	1632	ASP
2	E	2636	GLU
2	E	2652	GLN
2	E	2660	LEU
2	F	3661	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1552	GLN
1	A	1564	HIS
1	A	1577	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	2563	GLN
1	B	2564	HIS
1	B	2577	GLN
1	C	3551	GLN
1	C	3563	GLN
1	C	3564	HIS
1	C	3567	GLN
1	C	3577	GLN
2	D	1653	GLN
2	E	2652	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NLE	F	3629	2	6,7,8	0.55	0	2,7,9	0.45	0
2	NLE	D	1629	2	6,7,8	0.54	0	2,7,9	0.39	0
2	NLE	E	2629	2	6,7,8	0.43	0	2,7,9	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NLE	F	3629	2	-	0/5/6/8	-
2	NLE	D	1629	2	-	0/5/6/8	-
2	NLE	E	2629	2	-	0/5/6/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	2629	NLE	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	E	4001	-	4,4,4	0.27	0	6,6,6	0.25	0
3	ACY	C	5001	-	3,3,3	0.81	0	3,3,3	0.86	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	36/38 (94%)	-0.28	0 100 100	16, 21, 32, 38	0
1	B	36/38 (94%)	0.37	4 (11%) 10 10	19, 27, 45, 46	0
1	C	36/38 (94%)	-0.08	1 (2%) 55 57	16, 22, 29, 33	0
2	D	33/36 (91%)	0.91	4 (12%) 8 9	20, 32, 62, 68	0
2	E	33/36 (91%)	1.37	8 (24%) 2 2	21, 34, 54, 58	0
2	F	33/36 (91%)	0.49	2 (6%) 27 29	22, 26, 33, 37	0
All	All	207/222 (93%)	0.44	19 (9%) 14 15	16, 27, 47, 68	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	3644	GLU	3.9
2	D	1661	LYS	3.7
2	D	1658	LYS	3.5
2	D	1660	LEU	3.4
1	B	2546	SER	3.3
2	E	2633	ARG	3.3
2	E	2660	LEU	3.2
2	E	2659	GLU	3.1
2	E	2654	GLU	3.0
2	D	1654	GLU	3.0
1	B	2547	ASP	2.7
2	E	2637	GLU	2.7
1	B	2581	LEU	2.6
2	E	2643	GLU	2.5
1	B	2567	GLN	2.5
1	C	3550	GLN	2.4
2	E	2652	GLN	2.4
2	E	2661	LYS	2.3
2	F	3636	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NLE	F	3629	8/9	0.92	0.12	26,28,30,34	0
2	NLE	E	2629	8/9	0.95	0.08	21,23,27,29	0
2	NLE	D	1629	8/9	0.96	0.07	19,21,21,21	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ACY	C	5001	4/4	0.79	0.27	48,48,48,48	0
4	SO4	E	4001	5/5	0.99	0.05	26,27,28,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.