

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BTB	B	202	-	-	X	-

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:ASN:OD1	1:B:53:GLN:NE2[4_555]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	125/134 (93%)	125 (100%)	0	0	100	100
1	B	125/134 (93%)	124 (99%)	1 (1%)	0	100	100
All	All	250/268 (93%)	249 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	103/108 (95%)	102 (99%)	1 (1%)	68	52
1	B	103/108 (95%)	98 (95%)	5 (5%)	22	5
All	All	206/216 (95%)	200 (97%)	6 (3%)	43	14

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	104	GLN
1	B	10	ASP
1	B	22	ASN
1	B	23[A]	GLN
1	B	23[B]	GLN
1	B	108	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	ASN
1	A	51	ASN
1	B	19	ASN
1	B	22	ASN
1	B	40	GLN
1	B	104	GLN
1	B	127	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	202	BTB	C4-C2-C3-O3
3	A	202	BTB	N-C2-C3-O3
3	B	202	BTB	C1-C2-C4-O4
3	B	202	BTB	C3-C2-C4-O4
3	B	202	BTB	N-C2-C4-O4
3	B	202	BTB	C1-C2-N-C5
3	B	202	BTB	C1-C2-N-C7
3	B	202	BTB	C3-C2-N-C5
3	B	202	BTB	C3-C2-N-C7
3	B	202	BTB	C4-C2-N-C5
3	B	202	BTB	C4-C2-N-C7
3	B	202	BTB	C6-C5-N-C2
3	B	202	BTB	N-C7-C8-O8
3	B	202	BTB	N-C5-C6-O6
2	B	201	BTN	S1-C2-C7-C8
3	B	202	BTB	C8-C7-N-C2
3	A	202	BTB	N-C7-C8-O8
2	A	201	BTN	C9-C10-C11-O11
3	B	202	BTB	O1-C1-C2-C3
2	A	201	BTN	C9-C10-C11-O12

There are no ring outliers.

1 monomer is involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	202	BTB	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	127/134 (94%)	0.12	10 (7%)	18 21	23, 31, 60, 78	0
1	B	126/134 (94%)	0.02	4 (3%)	50 54	23, 30, 48, 78	1 (0%)
All	All	253/268 (94%)	0.07	14 (5%)	30 34	23, 31, 55, 78	1 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	9	PRO	4.0
1	A	15	ALA	3.5
1	A	133	SER	2.6
1	A	17	ALA	2.6
1	B	10	ASP	2.5
1	A	9	PRO	2.5
1	A	7	MET	2.4
1	B	11	MET	2.4
1	A	104	GLN	2.4
1	A	12	LYS	2.4
1	A	103	SER	2.3
1	B	134	VAL	2.3
1	A	16	GLY	2.0
1	A	53	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BTB	B	202	14/14	0.88	0.16	46,65,90,93	0
3	BTB	A	202	14/14	0.92	0.12	36,45,58,68	0
2	BTN	A	201	16/16	0.98	0.04	23,26,30,32	0
2	BTN	B	201	16/16	0.99	0.04	24,27,31,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.