



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 20, 2026 – 09:24 am BST

PDB ID : 7Z26 / pdb_00007z26
Title : Crystal structure of YTHDF2 YTH domain in complex with m6A RNA
Authors : Nai, F.; Nachawati, R.; Li, Y.; Caffisch, A.
Deposited on : 2022-02-25
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)
RNA backbone	3983	1098 (2.30-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	5% 83% 7% 10%
1	B	167	2% 84% 5% 11%
2	D	5	20% 40% 40%
2	E	5	40% 40% 20%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2707 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YTH domain-containing family protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	150	Total	C	N	O	S	0	2	0
			1188	753	211	219	5			
1	B	149	Total	C	N	O	S	0	2	0
			1201	766	209	220	6			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	386	MET	-	initiating methionine	UNP Q9Y5A9
A	387	GLY	-	expression tag	UNP Q9Y5A9
A	388	SER	-	expression tag	UNP Q9Y5A9
A	389	SER	-	expression tag	UNP Q9Y5A9
A	390	TYR	-	expression tag	UNP Q9Y5A9
A	391	HIS	-	expression tag	UNP Q9Y5A9
A	392	HIS	-	expression tag	UNP Q9Y5A9
A	393	HIS	-	expression tag	UNP Q9Y5A9
A	394	HIS	-	expression tag	UNP Q9Y5A9
A	395	HIS	-	expression tag	UNP Q9Y5A9
A	396	HIS	-	expression tag	UNP Q9Y5A9
A	397	SER	-	expression tag	UNP Q9Y5A9
A	398	SER	-	expression tag	UNP Q9Y5A9
A	399	GLY	-	expression tag	UNP Q9Y5A9
A	400	GLU	-	expression tag	UNP Q9Y5A9
A	401	ASN	-	expression tag	UNP Q9Y5A9
A	402	LEU	-	expression tag	UNP Q9Y5A9
A	403	TYR	-	expression tag	UNP Q9Y5A9
A	404	PHE	-	expression tag	UNP Q9Y5A9
A	405	GLN	-	expression tag	UNP Q9Y5A9
A	406	HIS	-	expression tag	UNP Q9Y5A9
A	407	MET	-	expression tag	UNP Q9Y5A9
B	386	MET	-	initiating methionine	UNP Q9Y5A9
B	387	GLY	-	expression tag	UNP Q9Y5A9
B	388	SER	-	expression tag	UNP Q9Y5A9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	389	SER	-	expression tag	UNP Q9Y5A9
B	390	TYR	-	expression tag	UNP Q9Y5A9
B	391	HIS	-	expression tag	UNP Q9Y5A9
B	392	HIS	-	expression tag	UNP Q9Y5A9
B	393	HIS	-	expression tag	UNP Q9Y5A9
B	394	HIS	-	expression tag	UNP Q9Y5A9
B	395	HIS	-	expression tag	UNP Q9Y5A9
B	396	HIS	-	expression tag	UNP Q9Y5A9
B	397	SER	-	expression tag	UNP Q9Y5A9
B	398	SER	-	expression tag	UNP Q9Y5A9
B	399	GLY	-	expression tag	UNP Q9Y5A9
B	400	GLU	-	expression tag	UNP Q9Y5A9
B	401	ASN	-	expression tag	UNP Q9Y5A9
B	402	LEU	-	expression tag	UNP Q9Y5A9
B	403	TYR	-	expression tag	UNP Q9Y5A9
B	404	PHE	-	expression tag	UNP Q9Y5A9
B	405	GLN	-	expression tag	UNP Q9Y5A9
B	406	HIS	-	expression tag	UNP Q9Y5A9
B	407	MET	-	expression tag	UNP Q9Y5A9

- Molecule 2 is a RNA chain called RNA (5'-R(P*(6MZ)P*CP*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	3	Total 63	C 29	N 10	O 21	P 3	0	0	0
2	E	4	Total 64	C 29	N 10	O 22	P 3	0	0	1

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	90	Total 90	O 90	0	0
5	B	75	Total 75	O 75	0	0
5	D	3	Total 3	O 3	0	0
5	E	7	Total 7	O 7	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: YTH domain-containing family protein 2



- Molecule 1: YTH domain-containing family protein 2



- Molecule 2: RNA (5'-R(P*(6MZ)P*CP*U)-3')



- Molecule 2: RNA (5'-R(P*(6MZ)P*CP*U)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	80.69Å 80.69Å 114.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.34 – 1.90 40.34 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.34-1.90) 99.9 (40.34-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 1.89Å)	Xtriage
Refinement program	PHENIX 1.20_4459	Depositor
R, R_{free}	0.197 , 0.230 0.197 , 0.230	Depositor DCC
R_{free} test set	1667 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	28.6	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.088 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2707	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6MZ, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/1217	0.55	0/1647
1	B	0.35	0/1232	0.55	0/1666
2	D	0.35	0/43	0.70	0/64
2	E	0.30	0/43	0.41	0/64
All	All	0.36	0/2535	0.55	0/3441

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1188	0	1092	9	0
1	B	1201	0	1121	11	0
2	D	63	0	35	0	0
2	E	64	0	35	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	B	6	0	8	2	0
5	A	90	0	0	3	1
5	B	75	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	3	0	0	0	0
5	E	7	0	0	0	0
All	All	2707	0	2291	20	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:489:ASP:H	4:B:602:GOL:H31	1.50	0.77
1:B:405:GLN:HE22	1:B:542:LYS:HE3	1.61	0.66
1:B:538:LYS:HG2	1:B:542:LYS:HZ3	1.59	0.65
1:A:399:GLY:N	5:A:702:HOH:O	2.30	0.64
1:B:538:LYS:O	1:B:542:LYS:HD3	2.02	0.60
1:B:538:LYS:CG	1:B:542:LYS:HZ3	2.15	0.59
1:B:536:LYS:HE2	5:B:748:HOH:O	2.04	0.57
1:B:488:GLN:HG2	4:B:602:GOL:H2	1.87	0.55
1:A:536:LYS:HE2	5:A:759:HOH:O	2.06	0.55
1:B:538:LYS:HE2	1:B:542:LYS:HZ3	1.75	0.52
1:A:516:GLU:OE1	1:A:536:LYS:NZ	2.44	0.51
1:B:458:LEU:HD13	1:B:540:VAL:HG11	1.92	0.50
1:B:538:LYS:HE2	1:B:542:LYS:NZ	2.27	0.49
1:B:401:ASN:O	1:B:405:GLN:HG2	2.14	0.47
1:A:474:LYS:HA	1:A:474:LYS:HD2	1.67	0.45
1:A:412:VAL:HG22	1:A:456:TYR:HB2	2.00	0.44
1:A:510:LEU:HD22	1:A:540:VAL:HG13	2.00	0.43
1:A:449:MET:HE2	1:A:449:MET:HA	2.02	0.42
1:A:520:ASN:ND2	5:A:701:HOH:O	2.23	0.41
1:A:412:VAL:HG21	1:A:537:ALA:HB1	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:780:HOH:O	5:A:782:HOH:O[5_555]	2.05	0.15

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/167 (90%)	146 (97%)	4 (3%)	0	100	100
1	B	149/167 (89%)	147 (99%)	2 (1%)	0	100	100
All	All	299/334 (90%)	293 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	119/149 (80%)	118 (99%)	1 (1%)	73	75
1	B	122/149 (82%)	122 (100%)	0	100	100
All	All	241/298 (81%)	240 (100%)	1 (0%)	84	87

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	400	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	488	GLN
1	A	525	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	405	GLN
1	B	488	GLN
1	B	539	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	D	2/5 (40%)	1 (50%)	0
2	E	2/5 (40%)	1 (50%)	0
All	All	4/10 (40%)	2 (50%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	D	4	C
2	E	4	C

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	6MZ	D	3	2	22,25,26	2.47	4 (18%)	30,36,39	3.22	13 (43%)
2	6MZ	E	3	2	22,25,26	2.53	4 (18%)	30,36,39	3.20	12 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6MZ	D	3	2	-	0/9/27/28	0/3/3/3
2	6MZ	E	3	2	-	0/9/27/28	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	3	6MZ	C6-N6	10.46	1.45	1.34
2	D	3	6MZ	C6-N6	10.19	1.45	1.34
2	E	3	6MZ	C5-N7	-2.61	1.34	1.39
2	D	3	6MZ	C5-N7	-2.33	1.34	1.39
2	D	3	6MZ	C2-N1	2.24	1.38	1.33
2	E	3	6MZ	C5-C4	-2.24	1.34	1.39
2	D	3	6MZ	C5-C4	-2.03	1.35	1.39
2	E	3	6MZ	C2-N1	2.00	1.37	1.33

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	3	6MZ	C4-C5-C6	7.25	122.43	116.81
2	E	3	6MZ	C5-C4-N3	-7.15	117.42	126.75
2	D	3	6MZ	C4-N9-C1'	-6.67	110.71	126.59
2	E	3	6MZ	C4-N9-C1'	-6.64	110.78	126.59
2	D	3	6MZ	C4-C5-C6	6.56	121.89	116.81
2	D	3	6MZ	C5-C4-N3	-6.47	118.31	126.75
2	E	3	6MZ	N3-C4-N9	6.18	137.27	127.08
2	E	3	6MZ	C1'-N9-C8	6.07	140.84	127.14
2	D	3	6MZ	C1'-N9-C8	5.90	140.46	127.14
2	D	3	6MZ	N3-C4-N9	5.81	136.66	127.08
2	D	3	6MZ	N1-C2-N3	-4.87	120.99	128.60
2	E	3	6MZ	C6-C5-N7	-4.54	127.47	132.39
2	D	3	6MZ	C6-C5-N7	-4.24	127.80	132.39
2	E	3	6MZ	N1-C2-N3	-4.18	122.07	128.60
2	D	3	6MZ	N9-C8-N7	-3.96	108.49	113.91
2	D	3	6MZ	C9-N6-C6	3.96	126.28	122.87
2	E	3	6MZ	N9-C8-N7	-3.38	109.29	113.91
2	E	3	6MZ	C2-N3-C4	3.35	119.66	111.75
2	D	3	6MZ	C2-N3-C4	3.27	119.48	111.75
2	D	3	6MZ	C5-N7-C8	2.79	107.48	103.51
2	D	3	6MZ	C4-N9-C8	2.70	108.65	105.73
2	E	3	6MZ	C5-N7-C8	2.56	107.14	103.51
2	E	3	6MZ	C5-C6-N1	2.51	120.87	118.20
2	E	3	6MZ	C4-N9-C8	2.19	108.10	105.73
2	D	3	6MZ	C5-C6-N1	2.15	120.49	118.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	B	602	-	5,5,5	0.83	0	5,5,5	1.16	0
3	SO4	B	601	-	4,4,4	0.15	0	6,6,6	0.17	0
3	SO4	A	601	-	4,4,4	0.15	0	6,6,6	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	602	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	602	GOL	O1-C1-C2-C3
4	B	602	GOL	O1-C1-C2-O2
4	B	602	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	602	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	150/167 (89%)	0.10	8 (5%) 32 34	12, 28, 47, 59	2 (1%)
1	B	149/167 (89%)	0.32	3 (2%) 65 69	11, 31, 49, 54	2 (1%)
2	D	2/5 (40%)	0.21	0 100 100	40, 40, 40, 47	0
2	E	3/5 (60%)	0.61	0 100 100	36, 36, 46, 51	0
All	All	304/344 (88%)	0.21	11 (3%) 46 49	11, 30, 49, 59	4 (1%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	406	HIS	3.8
1	A	409	HIS	3.0
1	A	407	MET	2.9
1	A	404	PHE	2.4
1	B	406	HIS	2.3
1	B	410	GLY	2.2
1	A	500	ILE	2.2
1	A	548	LYS	2.1
1	A	437	HIS	2.1
1	B	418[A]	TYR	2.1
1	A	489	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	6MZ	D	3	23/24	0.94	0.08	20,24,35,44	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	6MZ	E	3	23/24	0.96	0.06	20,23,45,54	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	B	602	6/6	0.87	0.12	38,43,51,55	0
3	SO4	B	601	5/5	0.92	0.08	48,50,61,61	0
3	SO4	A	601	5/5	0.93	0.09	35,46,52,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.