



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:31 PM UTC

PDB ID : 8Z2V / pdb_00008z2v
Title : Crystal structure of Sonic hedgehog in complex with antibody 5E1 mutant H-R102A with metals
Authors : Caaveiro, J.M.M.; Kaneda, I.; Tsumoto, K.
Deposited on : 2024-04-13
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	169	8% 72% 15% • 11%
2	H	241	5% 67% 20% 12%
3	L	235	% 74% 14% • 10%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4913 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sonic hedgehog protein N-product.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	150	Total	C	N	O	S	0	0	0
			1205	753	214	233	5			

- Molecule 2 is a protein called Heavy chain of antibody 5E1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	213	Total	C	N	O	S	0	3	0
			1636	1041	266	324	5			

- Molecule 3 is a protein called Light chain of antibody 5E1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	211	Total	C	N	O	S	0	3	0
			1648	1041	271	330	6			

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Ca	0	0
			2	2		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	83	Total	O	0	2
			85	85		
8	H	163	Total	O	0	2
			165	165		
8	L	163	Total	O	0	1
			164	164		

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	96.64Å 69.75Å 119.27Å 90.00° 108.73° 90.00°	Depositor
Resolution (Å)	56.48 – 1.89 56.48 – 1.89	Depositor EDS
% Data completeness (in resolution range)	97.0 (56.48-1.89) 97.0 (56.48-1.89)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.186 , 0.225 (Not available) , 0.227	Depositor DCC
R_{free} test set	2384 reflections (3.96%)	wwPDB-VP
Wilson B-factor (Å ²)	28.0	Xtriage
Anisotropy	0.579	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4913	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, CA, ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.34	5/1231 (0.4%)	1.57	14/1661 (0.8%)
2	H	1.47	14/1686 (0.8%)	1.59	15/2298 (0.7%)
3	L	1.30	7/1694 (0.4%)	1.47	10/2302 (0.4%)
All	All	1.37	26/4611 (0.6%)	1.54	39/6261 (0.6%)

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	44	SER	C-O	10.66	1.34	1.24
1	A	140	HIS	CE1-NE2	9.21	1.41	1.32
3	L	135	LEU	C-O	8.21	1.33	1.24
2	H	140	GLY	N-CA	7.08	1.56	1.45
2	H	154	GLU	CD-OE1	6.83	1.38	1.25
2	H	73	ASP	C-O	6.44	1.31	1.24
2	H	33	ALA	C-O	6.08	1.31	1.23
1	A	84	ILE	C-O	6.05	1.30	1.24
2	H	125	PRO	C-O	-6.05	1.16	1.23
3	L	45	LYS	C-O	5.94	1.31	1.24
2	H	51	ILE	C-O	5.74	1.29	1.24
2	H	19	LYS	C-O	-5.73	1.17	1.24
3	L	198	HIS	CG-ND1	5.66	1.44	1.38
2	H	193	SER	CB-OG	5.65	1.53	1.42
2	H	181	LEU	C-O	5.64	1.30	1.23
2	H	191	PRO	CA-C	5.64	1.59	1.52
3	L	177	SER	C-O	5.56	1.30	1.24
2	H	15	GLY	C-O	-5.55	1.15	1.24
1	A	48	ILE	N-CA	5.52	1.50	1.45
2	H	168	GLY	N-CA	5.40	1.53	1.45
3	L	164	THR	C-O	5.33	1.31	1.23
1	A	147	ASP	CG-OD1	5.21	1.35	1.25
2	H	92	ALA	C-O	5.17	1.29	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	L	111	ALA	C-O	5.12	1.29	1.23
3	L	198	HIS	CE1-NE2	5.05	1.37	1.32
1	A	102	CYS	C-O	5.04	1.29	1.24

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	14	PRO	CA-C-O	-7.57	112.81	121.36
3	L	17	ASP	CA-CB-CG	7.38	119.98	112.60
1	A	40	THR	CA-CB-OG1	-6.83	99.36	109.60
1	A	71	GLU	CB-CA-C	6.55	121.34	110.92
2	H	89	GLU	N-CA-CB	6.53	121.23	110.39
2	H	32	GLU	CB-CG-CD	6.31	123.32	112.60
1	A	89	GLU	CB-CG-CD	6.11	122.98	112.60
1	A	92	THR	CA-CB-OG1	-6.07	100.50	109.60
1	A	50	ASN	CB-CA-C	-6.01	98.45	110.42
1	A	163	ARG	N-CA-C	-5.90	104.75	111.07
2	H	11	LEU	CA-C-O	-5.89	113.96	120.38
1	A	41	PRO	CA-C-O	-5.78	114.75	121.34
1	A	166	VAL	CA-C-O	-5.73	115.16	121.29
2	H	87	THR	CA-CB-OG1	-5.71	101.04	109.60
3	L	77	THR	CA-CB-OG1	-5.65	101.12	109.60
2	H	52	ARG	CB-CA-C	5.60	115.48	110.44
1	A	104	ASP	CB-CA-C	5.58	119.63	110.88
2	H	163	GLY	N-CA-C	-5.58	107.67	114.92
3	L	151	ASP	CA-CB-CG	5.56	118.16	112.60
3	L	206	THR	CA-CB-OG1	-5.43	101.45	109.60
2	H	89	GLU	CB-CA-C	-5.41	99.01	110.31
1	A	175	TYR	CA-C-N	5.40	127.96	120.29
1	A	175	TYR	C-N-CA	5.40	127.96	120.29
2	H	171	THR	CA-C-O	-5.30	114.79	120.46
2	H	156	VAL	CA-C-O	-5.28	115.69	121.28
3	L	197	THR	CA-CB-OG1	-5.28	101.69	109.60
1	A	47	PHE	CA-C-N	5.27	127.43	122.85
1	A	47	PHE	C-N-CA	5.27	127.43	122.85
2	H	31	ASP	CA-C-O	-5.26	114.15	120.10
2	H	190	VAL	O-C-N	5.24	127.07	121.10
3	L	206	THR	CA-C-O	-5.23	115.17	120.71
2	H	14	PRO	O-C-N	5.22	129.52	122.89
2	H	197	THR	CB-CA-C	5.22	117.43	109.34
3	L	59	PRO	CA-C-N	5.15	127.70	120.28
3	L	59	PRO	C-N-CA	5.15	127.70	120.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	105	GLU	CB-CG-CD	5.10	121.27	112.60
3	L	201	LEU	CA-C-O	-5.09	114.83	120.38
1	A	170	PHE	CB-CA-C	5.05	117.78	109.90
2	H	119	SER	CB-CA-C	5.01	118.46	110.09

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1205	0	1164	8	0
2	H	1636	0	1600	16	0
3	L	1648	0	1612	14	0
4	A	1	0	0	1	0
5	A	1	0	0	0	0
6	A	2	0	0	0	0
7	H	6	0	8	0	0
8	A	85	0	0	3	0
8	H	165	0	0	3	0
8	L	164	0	0	2	0
All	All	4913	0	4384	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:52:ALA:HB3	1:A:55:THR:HG23	1.67	0.76
3:L:4:MET:HE3	3:L:23[A]:CYS:SG	2.28	0.73
4:A:201:CL:CL	8:A:348:HOH:O	2.47	0.69
3:L:198:HIS:CD2	3:L:200:GLY:H	2.11	0.68
2:H:98:ARG:NE	8:H:402:HOH:O	2.30	0.62
2:H:98:ARG:NH2	8:H:402:HOH:O	2.29	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:4:MET:CE	3:L:23[A]:CYS:SG	2.91	0.59
1:A:77:THR:HG21	1:A:103:LYS:HD2	1.85	0.57
1:A:78:PRO:HB2	1:A:80:TYR:CE2	2.42	0.54
2:H:66:ASP:O	2:H:85:ARG:NH2	2.41	0.54
3:L:149:LYS:HE2	3:L:154:LEU:HD11	1.91	0.53
3:L:161:GLU:OE2	8:L:301:HOH:O	2.19	0.53
3:L:37:GLN:HB2	3:L:47:LEU:HD11	1.91	0.52
2:H:52:ARG:HD3	2:H:54:TYR:OH	2.11	0.51
3:L:198:HIS:HD2	3:L:200:GLY:H	1.55	0.50
1:A:176:GLU:OE2	8:A:349[B]:HOH:O	2.18	0.49
2:H:98:ARG:CZ	8:H:402:HOH:O	2.59	0.49
3:L:141:PRO:O	3:L:198:HIS:HE1	1.95	0.49
3:L:203:SER:HB2	8:L:318:HOH:O	2.14	0.48
3:L:35:TRP:CE2	3:L:73:PHE:HB2	2.49	0.47
3:L:24:LYS:HA	3:L:69:THR:O	2.15	0.46
2:H:47:TRP:CE3	3:L:96:PRO:HD2	2.50	0.46
1:A:49:PRO:HD3	8:A:366:HOH:O	2.16	0.45
1:A:42:LEU:HD11	1:A:48:ILE:HD12	1.98	0.45
2:H:89:GLU:OE1	2:H:89:GLU:CA	2.64	0.45
2:H:207:LYS:N	2:H:208:PRO:CD	2.80	0.45
2:H:100:TRP:CH2	2:H:105:PHE:HE1	2.35	0.44
1:A:74:LYS:N	1:A:74:LYS:HD3	2.33	0.43
1:A:77:THR:HG21	1:A:103:LYS:CD	2.46	0.43
2:H:97:ALA:HB1	2:H:106:PHE:HB3	2.00	0.43
2:H:16:VAL:O	2:H:86:LEU:HD12	2.17	0.43
2:H:141:THR:HG22	2:H:142:ALA:N	2.33	0.43
2:H:5:GLN:NE2	2:H:5:GLN:HA	2.33	0.42
2:H:52:ARG:HB3	2:H:54:TYR:CZ	2.55	0.41
3:L:64:GLY:HA2	3:L:72:THR:O	2.21	0.41
2:H:132:PRO:HD3	2:H:144:LEU:HB2	2.02	0.40
2:H:89:GLU:OE1	2:H:89:GLU:HA	2.21	0.40
3:L:190:LYS:O	3:L:210:ASN:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	148/169 (88%)	143 (97%)	4 (3%)	1 (1%)	18	10
2	H	212/241 (88%)	209 (99%)	2 (1%)	1 (0%)	24	16
3	L	212/235 (90%)	206 (97%)	6 (3%)	0	100	100
All	All	572/645 (89%)	558 (98%)	12 (2%)	2 (0%)	30	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
2	H	150	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	128/141 (91%)	126 (98%)	2 (2%)	55	54
2	H	186/209 (89%)	179 (96%)	7 (4%)	29	21
3	L	189/208 (91%)	183 (97%)	6 (3%)	34	27
All	All	503/558 (90%)	488 (97%)	15 (3%)	38	30

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	LYS
1	A	135	SER
2	H	43	GLU
2	H	62	GLN
2	H	89	GLU
2	H	149[A]	LYS
2	H	149[B]	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	184	LEU
2	H	218	GLU
3	L	56	THR
3	L	83	LEU
3	L	105	GLU
3	L	145	LYS
3	L	202	SER
3	L	203	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	5	GLN
2	H	39	GLN
2	H	59	ASN
3	L	38	GLN
3	L	198	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	GOL	H	301	-	5,5,5	0.13	0	5,5,5	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	H	301	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	H	301	GOL	O1-C1-C2-C3
7	H	301	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	150/169 (88%)	0.62	14 (9%) 14 15	20, 35, 63, 83	0
2	H	213/241 (88%)	0.39	11 (5%) 33 35	11, 27, 52, 69	3 (1%)
3	L	211/235 (89%)	0.22	2 (0%) 81 83	14, 30, 42, 52	3 (1%)
All	All	574/645 (88%)	0.39	27 (4%) 36 39	11, 31, 54, 83	6 (1%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	L	10	PHE	4.0
1	A	56	LEU	3.7
1	A	69	ASN	3.7
2	H	133	SER	3.6
1	A	39	LEU	3.5
1	A	77	THR	3.4
2	H	41	HIS	3.4
2	H	197	THR	3.4
2	H	141	THR	3.0
2	H	140	GLY	3.0
2	H	42	ALA	3.0
1	A	47	PHE	2.8
2	H	30	ILE	2.7
1	A	74	LYS	2.6
1	A	70	SER	2.6
2	H	196	GLY	2.6
2	H	142	ALA	2.6
1	A	49	PRO	2.3
1	A	50	ASN	2.3
1	A	76	LEU	2.3
1	A	61	ARG	2.2
2	H	191	PRO	2.2
2	H	54	TYR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	62	TYR	2.1
3	L	154	LEU	2.1
1	A	187	ALA	2.1
1	A	65	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	GOL	H	301	6/6	0.84	0.16	50,51,51,52	0
4	CL	A	201	1/1	0.94	0.10	47,47,47,47	0
6	CA	A	204	1/1	0.98	0.05	30,30,30,30	0
5	ZN	A	202	1/1	0.99	0.03	28,28,28,28	0
6	CA	A	203	1/1	0.99	0.03	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CA A 204:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CA A 203:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.