



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 09:25 PM UTC

PDB ID : 4Z34 / pdb_00004z34
Title : Crystal Structure of Human Lysophosphatidic Acid Receptor 1 in complex with ONO9780307
Authors : Chrencik, J.E.; Roth, C.B.; Terakado, M.; Kurata, H.; Omi, R.; Kihara, Y.; Warshaviak, D.; Nakade, S.; Asmar-Rovira, G.; Mileni, M.; Mizuno, H.; Griffith, M.T.; Rodgers, C.; Han, G.W.; Velasquez, J.; Chun, J.; Stevens, R.C.; Hanson, M.A.; GPCR Network (GPCR)
Deposited on : 2015-03-30
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	8% 70% 11% • 17%

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3047 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysophosphatidic acid receptor 1, Soluble cytochrome b562.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	385	2991	1945	492	529	25	0	0	0

There are 59 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	initiating methionine	UNP Q92633
A	-16	LYS	-	expression tag	UNP Q92633
A	-15	THR	-	expression tag	UNP Q92633
A	-14	ILE	-	expression tag	UNP Q92633
A	-13	ILE	-	expression tag	UNP Q92633
A	-12	ALA	-	expression tag	UNP Q92633
A	-11	LEU	-	expression tag	UNP Q92633
A	-10	SER	-	expression tag	UNP Q92633
A	-9	TYR	-	expression tag	UNP Q92633
A	-8	ILE	-	expression tag	UNP Q92633
A	-7	PHE	-	expression tag	UNP Q92633
A	-6	CYS	-	expression tag	UNP Q92633
A	-5	LEU	-	expression tag	UNP Q92633
A	-4	VAL	-	expression tag	UNP Q92633
A	-3	PHE	-	expression tag	UNP Q92633
A	-2	ALA	-	expression tag	UNP Q92633
A	-1	GLY	-	expression tag	UNP Q92633
A	0	ALA	-	expression tag	UNP Q92633
A	1	PRO	-	expression tag	UNP Q92633
A	1007	TRP	MET	engineered mutation	UNP P0ABE7
A	1043	ALA	-	linker	UNP P0ABE7
A	1044	THR	-	linker	UNP P0ABE7
A	1045	PRO	-	linker	UNP P0ABE7
A	1046	PRO	-	linker	UNP P0ABE7
A	1047	LYS	-	linker	UNP P0ABE7
A	1048	LEU	-	linker	UNP P0ABE7
A	1049	GLU	-	linker	UNP P0ABE7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	1050	ASP	-	linker	UNP P0ABE7
A	1102	ILE	HIS	engineered mutation	UNP P0ABE7
A	1106	LEU	-	linker	UNP P0ABE7
A	327	GLY	-	expression tag	UNP Q92633
A	328	ARG	-	expression tag	UNP Q92633
A	329	PRO	-	expression tag	UNP Q92633
A	330	LEU	-	expression tag	UNP Q92633
A	331	GLU	-	expression tag	UNP Q92633
A	332	VAL	-	expression tag	UNP Q92633
A	333	LEU	-	expression tag	UNP Q92633
A	334	PHE	-	expression tag	UNP Q92633
A	335	GLN	-	expression tag	UNP Q92633
A	336	GLY	-	expression tag	UNP Q92633
A	337	PRO	-	expression tag	UNP Q92633
A	338	HIS	-	expression tag	UNP Q92633
A	339	HIS	-	expression tag	UNP Q92633
A	340	HIS	-	expression tag	UNP Q92633
A	341	HIS	-	expression tag	UNP Q92633
A	342	HIS	-	expression tag	UNP Q92633
A	343	HIS	-	expression tag	UNP Q92633
A	344	HIS	-	expression tag	UNP Q92633
A	345	HIS	-	expression tag	UNP Q92633
A	346	HIS	-	expression tag	UNP Q92633
A	347	HIS	-	expression tag	UNP Q92633
A	348	ASP	-	expression tag	UNP Q92633
A	349	TYR	-	expression tag	UNP Q92633
A	350	LYS	-	expression tag	UNP Q92633
A	351	ASP	-	expression tag	UNP Q92633
A	352	ASP	-	expression tag	UNP Q92633
A	353	ASP	-	expression tag	UNP Q92633
A	354	ASP	-	expression tag	UNP Q92633
A	355	LYS	-	expression tag	UNP Q92633

- Molecule 2 is {1-[(2S,3S)-2-(2,3-dihydro-1H-inden-2-ylmethyl)-3-(3,5-dimethoxy-4-methylphenyl)-3-hydroxypropyl]-4-(methoxycarbonyl)-1 H-pyrrol-3-yl}acetic acid (CCD ID: ON7) (formula: C₃₀H₃₅NO₇).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			38	30	1	7		

- Molecule 3 is (2S)-2,3-dihydroxypropyl (7Z)-tetradec-7-enoate (CCD ID: 1WV) (formula: $C_{17}H_{32}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			18	14	4		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lysophosphatidic acid receptor 1, Soluble cytochrome b562



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	34.28Å 112.15Å 154.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.00 30.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	83.6 (30.00-3.00) 83.5 (30.00-3.00)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 3.00Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.254 , 0.281 0.325 , 0.358	Depositor DCC
R_{free} test set	531 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	52.5	Xtriage
Anisotropy	0.281	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 34.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	3047	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1WV, ON7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	1/3052 (0.0%)	1.47	10/4152 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	181	ILE	CA-CB	5.90	1.57	1.54

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	32	PHE	CA-CB-CG	12.94	126.74	113.80
1	A	302	PHE	CA-CB-CG	9.23	123.03	113.80
1	A	156	HIS	CA-C-N	7.04	133.63	122.36
1	A	156	HIS	C-N-CA	7.04	133.63	122.36
1	A	30	ILE	N-CA-CB	6.33	119.15	110.54
1	A	191	ASP	CA-C-N	6.14	130.70	120.62
1	A	191	ASP	C-N-CA	6.14	130.70	120.62
1	A	180	ALA	CA-C-N	5.25	125.47	120.43
1	A	180	ALA	C-N-CA	5.25	125.47	120.43
1	A	293	GLU	N-CA-CB	5.05	117.31	110.59

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2991	0	2987	27	0
2	A	38	0	34	4	0
3	A	18	0	23	0	0
All	All	3047	0	3044	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:ILE:HD13	1:A:256:LEU:HD21	1.53	0.90
1:A:124:ARG:O	1:A:128:ILE:HD12	1.87	0.75
1:A:208:VAL:CG2	1:A:282:VAL:HG21	2.21	0.70
1:A:84:ILE:HD13	1:A:256:LEU:CD2	2.21	0.69
1:A:61:ILE:O	1:A:65:LEU:HD13	2.00	0.62
1:A:208:VAL:HG23	1:A:282:VAL:HG21	1.81	0.61
1:A:184:VAL:HG12	1:A:184:VAL:O	2.02	0.60
1:A:69:LEU:HD22	1:A:325:ILE:HG21	1.86	0.56
1:A:1075:ALA:HA	1:A:1078:LEU:HD12	1.88	0.56
1:A:297:LEU:HD23	2:A:2001:ON7:H24	1.88	0.55
1:A:29:SER:O	1:A:33:PHE:CD2	2.66	0.48
1:A:26:TYR:O	1:A:29:SER:HB2	2.14	0.48
1:A:258:LYS:O	1:A:261:VAL:HG22	2.13	0.48
1:A:278:LEU:HD11	2:A:2001:ON7:H30	1.94	0.47
1:A:29:SER:HB3	1:A:33:PHE:HE2	1.78	0.47
1:A:84:ILE:CD1	1:A:256:LEU:HD21	2.36	0.47
1:A:297:LEU:CD2	2:A:2001:ON7:H24	2.45	0.46
1:A:221:MET:HA	1:A:224:LEU:HD12	1.98	0.45
1:A:1078:LEU:HD13	1:A:1086:GLU:HB3	1.99	0.45
2:A:2001:ON7:C21	2:A:2001:ON7:H23	2.47	0.44
1:A:1106:LEU:C	1:A:249:ASN:H	2.26	0.44
1:A:277:LEU:HG	1:A:290:LEU:HD23	2.00	0.43
1:A:39:LYS:HG3	1:A:197:ASN:O	2.19	0.42
1:A:171:ILE:O	1:A:174:MET:HG2	2.19	0.41
1:A:24:CYS:O	1:A:189:ILE:HD11	2.21	0.41
1:A:275:LEU:HA	1:A:278:LEU:HD12	2.03	0.41
1:A:37:SER:HB3	1:A:197:ASN:HA	2.03	0.41
1:A:258:LYS:HG3	1:A:259:THR:N	2.35	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	381/464 (82%)	366 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/400 (78%)	292 (93%)	21 (7%)	15	46

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	CYS
1	A	30	ILE
1	A	39	LYS
1	A	51	LEU
1	A	84	ILE
1	A	128	ILE
1	A	133	THR
1	A	149	THR
1	A	190	CYS
1	A	198	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	216	VAL
1	A	217	THR
1	A	1005	ASP
1	A	1010	LEU
1	A	1102	ILE
1	A	264	LEU
1	A	287	CYS
1	A	297	LEU
1	A	315	ASP
1	A	318	MET
1	A	325	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	156	HIS
1	A	214	ASN
1	A	1099	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	1WV	A	2002	-	17,17,20	1.40	2 (11%)	18,18,21	1.18	2 (11%)
2	ON7	A	2001	-	41,41,41	1.74	8 (19%)	48,58,58	1.30	5 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1WV	A	2002	-	-	10/17/17/20	-
2	ON7	A	2001	-	-	12/30/38/38	0/4/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	ON7	C8-C6	5.00	1.60	1.53
2	A	2001	ON7	C30-C31	4.85	1.58	1.51
3	A	2002	1WV	C08-C07	4.08	1.54	1.31
2	A	2001	ON7	C1-C9	2.97	1.44	1.40
2	A	2001	ON7	O4-C28	2.56	1.39	1.33
3	A	2002	1WV	C2-C1	-2.43	1.35	1.51
2	A	2001	ON7	C21-C22	2.38	1.43	1.38
2	A	2001	ON7	C23-N1	2.22	1.40	1.37
2	A	2001	ON7	C13-C8	2.21	1.55	1.52
2	A	2001	ON7	O4-C29	-2.09	1.40	1.45

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	ON7	O3-C7-C9	3.81	117.87	115.03
2	A	2001	ON7	C29-O4-C28	2.99	121.39	115.85
2	A	2001	ON7	C12-C14-C15	-2.69	107.60	115.34
3	A	2002	1WV	C2-C1-C07	2.48	127.55	112.99
3	A	2002	1WV	C3-C2-C1	2.45	126.98	112.50
2	A	2001	ON7	O7-C31-C30	-2.16	115.96	122.11
2	A	2001	ON7	O3-C7-C5	-2.13	120.40	124.08

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2002	1WV	O16-C17-C18-O19
3	A	2002	1WV	C07-C1-C2-C3
3	A	2002	1WV	C1-C07-C08-C09
3	A	2002	1WV	O15-C14-O16-C17
3	A	2002	1WV	C13-C14-O16-C17
2	A	2001	ON7	C2-C1-O1-C4
2	A	2001	ON7	C5-C7-O3-C11
2	A	2001	ON7	C9-C7-O3-C11
2	A	2001	ON7	C9-C1-O1-C4
3	A	2002	1WV	O16-C17-C18-C20
2	A	2001	ON7	C24-C25-C28-O5
2	A	2001	ON7	C24-C25-C28-O4
2	A	2001	ON7	C14-C12-C8-C6
3	A	2002	1WV	C07-C08-C09-C10
2	A	2001	ON7	C24-C30-C31-O7
2	A	2001	ON7	C24-C30-C31-O6
2	A	2001	ON7	C26-C25-C28-O5
2	A	2001	ON7	C26-C25-C28-O4
3	A	2002	1WV	C08-C07-C1-C2
3	A	2002	1WV	C12-C13-C14-O16
2	A	2001	ON7	C14-C12-C8-C13
3	A	2002	1WV	C12-C13-C14-O15


There are no ring outliers.


1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2001	ON7	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	385/464 (82%)	0.81	37 (9%) 13 7	33, 74, 120, 156	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	27	ASN	3.8
1	A	327	GLY	3.7
1	A	305	ALA	3.7
1	A	26	TYR	3.5
1	A	189	ILE	3.5
1	A	113	THR	3.3
1	A	84	ILE	3.3
1	A	28	GLU	3.3
1	A	1083	LYS	3.2
1	A	1086	GLU	3.2
1	A	187	ASN	3.2
1	A	192	ILE	3.1
1	A	160	SER	3.1
1	A	45	TRP	2.9
1	A	248	ARG	2.8
1	A	190	CYS	2.8
1	A	194	ASN	2.7
1	A	200	PRO	2.7
1	A	211	ALA	2.7
1	A	181	ILE	2.6
1	A	191	ASP	2.4
1	A	41	LEU	2.4
1	A	322	PHE	2.3
1	A	325	ILE	2.3
1	A	1101	TYR	2.2
1	A	287	CYS	2.2
1	A	318	MET	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	157	THR	2.2
1	A	275	LEU	2.1
1	A	145	GLU	2.1
1	A	1081	GLU	2.1
1	A	87	LEU	2.1
1	A	289	VAL	2.1
1	A	161	ASN	2.1
1	A	1059	LYS	2.0
1	A	85	TYR	2.0
1	A	295	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	1WV	A	2002	18/21	0.89	0.12	55,63,66,68	0
2	ON7	A	2001	38/38	0.91	0.12	39,44,64,66	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 1WV A 2002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ON7 A 2001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.