



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:34 AM UTC

PDB ID : 4Z4K / pdb_00004z4k
Title : Crystal structure of GFP-TAX1BP1 UBZ1+2 domain fusion protein
Authors : Rohaim, A.; Kawasaki, M.; Wakatsuki, S.
Deposited on : 2015-04-02
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	74% 21% ..
1	B	290	78% 17% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4642 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein,Tax1-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	0	0
			2293	1460	386	435	12			
1	B	282	Total	C	N	O	S	0	0	0
			2287	1457	385	433	12			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P42212
A	-1	SER	-	expression tag	UNP P42212
A	0	HIS	-	expression tag	UNP P42212
A	66	CR2	SER	chromophore	UNP P42212
A	66	CR2	TYR	chromophore	UNP P42212
A	66	CR2	GLY	chromophore	UNP P42212
A	72	ALA	SER	engineered mutation	UNP P42212
A	231	GLY	-	linker	UNP P42212
A	232	SER	-	linker	UNP P42212
B	-2	GLY	-	expression tag	UNP P42212
B	-1	SER	-	expression tag	UNP P42212
B	0	HIS	-	expression tag	UNP P42212
B	66	CR2	SER	chromophore	UNP P42212
B	66	CR2	TYR	chromophore	UNP P42212
B	66	CR2	GLY	chromophore	UNP P42212
B	72	ALA	SER	engineered mutation	UNP P42212
B	231	GLY	-	linker	UNP P42212
B	232	SER	-	linker	UNP P42212

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Zn	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	39	Total	O	0	0
			39	39		
3	B	19	Total	O	0	0
			19	19		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein,Tax1-binding protein 1



- Molecule 1: Green fluorescent protein,Tax1-binding protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.88Å 83.33Å 128.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 50.00 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.6 (50.00-2.80) 99.6 (50.00-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.87 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, R_{free}	0.206 , 0.280 0.205 , 0.281	Depositor DCC
R_{free} test set	778 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	41.7	Xtriage
Anisotropy	0.413	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 26.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4642	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CR2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/2334	0.93	1/3155 (0.0%)
1	B	0.74	0/2328	0.94	3/3147 (0.1%)
All	All	0.78	0/4662	0.93	4/6302 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	158	LYS	N-CA-C	-9.71	95.17	109.63
1	B	186	THR	CB-CA-C	5.58	117.58	109.20
1	B	241	CYS	N-CA-C	-5.18	102.50	109.95
1	A	197	ASP	N-CA-C	-5.13	103.62	110.55

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2293	0	2180	32	0
1	B	2287	0	2175	29	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	39	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	19	0	0	1	0
All	All	4642	0	4355	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:VAL:HG22	1:A:235:HIS:H	1.53	0.70
1:A:90:GLU:HG2	3:A:405:HOH:O	1.92	0.68
1:A:162:LYS:HD3	1:A:184:GLN:HG2	1.78	0.66
1:A:153:MET:HE1	1:A:258:VAL:HG11	1.78	0.66
1:A:88:MET:HE3	1:A:89:PRO:HA	1.78	0.65
1:B:262:TRP:HB2	1:B:271:GLN:HG2	1.79	0.64
1:B:207:LEU:HD13	1:B:218:MET:HE2	1.80	0.64
1:A:221:LEU:HD22	1:A:223:PHE:CE1	2.33	0.64
1:A:164:ASN:HD21	1:A:262:TRP:HE1	1.44	0.62
1:B:88:MET:HE3	1:B:89:PRO:HA	1.82	0.61
1:A:237:LYS:HE2	1:A:242:GLU:HA	1.83	0.61
1:A:207:LEU:HD13	1:A:218:MET:HE2	1.81	0.61
1:A:155:ASP:OD2	1:A:162:LYS:NZ	2.33	0.59
1:B:88:MET:HE1	1:B:113:LYS:HA	1.85	0.58
1:B:207:LEU:HD13	1:B:218:MET:CE	2.36	0.56
1:A:207:LEU:HD13	1:A:218:MET:CE	2.36	0.56
1:A:112:VAL:HA	1:A:120:VAL:O	2.07	0.55
1:B:57:TRP:HB3	1:B:218:MET:HE3	1.89	0.54
1:B:112:VAL:HA	1:B:120:VAL:O	2.06	0.54
1:B:233:ASP:O	1:B:248:ASN:ND2	2.41	0.54
1:B:3:LYS:HB2	1:B:7:LEU:HD21	1.88	0.54
1:A:88:MET:HE1	1:A:113:LYS:HA	1.90	0.53
1:A:125:LEU:C	1:A:125:LEU:HD23	2.33	0.53
1:A:2:SER:N	3:A:402:HOH:O	2.42	0.53
1:A:203:THR:HG22	1:A:224:VAL:HG13	1.90	0.52
1:B:125:LEU:C	1:B:125:LEU:HD23	2.35	0.52
1:B:234:VAL:HG13	1:B:247:PRO:HB3	1.91	0.51
1:B:203:THR:HG22	1:B:224:VAL:HG13	1.91	0.51
1:A:267:MET:N	1:A:267:MET:HE2	2.26	0.51
1:A:162:LYS:CD	1:A:184:GLN:HG2	2.39	0.50
1:B:262:TRP:CB	1:B:271:GLN:HG2	2.41	0.49
1:A:232:SER:OG	1:A:236:LYS:HE3	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29:VAL:HG11	1:A:64:PHE:CZ	2.48	0.48
1:B:5:GLU:HG2	3:B:406:HOH:O	2.13	0.48
1:B:119:LEU:C	1:B:119:LEU:HD13	2.39	0.48
1:A:57:TRP:HB3	1:A:218:MET:HE3	1.96	0.48
1:A:2:SER:N	1:A:5:GLU:OE2	2.47	0.47
1:A:119:LEU:C	1:A:119:LEU:HD13	2.39	0.47
1:B:237:LYS:HE2	1:B:242:GLU:HA	1.97	0.47
1:B:237:LYS:NZ	1:B:242:GLU:OE2	2.47	0.47
1:A:92:TYR:CE1	1:A:112:VAL:HG13	2.50	0.46
1:B:171:ILE:HD12	1:B:175:SER:OG	2.16	0.46
1:B:92:TYR:CE1	1:B:112:VAL:HG13	2.51	0.46
1:A:151:TYR:CD1	1:A:200:TYR:HB3	2.50	0.46
1:A:42:LEU:HB3	1:A:222:GLU:HB3	1.98	0.45
1:A:57:TRP:HB2	1:A:58:PRO:HD3	1.99	0.45
1:B:3:LYS:HB2	1:B:7:LEU:CD2	2.47	0.45
1:A:141:LEU:HD22	1:A:171:ILE:CD1	2.47	0.44
1:A:173:ASP:HB2	1:B:235:HIS:HB3	1.98	0.44
1:B:43:THR:HG23	1:B:221:LEU:HD22	2.00	0.44
1:B:57:TRP:HB3	1:B:218:MET:CE	2.48	0.44
1:B:284:HIS:O	1:B:287:THR:OG1	2.29	0.43
1:A:264:VAL:HG12	1:A:271:GLN:HG3	2.00	0.42
1:A:247:PRO:O	1:A:248:ASN:C	2.62	0.42
1:B:92:TYR:CE1	1:B:112:VAL:CG1	3.03	0.42
1:B:166:LYS:HD3	1:B:178:LEU:HB3	2.02	0.42
1:A:92:TYR:CE1	1:A:112:VAL:CG1	3.02	0.42
1:B:57:TRP:HB2	1:B:58:PRO:HD3	2.03	0.41
1:B:166:LYS:HE3	1:B:240:LEU:O	2.20	0.40
1:B:66:CR2:N2	1:B:66:CR2:HD1	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/290 (95%)	268 (97%)	8 (3%)	0	100	100
1	B	275/290 (95%)	265 (96%)	10 (4%)	0	100	100
All	All	551/580 (95%)	533 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/258 (98%)	225 (89%)	28 (11%)	6	20
1	B	252/258 (98%)	229 (91%)	23 (9%)	9	28
All	All	505/516 (98%)	454 (90%)	51 (10%)	7	23

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LYS
1	A	6	GLU
1	A	9	THR
1	A	15	LEU
1	A	17	GLU
1	A	32	GLU
1	A	43	THR
1	A	44	LEU
1	A	52	LYS
1	A	101	LYS
1	A	107	LYS
1	A	112	VAL
1	A	113	LYS
1	A	123	ILE
1	A	153	MET
1	A	156	LYS
1	A	157	GLN
1	A	158	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	186	THR
1	A	209	LYS
1	A	221	LEU
1	A	225	THR
1	A	242	GLU
1	A	243	LEU
1	A	267	MET
1	A	271	GLN
1	A	283	ARG
1	A	287	THR
1	B	9	THR
1	B	15	LEU
1	B	42	LEU
1	B	43	THR
1	B	44	LEU
1	B	52	LYS
1	B	80	GLN
1	B	101	LYS
1	B	112	VAL
1	B	123	ILE
1	B	126	LYS
1	B	156	LYS
1	B	158	LYS
1	B	166	LYS
1	B	184	GLN
1	B	190	ASP
1	B	221	LEU
1	B	225	THR
1	B	233	ASP
1	B	251	GLN
1	B	267	MET
1	B	279	GLN
1	B	286	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	135	ASN
1	A	149	ASN
1	A	157	GLN
1	A	164	ASN
1	A	170	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	198	ASN
1	A	248	ASN
1	A	251	GLN
1	A	271	GLN
1	A	286	GLN
1	B	80	GLN
1	B	170	ASN
1	B	212	ASN
1	B	235	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CR2	A	66	1	20,20,21	3.52	4 (20%)	25,27,29	4.57	6 (24%)
1	CR2	B	66	1	20,20,21	3.60	4 (20%)	25,27,29	3.89	10 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CR2	A	66	1	-	0/6/25/26	0/2/2/2
1	CR2	B	66	1	-	0/6/25/26	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	66	CR2	CB2-CA2	14.64	1.49	1.35
1	A	66	CR2	CB2-CA2	13.91	1.48	1.35
1	A	66	CR2	CA2-C2	-4.45	1.43	1.48
1	B	66	CR2	C2-N3	-3.74	1.31	1.40
1	A	66	CR2	C2-N3	-3.56	1.31	1.40
1	B	66	CR2	C1-N2	3.47	1.38	1.32
1	A	66	CR2	C1-N2	2.75	1.37	1.32
1	B	66	CR2	CA1-C1	-2.04	1.47	1.49

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CR2	O2-C2-CA2	-15.28	121.27	131.02
1	A	66	CR2	C2-N3-C1	-11.40	102.95	108.08
1	B	66	CR2	C2-N3-C1	-10.97	103.14	108.08
1	A	66	CR2	CA2-C2-N3	10.39	112.23	103.50
1	B	66	CR2	CA2-C2-N3	9.72	111.66	103.50
1	B	66	CR2	O2-C2-CA2	-6.88	126.63	131.02
1	B	66	CR2	CB2-CA2-C2	5.20	128.66	122.36
1	B	66	CR2	C1-CA1-N1	-5.16	101.44	112.85
1	B	66	CR2	C2-CA2-N2	-5.01	105.36	108.95
1	A	66	CR2	C2-CA2-N2	-4.72	105.57	108.95
1	B	66	CR2	N3-C1-N2	3.00	115.22	111.75
1	B	66	CR2	CG2-CB2-CA2	-2.90	126.42	129.87
1	B	66	CR2	C3-CA3-N3	2.74	118.67	112.43
1	B	66	CR2	CB2-CA2-N2	-2.04	125.98	128.76
1	A	66	CR2	CA1-C1-N2	-2.01	121.58	124.28
1	A	66	CR2	CA1-C1-N3	2.01	125.21	122.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	66	CR2	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/290 (97%)	-0.23	3 (1%) 78 70	14, 29, 72, 97	0
1	B	281/290 (96%)	0.05	3 (1%) 78 70	24, 52, 83, 119	0
All	All	563/580 (97%)	-0.09	6 (1%) 78 70	14, 43, 81, 119	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	232	SER	3.1
1	B	228	GLY	2.6
1	A	279	GLN	2.6
1	B	276	TYR	2.5
1	B	289	PHE	2.3
1	A	129	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CR2	A	66	19/20	0.92	0.08	17,18,20,20	0
1	CR2	B	66	19/20	0.94	0.08	29,30,39,41	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	B	302	1/1	0.97	0.04	75,75,75,75	0
2	ZN	B	301	1/1	0.99	0.04	52,52,52,52	0
2	ZN	A	302	1/1	0.99	0.04	64,64,64,64	0
2	ZN	A	301	1/1	1.00	0.05	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.