



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:33 PM UTC

PDB ID : 5Z4P / pdb_00005z4p
Title : Crystal structure of tubulin-stathmin-TTL-Compound TCA complex
Authors : Zhang, H.; Luo, C.
Deposited on : 2018-01-12
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	4% 70% 29%
1	C	440	3% 77% 23%
2	B	431	4% 69% 29% .
2	D	431	5% 62% 34% ..
3	E	185	5% 45% 20% 35%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	378	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
13	ACP	F	401	-	-	X	-
7	GOL	C	505	-	-	X	-

2 Entry composition [i](#)

There are 14 unique types of molecules in this entry. The entry contains 17477 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	438	Total	C	N	O	S	0	2	0
			3433	2175	582	653	23			
1	C	440	Total	C	N	O	S	0	3	0
			3451	2184	586	658	23			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	424	Total	C	N	O	S	0	2	0
			3347	2104	568	648	27			
2	D	421	Total	C	N	O	S	0	0	0
			3309	2080	562	640	27			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	121	Total	C	N	O	S	0	2	0
			1011	624	183	199	5			

- Molecule 4 is a protein called Tubulin tyrosine ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	317	Total	C	N	O	S	0	0	0
			2596	1677	434	471	14			

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total 32	C 10	N 5	O 14	P 3	0	0
5	C	1	Total 32	C 10	N 5	O 14	P 3	0	0

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Mg 1 1	0	0
6	B	1	Total Mg 1 1	0	0
6	C	1	Total Mg 1 1	0	0
6	D	1	Total Mg 1 1	0	0

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Ca	0	0
			1	1		
8	C	1	Total	Ca	0	0
			1	1		


- Molecule 9 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).






Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 11 is 6,7,8-trimethoxy-1-(4-methoxyphenyl)-4,5-dihydro-2 {H}-benzo[e]indazole (CCD ID: 97O) (formula: $C_{21}H_{22}N_2O_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	B	1	Total	C	N	O	0	0
			27	21	2	4		

- Molecule 12 is IMIDAZOLE (CCD ID: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	C	1	Total	C	N	0	0
			5	3	2		
12	C	1	Total	C	N	0	0
			5	3	2		
12	C	1	Total	C	N	0	0
			5	3	2		

- Molecule 13 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
13	F	1	Total	C	N	O	P	0	0
			31	11	5	12	3		


- Molecule 14 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	A	11	Total	O	0	0
			11	11		
14	B	13	Total	O	0	0
			13	13		
14	C	17	Total	O	0	0
			17	17		
14	D	4	Total	O	0	0
			4	4		
14	F	7	Total	O	0	0
			7	7		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tubulin alpha-1B chain




- Molecule 1: Tubulin alpha-1B chain




- Molecule 2: Tubulin beta-2B chain







• Molecule 2: Tubulin beta-2B chain



• Molecule 3: Stathmin-4



• Molecule 4: Tubulin tyrosine ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	103.83Å 154.84Å 181.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.64 – 2.50 49.64 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.64-2.50) 99.7 (49.64-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.05 (at 2.48Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.222 , 0.286 0.223 , 0.287	Depositor DCC
R_{free} test set	2000 reflections (0.80%)	wwPDB-VP
Wilson B-factor (Å ²)	61.3	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	17477	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, MG, GOL, IMD, MES, GDP, CA, 97O, ACP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/3517	0.59	0/4774
1	C	0.46	0/3538	0.65	2/4804 (0.0%)
2	B	0.43	0/3424	0.66	0/4638
2	D	0.39	0/3382	0.63	0/4581
3	E	0.44	0/1025	0.60	0/1360
4	F	0.37	0/2652	0.62	0/3579
All	All	0.42	0/17538	0.63	2/23736 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	C	171	ILE	CA-C-N	8.07	136.28	120.94
1	C	171	ILE	C-N-CA	8.07	136.28	120.94

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	247	GLN	Peptide
2	D	93	VAL	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	D	95	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3433	0	3352	103	0
1	C	3451	0	3368	83	0
2	B	3347	0	3220	106	0
2	D	3309	0	3189	129	0
3	E	1011	0	1032	42	0
4	F	2596	0	2570	132	0
5	A	32	0	12	2	0
5	C	32	0	12	1	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	18	0	24	1	0
7	B	6	0	8	0	0
7	C	30	0	40	8	0
7	D	12	0	16	0	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
9	B	28	0	12	1	0
9	D	28	0	12	1	0
10	B	12	0	12	0	0
11	B	27	0	0	2	0
12	C	15	0	15	5	0
13	F	31	0	12	13	0
14	A	11	0	0	2	0
14	B	13	0	0	2	0
14	C	17	0	0	1	0
14	D	4	0	0	0	0
14	F	7	0	0	1	0
All	All	17477	0	16906	573	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 573 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
13:F:401:ACP:O3'	13:F:401:ACP:C3'	1.80	1.30
4:F:241:THR:N	13:F:401:ACP:O3'	1.73	1.21
3:E:103:GLN:OE1	3:E:104:LYS:CE	1.95	1.14
4:F:148:ILE:HA	4:F:162:ILE:HD12	1.30	1.09
4:F:241:THR:HG22	13:F:401:ACP:HO3'	1.24	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/440 (100%)	412 (94%)	24 (6%)	2 (0%)	24	43
1	C	441/440 (100%)	428 (97%)	13 (3%)	0	100	100
2	B	422/431 (98%)	395 (94%)	25 (6%)	2 (0%)	24	43
2	D	417/431 (97%)	387 (93%)	23 (6%)	7 (2%)	7	13
3	E	119/185 (64%)	112 (94%)	7 (6%)	0	100	100
4	F	299/378 (79%)	246 (82%)	47 (16%)	6 (2%)	6	10
All	All	2136/2305 (93%)	1980 (93%)	139 (6%)	17 (1%)	16	31

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	57	THR
2	B	82	PRO
2	D	177	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	F	254	GLY
1	A	177	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	371/371 (100%)	368 (99%)	3 (1%)	73	88
1	C	374/371 (101%)	372 (100%)	2 (0%)	81	92
2	B	368/372 (99%)	368 (100%)	0	100	100
2	D	364/372 (98%)	363 (100%)	1 (0%)	86	94
3	E	111/168 (66%)	111 (100%)	0	100	100
4	F	286/336 (85%)	285 (100%)	1 (0%)	86	94
All	All	1874/1990 (94%)	1867 (100%)	7 (0%)	84	93

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	241	SER
1	C	381	THR
4	F	12	SER
2	D	248	LEU
1	A	419	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
3	E	92	ASN
3	E	12	ASN
2	D	50	ASN
2	D	249	ASN
2	D	15	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 28 ligands modelled in this entry, 7 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	GDP	D	501	6	29,30,30	1.25	3 (10%)	45,47,47	1.84	8 (17%)
12	IMD	C	509	-	5,5,5	0.76	0	5,5,5	0.56	0
5	GTP	C	503	6	33,34,34	0.88	2 (6%)	50,54,54	1.70	11 (22%)
10	MES	B	505	-	12,12,12	2.28	1 (8%)	15,16,16	1.87	3 (20%)
7	GOL	A	506	-	5,5,5	1.17	0	5,5,5	0.94	0
7	GOL	C	501	-	5,5,5	0.63	0	5,5,5	1.42	1 (20%)
7	GOL	A	503	-	5,5,5	1.01	0	5,5,5	0.89	0
11	97O	B	506	-	29,30,30	6.24	21 (72%)	38,43,43	3.45	16 (42%)
7	GOL	C	506	-	5,5,5	1.63	2 (40%)	5,5,5	0.78	0
7	GOL	D	504	-	5,5,5	0.88	0	5,5,5	1.21	0
5	GTP	A	501	6	33,34,34	0.97	2 (6%)	50,54,54	1.71	11 (22%)
7	GOL	B	503	-	5,5,5	1.26	1 (20%)	5,5,5	0.70	0
7	GOL	C	507	-	5,5,5	1.43	1 (20%)	5,5,5	0.82	0
7	GOL	D	503	-	5,5,5	1.04	0	5,5,5	1.04	0
7	GOL	C	505	-	5,5,5	0.92	0	5,5,5	1.06	1 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
12	IMD	C	511	-	5,5,5	0.80	0	5,5,5	0.38	0
7	GOL	C	502	-	5,5,5	1.20	0	5,5,5	0.93	0
12	IMD	C	510	-	5,5,5	0.65	0	5,5,5	0.71	0
9	GDP	B	501	6	29,30,30	1.32	3 (10%)	45,47,47	1.80	7 (15%)
13	ACP	F	401	-	31,33,33	4.57	13 (41%)	47,52,52	3.10	15 (31%)
7	GOL	A	504	-	5,5,5	1.51	2 (40%)	5,5,5	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	GDP	D	501	6	-	3/16/32/32	0/3/3/3
12	IMD	C	509	-	-	-	0/1/1/1
5	GTP	C	503	6	-	8/22/38/38	0/3/3/3
10	MES	B	505	-	-	4/6/14/14	0/1/1/1
7	GOL	A	506	-	-	0/4/4/4	-
7	GOL	C	501	-	-	2/4/4/4	-
7	GOL	A	503	-	-	2/4/4/4	-
11	97O	B	506	-	-	2/12/21/21	0/4/4/4
7	GOL	C	506	-	-	2/4/4/4	-
7	GOL	D	504	-	-	4/4/4/4	-
5	GTP	A	501	6	-	6/22/38/38	0/3/3/3
7	GOL	B	503	-	-	0/4/4/4	-
7	GOL	C	507	-	-	2/4/4/4	-
7	GOL	D	503	-	-	1/4/4/4	-
7	GOL	C	505	-	-	2/4/4/4	-
12	IMD	C	511	-	-	-	0/1/1/1
7	GOL	C	502	-	-	4/4/4/4	-
12	IMD	C	510	-	-	-	0/1/1/1
9	GDP	B	501	6	-	5/16/32/32	0/3/3/3
13	ACP	F	401	-	-	4/19/38/38	0/3/3/3
7	GOL	A	504	-	-	2/4/4/4	-

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	B	506	97O	C3-C2	-15.77	1.08	1.41
13	F	401	ACP	O3'-C3'	14.97	1.80	1.43
11	B	506	97O	C8-C9	13.99	1.72	1.50
13	F	401	ACP	C2'-C3'	-12.88	1.18	1.53
11	B	506	97O	C4-C3	-11.62	1.19	1.38

The worst 5 of 73 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	B	506	97O	C5-C6-C1	-8.15	106.49	118.42
13	F	401	ACP	C4-N9-C8	8.05	114.19	105.74
11	B	506	97O	C4-C3-C2	7.42	128.47	120.22
13	F	401	ACP	O3'-C3'-C4'	7.37	132.25	111.08
13	F	401	ACP	C2'-C3'-C4'	6.96	116.07	102.61

There are no chirality outliers.

5 of 53 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	501	GTP	PB-O3B-PG-O3G
5	A	501	GTP	C5'-O5'-PA-O3A
5	A	501	GTP	C5'-O5'-PA-O1A
5	A	501	GTP	C5'-O5'-PA-O2A
5	C	503	GTP	PB-O3B-PG-O3G


There are no ring outliers.

12 monomers are involved in 34 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	D	501	GDP	1	0
5	C	503	GTP	1	0
7	C	501	GOL	2	0
7	A	503	GOL	1	0
11	B	506	97O	2	0
7	C	506	GOL	1	0
5	A	501	GTP	2	0
7	C	505	GOL	5	0
12	C	511	IMD	3	0
12	C	510	IMD	2	0
9	B	501	GDP	1	0
13	F	401	ACP	13	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






Ligand 97O B 506







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	438/440 (99%)	0.38	17 (3%) 43 38	44, 63, 79, 146	2 (0%)
1	C	440/440 (100%)	0.07	15 (3%) 48 43	37, 51, 70, 149	3 (0%)
2	B	424/431 (98%)	0.31	16 (3%) 44 39	38, 58, 91, 135	2 (0%)
2	D	421/431 (97%)	0.61	22 (5%) 33 29	46, 70, 102, 118	0
3	E	121/185 (65%)	0.62	10 (8%) 17 15	43, 71, 94, 148	2 (1%)
4	F	317/378 (83%)	1.16	67 (21%) 2 2	52, 81, 247, 320	0
All	All	2161/2305 (93%)	0.47	147 (6%) 23 20	37, 63, 117, 320	9 (0%)

The worst 5 of 147 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	68	VAL	5.6
2	D	249	ASN	5.4
4	F	182	ILE	5.2
4	F	170	LEU	5.1
4	F	173	ILE	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
13	ACP	F	401	31/31	0.59	0.14	86,205,221,231	0
7	GOL	A	504	6/6	0.75	0.19	62,74,81,82	0
7	GOL	A	506	6/6	0.76	0.16	68,71,73,76	0
7	GOL	C	506	6/6	0.77	0.20	63,69,69,69	0
7	GOL	D	503	6/6	0.78	0.17	82,90,92,96	0
7	GOL	C	501	6/6	0.81	0.25	50,56,66,76	0
6	MG	D	502	1/1	0.82	0.19	76,76,76,76	0
12	IMD	C	510	5/5	0.84	0.17	61,65,71,74	0
7	GOL	A	503	6/6	0.84	0.14	67,72,75,79	0
7	GOL	B	503	6/6	0.85	0.18	57,71,76,89	0
7	GOL	C	502	6/6	0.86	0.20	64,68,73,78	0
7	GOL	D	504	6/6	0.87	0.17	67,68,75,79	0
7	GOL	C	507	6/6	0.87	0.13	56,63,66,66	0
6	MG	A	502	1/1	0.87	0.31	59,59,59,59	0
11	97O	B	506	27/27	0.88	0.14	58,73,78,80	0
12	IMD	C	509	5/5	0.89	0.13	51,53,67,67	0
12	IMD	C	511	5/5	0.90	0.25	65,69,75,76	0
7	GOL	C	505	6/6	0.91	0.19	66,72,74,78	0
6	MG	C	504	1/1	0.91	0.16	50,50,50,50	0
9	GDP	D	501	28/28	0.92	0.11	66,71,81,92	0
6	MG	B	502	1/1	0.92	0.18	51,51,51,51	0
8	CA	A	505	1/1	0.94	0.08	82,82,82,82	0
10	MES	B	505	12/12	0.95	0.08	45,54,69,70	0
8	CA	B	504	1/1	0.96	0.08	97,97,97,97	0
8	CA	C	508	1/1	0.96	0.07	85,85,85,85	0
5	GTP	A	501	32/32	0.96	0.07	46,53,58,60	0
9	GDP	B	501	28/28	0.97	0.07	38,47,50,58	0
5	GTP	C	503	32/32	0.97	0.07	35,44,50,57	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ACP F 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 97O B 506:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GTP A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GTP C 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.