



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:50 AM UTC

PDB ID : 6Z45 / pdb_00006z45
Title : CDK9-Cyclin-T1 complex bound by compound 24
Authors : Ferguson, A.; Collie, G.W.
Deposited on : 2020-05-22
Resolution : 3.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1047 (3.42-3.34)
Clashscore	190562	1067 (3.42-3.34)
Ramachandran outliers	187476	1056 (3.42-3.34)
Sidechain outliers	187428	1056 (3.42-3.34)
RSRZ outliers	180081	1047 (3.42-3.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	
2	B	258	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4646 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 9.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	314	2546	1631	438	460	1	16	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ASP	SER	conflict	UNP P50750
A	8	ASN	VAL	conflict	UNP P50750
A	44	ARG	LYS	conflict	UNP P50750
A	138	PHE	TYR	conflict	UNP P50750
A	181	ASN	GLN	conflict	UNP P50750
A	280	ALA	LYS	conflict	UNP P50750
A	307	GLU	ASP	conflict	UNP P50750
A	311	GLU	ASN	conflict	UNP P50750

- Molecule 2 is a protein called Cyclin-T1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	252	2057	1315	359	374	9	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	26	ALA	ARG	conflict	UNP O60563
B	77	ARG	GLN	conflict	UNP O60563
B	96	GLY	GLU	conflict	UNP O60563
B	106	ARG	LYS	conflict	UNP O60563
B	241	LEU	PHE	conflict	UNP O60563


- Molecule 3 is (1 {S},3 {R})-3-acetamido- {N}-[5-chloranyl-4-(5,5-dimethyl-4,6-dihydropyrrlo[1,2-b]pyrazol-3-yl)pyridin-2-yl]cyclohexane-1-carboxamide (CCD ID: Q6E) (formula:

C₂₂H₂₈ClN₅O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0
			30	22	1	5	2	

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 5 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	P	0	0
			5	4	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cyclin-dependent kinase 9



• Molecule 2: Cyclin-T1



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	171.76Å 171.76Å 97.79Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	85.88 – 3.37 85.88 – 3.37	Depositor EDS
% Data completeness (in resolution range)	99.2 (85.88-3.37) 99.5 (85.88-3.37)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 3.40Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.173 , 0.188 (Not available) , 0.191	Depositor DCC
R_{free} test set	591 reflections (3.92%)	wwPDB-VP
Wilson B-factor (Å ²)	141.1	Xtriage
Anisotropy	0.311	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 138.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4646	wwPDB-VP
Average B, all atoms (Å ²)	160.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, TRS, PO4, Q6E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.83	0/2585	1.27	12/3486 (0.3%)
2	B	0.81	0/2110	1.21	5/2875 (0.2%)
All	All	0.82	0/4695	1.25	17/6361 (0.3%)

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	GLU	CB-CG-CD	8.66	127.33	112.60
2	B	30	ASP	CA-CB-CG	7.58	120.18	112.60
1	A	50	VAL	N-CA-CB	7.36	119.39	111.00
2	B	102	GLU	CB-CG-CD	7.03	124.55	112.60
1	A	98	SER	N-CA-C	6.69	118.64	110.41
1	A	324	LEU	CA-C-N	6.47	133.26	121.41
1	A	324	LEU	C-N-CA	6.47	133.26	121.41
1	A	6	ASP	CA-CB-CG	6.39	118.99	112.60
1	A	30	PHE	CA-CB-CG	-6.01	107.79	113.80
2	B	217	ASP	CA-CB-CG	5.79	118.39	112.60
1	A	257	ASP	CA-CB-CG	5.77	118.37	112.60
2	B	168	LYS	N-CA-C	-5.48	104.53	111.11
1	A	87	THR	CB-CA-C	5.36	118.24	110.79
1	A	50	VAL	CB-CA-C	5.28	117.35	110.96
2	B	71	MET	N-CA-C	-5.19	106.95	113.28
1	A	28	GLY	CA-C-N	5.09	131.26	121.54
1	A	28	GLY	C-N-CA	5.09	131.26	121.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2546	0	2582	39	0
2	B	2057	0	2041	17	0
3	A	30	0	0	0	0
4	B	8	0	12	0	0
5	B	5	0	0	0	0
All	All	4646	0	4635	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:327:MET:HB2	1:A:328:LEU:HA	1.43	1.00
1:A:132:MET:HE1	1:A:162:VAL:HA	1.55	0.89
1:A:224:THR:HG22	1:A:282:TYR:CD2	2.10	0.85
1:A:55:GLU:HG2	1:A:60:PRO:HG3	1.58	0.85
1:A:261:LEU:HA	1:A:267:LEU:HD21	1.66	0.78
1:A:249:THR:HG22	1:A:251:GLU:H	1.52	0.75
1:A:248:ILE:HG22	1:A:256:VAL:HG21	1.69	0.73
2:B:44:LEU:HG	2:B:48:MET:HE2	1.76	0.66
1:A:196:PRO:HB2	1:A:198:GLU:OE1	1.95	0.66
1:A:144:LYS:HE2	1:A:176:LEU:HD11	1.80	0.64
1:A:189:VAL:HG23	1:A:199:LEU:HD13	1.79	0.63
1:A:327:MET:CB	1:A:328:LEU:HA	2.26	0.60
2:B:164:VAL:HG22	2:B:212:ILE:HG21	1.83	0.60
1:A:248:ILE:CG2	1:A:256:VAL:HG21	2.32	0.57
1:A:224:THR:HG22	1:A:282:TYR:HD2	1.71	0.55
2:B:166:ALA:HB1	2:B:170:LEU:HB3	1.90	0.53
1:A:107:GLU:HG2	1:A:159:ARG:HB3	1.89	0.53
1:A:268:VAL:HG21	1:A:271:GLN:HG3	1.91	0.53
1:A:260:GLU:HA	1:A:263:GLU:HB3	1.92	0.52
2:B:105:ILE:CD1	2:B:130:VAL:HA	2.40	0.52
1:A:242:SER:HB3	1:A:268:VAL:HG21	1.95	0.49
1:A:327:MET:HB2	1:A:328:LEU:CA	2.28	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:GLU:HG2	1:A:70:LEU:HD22	1.95	0.47
1:A:242:SER:HA	1:A:246:GLY:O	2.14	0.47
1:A:38:HIS:CE1	1:A:40:LYS:HB2	2.49	0.47
1:A:86:ARG:HA	1:A:99:ILE:HD13	1.96	0.47
2:B:160:CYS:SG	2:B:201:ILE:HD13	2.56	0.46
2:B:113:HIS:HB3	2:B:116:GLU:HB2	1.97	0.46
2:B:120:ASP:OD2	2:B:122:ARG:HD3	2.16	0.46
2:B:258:TRP:CD1	2:B:258:TRP:H	2.34	0.46
2:B:105:ILE:HD11	2:B:130:VAL:HA	1.98	0.46
1:A:249:THR:HG22	1:A:251:GLU:HB3	1.99	0.45
2:B:92:ALA:HB1	2:B:97:GLN:O	2.16	0.45
2:B:118:LEU:HD12	2:B:119:PRO:HD2	1.98	0.45
2:B:84:ALA:HB3	2:B:85:PRO:HD3	1.97	0.45
1:A:44:ARG:HH22	1:A:107:GLU:HG3	1.82	0.45
1:A:183:ASN:HB2	1:A:205:ASP:OD1	2.17	0.45
1:A:249:THR:H	1:A:252:VAL:HG22	1.82	0.45
1:A:240:LEU:HD23	1:A:243:GLN:NE2	2.32	0.44
1:A:276:LYS:O	1:A:280:ALA:HB2	2.16	0.44
1:A:254:PRO:O	1:A:255:ASN:HB2	2.18	0.44
2:B:126:TYR:HA	2:B:129:GLN:HE21	1.83	0.44
1:A:272:LYS:O	1:A:274:LYS:HG3	2.17	0.43
2:B:19:LEU:HD11	2:B:190:GLN:HG2	1.99	0.43
2:B:121:THR:HA	2:B:126:TYR:CD2	2.52	0.43
1:A:150:MET:HA	1:A:150:MET:HE2	2.01	0.43
1:A:10:CYS:SG	2:B:142:GLN:HB3	2.58	0.43
1:A:38:HIS:HE1	1:A:40:LYS:HB2	1.82	0.43
1:A:299:ASP:HA	1:A:300:PRO:HD2	1.93	0.42
1:A:247:SER:HB3	1:A:268:VAL:HG23	2.01	0.42
1:A:10:CYS:SG	1:A:13:CYS:HB3	2.60	0.42
1:A:187:ASN:ND2	1:A:202:GLY:HA2	2.36	0.41
2:B:247:LYS:O	2:B:249:PRO:HD3	2.20	0.41
1:A:240:LEU:HA	1:A:243:GLN:HE21	1.86	0.40
1:A:151:LYS:HB2	1:A:194:TYR:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/330 (93%)	273 (89%)	29 (9%)	5 (2%)	7	29
2	B	250/258 (97%)	245 (98%)	5 (2%)	0	100	100
All	All	557/588 (95%)	518 (93%)	34 (6%)	5 (1%)	14	41

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	53	GLU
1	A	190	VAL
1	A	266	GLU
1	A	257	ASP
1	A	31	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/290 (96%)	252 (91%)	25 (9%)	9	30
2	B	229/234 (98%)	221 (96%)	8 (4%)	32	56
All	All	506/524 (97%)	473 (94%)	33 (6%)	15	42

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	24	LYS
1	A	30	PHE
1	A	44	ARG
1	A	48	LYS
1	A	50	VAL
1	A	53	GLU
1	A	55	GLU
1	A	62	THR
1	A	70	LEU
1	A	76	GLU
1	A	96	LYS
1	A	124	SER
1	A	128	ARG
1	A	144	LYS
1	A	172	ARG
1	A	176	LEU
1	A	195	ARG
1	A	235	GLN
1	A	248	ILE
1	A	258	ASN
1	A	264	LYS
1	A	267	LEU
1	A	311	GLU
1	A	328	LEU
2	B	17	GLU
2	B	30	ASP
2	B	35	LEU
2	B	106	ARG
2	B	136	LEU
2	B	149	THR
2	B	164	VAL
2	B	217	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	GLN
1	A	187	ASN
1	A	235	GLN
1	A	237	GLN
1	A	243	GLN
1	A	258	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	43	ASN
2	B	46	GLN
2	B	50	GLN
2	B	56	GLN
2	B	60	ASN
2	B	129	GLN
2	B	152	HIS
2	B	239	HIS
2	B	243	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	A	186	1	8,10,11	0.89	0	10,14,16	1.90	4 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	186	1	-	1/9/11/13	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	186	TPO	O2P-P-OG1	3.05	117.72	105.85
1	A	186	TPO	O-C-CA	-2.47	118.43	124.77
1	A	186	TPO	O3P-P-OG1	2.21	114.46	105.85
1	A	186	TPO	O2P-P-O1P	-2.05	102.84	110.83

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	186	TPO	CB-OG1-P-O3P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TRS	B	301	-	7,7,7	0.33	0	9,9,9	0.44	0
3	Q6E	A	401	-	33,33,33	0.31	0	42,49,49	0.89	3 (7%)
5	PO4	B	302	-	4,4,4	2.06	3 (75%)	6,6,6	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TRS	B	301	-	-	9/9/9/9	-
3	Q6E	A	401	-	-	2/16/36/36	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	302	PO4	P-O1	2.20	1.55	1.50
5	B	302	PO4	P-O4	2.02	1.60	1.54
5	B	302	PO4	P-O2	2.01	1.60	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	Q6E	C10-C9-C11	-2.86	104.15	110.04
3	A	401	Q6E	C9-C10-C5	2.63	113.25	110.82
3	A	401	Q6E	C16-C20-C21	2.35	130.53	126.71

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	301	TRS	C1-C-C2-O2
4	B	301	TRS	C2-C-C1-O1
4	B	301	TRS	N-C-C2-O2
4	B	301	TRS	C1-C-C3-O3
4	B	301	TRS	C2-C-C3-O3
3	A	401	Q6E	C17-C16-C20-C24
3	A	401	Q6E	C17-C16-C20-C21
4	B	301	TRS	C3-C-C1-O1
4	B	301	TRS	C3-C-C2-O2
4	B	301	TRS	N-C-C1-O1
4	B	301	TRS	N-C-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/330 (94%)	-0.32	7 (2%) 62 46	109, 159, 223, 257	0
2	B	252/258 (97%)	-0.58	0 100 100	111, 146, 214, 235	0
All	All	565/588 (96%)	-0.44	7 (1%) 76 62	109, 153, 219, 257	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	228	ILE	3.9
1	A	30	PHE	3.5
1	A	5	TYR	3.1
1	A	328	LEU	2.7
1	A	200	LEU	2.3
1	A	222	MET	2.1
1	A	244	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	A	186	11/12	0.96	0.06	160,166,174,178	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PO4	B	302	5/5	0.74	0.06	220,221,221,222	0
4	TRS	B	301	8/8	0.90	0.14	165,169,171,171	0
3	Q6E	A	401	30/30	0.95	0.09	121,127,141,143	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.