



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 11:19 AM UTC

PDB ID : 7Z43 / pdb_00007z43
Title : Influenza B polymerase with Pol II pSer5 CTD peptide mimic bound in site 1B and 2B
Authors : Cusack, S.; Drncova, P.
Deposited on : 2022-03-03
Resolution : 3.12 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1816 (3.14-3.10)
Clashscore	190562	1906 (3.14-3.10)
Ramachandran outliers	187476	1802 (3.14-3.10)
Sidechain outliers	187428	1802 (3.14-3.10)
RSRZ outliers	180081	1816 (3.14-3.10)
RNA backbone	3983	1017 (3.34-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	751	4% 91% 5%
2	BBB	772	4% 91% 5%
3	CCC	798	4% 86% 5% 8%
4	RRR	18	22% 33% 56% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	VVV	14	43% 29% 29%
6	MaM	13	8% 23% 15% 62%
7	XXX	28	11% 25% 75%
7	YYY	28	4% 57% 39%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 18464 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polymerase acidic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	717	Total	C	N	O	S	0	0	0
			5755	3657	963	1095	40			

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	-13	GLY	-	expression tag	UNP Q5V8Z9
AAA	-12	SER	-	expression tag	UNP Q5V8Z9
AAA	-11	HIS	-	expression tag	UNP Q5V8Z9
AAA	-10	HIS	-	expression tag	UNP Q5V8Z9
AAA	-9	HIS	-	expression tag	UNP Q5V8Z9
AAA	-8	HIS	-	expression tag	UNP Q5V8Z9
AAA	-7	HIS	-	expression tag	UNP Q5V8Z9
AAA	-6	HIS	-	expression tag	UNP Q5V8Z9
AAA	-5	HIS	-	expression tag	UNP Q5V8Z9
AAA	-4	HIS	-	expression tag	UNP Q5V8Z9
AAA	-3	GLY	-	expression tag	UNP Q5V8Z9
AAA	-2	SER	-	expression tag	UNP Q5V8Z9
AAA	-1	GLY	-	expression tag	UNP Q5V8Z9
AAA	0	SER	-	expression tag	UNP Q5V8Z9
AAA	727	GLY	-	expression tag	UNP Q5V8Z9
AAA	728	SER	-	expression tag	UNP Q5V8Z9
AAA	729	GLY	-	expression tag	UNP Q5V8Z9
AAA	730	SER	-	expression tag	UNP Q5V8Z9
AAA	731	GLY	-	expression tag	UNP Q5V8Z9
AAA	732	GLU	-	expression tag	UNP Q5V8Z9
AAA	733	ASN	-	expression tag	UNP Q5V8Z9
AAA	734	LEU	-	expression tag	UNP Q5V8Z9
AAA	735	TYR	-	expression tag	UNP Q5V8Z9
AAA	736	PHE	-	expression tag	UNP Q5V8Z9
AAA	737	GLN	-	expression tag	UNP Q5V8Z9

- Molecule 2 is a protein called RNA-directed RNA polymerase catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	BBB	750	Total	C	N	O	S	0	0	0
			5881	3711	1020	1098	52			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
BBB	-8	GLY	-	expression tag	UNP Q5V8Y6
BBB	-7	SER	-	expression tag	UNP Q5V8Y6
BBB	-6	GLY	-	expression tag	UNP Q5V8Y6
BBB	-5	SER	-	expression tag	UNP Q5V8Y6
BBB	-4	GLY	-	expression tag	UNP Q5V8Y6
BBB	-3	SER	-	expression tag	UNP Q5V8Y6
BBB	-2	GLY	-	expression tag	UNP Q5V8Y6
BBB	-1	SER	-	expression tag	UNP Q5V8Y6
BBB	0	GLY	-	expression tag	UNP Q5V8Y6
BBB	753	GLY	-	expression tag	UNP Q5V8Y6
BBB	754	SER	-	expression tag	UNP Q5V8Y6
BBB	755	GLY	-	expression tag	UNP Q5V8Y6
BBB	756	SER	-	expression tag	UNP Q5V8Y6
BBB	757	GLY	-	expression tag	UNP Q5V8Y6
BBB	758	GLU	-	expression tag	UNP Q5V8Y6
BBB	759	ASN	-	expression tag	UNP Q5V8Y6
BBB	760	LEU	-	expression tag	UNP Q5V8Y6
BBB	761	TYR	-	expression tag	UNP Q5V8Y6
BBB	762	PHE	-	expression tag	UNP Q5V8Y6
BBB	763	GLN	-	expression tag	UNP Q5V8Y6

- Molecule 3 is a protein called Polymerase basic protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	CCC	732	Total	C	N	O	S	0	0	0
			5858	3723	1028	1067	40			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
CCC	-8	GLY	-	expression tag	UNP Q5V8X3
CCC	-7	SER	-	expression tag	UNP Q5V8X3
CCC	-6	GLY	-	expression tag	UNP Q5V8X3
CCC	-5	SER	-	expression tag	UNP Q5V8X3
CCC	-4	GLY	-	expression tag	UNP Q5V8X3
CCC	-3	SER	-	expression tag	UNP Q5V8X3
CCC	-2	GLY	-	expression tag	UNP Q5V8X3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
CCC	-1	SER	-	expression tag	UNP Q5V8X3
CCC	0	GLY	-	expression tag	UNP Q5V8X3
CCC	771	GLY	-	expression tag	UNP Q5V8X3
CCC	772	TRP	-	expression tag	UNP Q5V8X3
CCC	773	SER	-	expression tag	UNP Q5V8X3
CCC	774	HIS	-	expression tag	UNP Q5V8X3
CCC	775	PRO	-	expression tag	UNP Q5V8X3
CCC	776	GLN	-	expression tag	UNP Q5V8X3
CCC	777	PHE	-	expression tag	UNP Q5V8X3
CCC	778	GLU	-	expression tag	UNP Q5V8X3
CCC	779	LYS	-	expression tag	UNP Q5V8X3
CCC	780	GLY	-	expression tag	UNP Q5V8X3
CCC	781	SER	-	expression tag	UNP Q5V8X3
CCC	782	GLY	-	expression tag	UNP Q5V8X3
CCC	783	SER	-	expression tag	UNP Q5V8X3
CCC	784	GLU	-	expression tag	UNP Q5V8X3
CCC	785	ASN	-	expression tag	UNP Q5V8X3
CCC	786	LEU	-	expression tag	UNP Q5V8X3
CCC	787	TYR	-	expression tag	UNP Q5V8X3
CCC	788	PHE	-	expression tag	UNP Q5V8X3
CCC	789	GLN	-	expression tag	UNP Q5V8X3

- Molecule 4 is a RNA chain called RNA (5'-R(*UP*AP*UP*AP*CP*CP*UP*CP*UP*GP*CP*UP*CP*CP*UP*GP*CP*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	RRR	18	Total	C	N	O	P	0	0	0
			349	156	52	124	17			

- Molecule 5 is a RNA chain called RNA (5'-R(P*AP*GP*UP*AP*GP*UP*AP*AP*CP*AP*AP*GP*AP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	VVV	14	Total	C	N	O	P	0	0	0
			307	137	62	94	14			


- Molecule 6 is a RNA chain called RNA (5'-R(P*AP*AP*UP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	MaM	5	Total	C	N	O	P	0	0	0
			89	39	15	30	5			

- Molecule 7 is a protein called SER-TYR-SER-PRO-THR-SEP-PRO.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	XXX	7	Total	C	N	O	P	0	0	0
			55	32	7	15	1			
7	YYY	17	Total	C	N	O	P	0	0	0
			135	81	17	35	2			

- Molecule 8 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	BBB	1	Total	O	P	0	0
			5	4	1		

- Molecule 9 is [(2 {R},3 {S},4 {R},5 {R})-5-(2-azanyl-7-methyl-6-oxidanylidene-1 {H}-purin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methyl phosphono hydrogen phosphate (CCD ID: IC5) (formula: C₁₁H₁₈N₅O₁₁P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	MaM	1	Total	C	N	O	P	0	0
			29	11	5	11	2		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	YYY	1	Total	O	0	0
			1	1		

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	200.82Å 200.82Å 257.02Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	76.97 – 3.12 76.97 – 3.12	Depositor EDS
% Data completeness (in resolution range)	73.1 (76.97-3.12) 73.1 (76.97-3.12)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 3.13Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.240 , 0.272 0.240 , 0.272	Depositor DCC
R_{free} test set	3951 reflections (3.70%)	wwPDB-VP
Wilson B-factor (Å ²)	98.3	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 60.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.007 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	18464	wwPDB-VP
Average B, all atoms (Å ²)	116.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IC5, SEP, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	1.02	0/5871	1.51	3/7918 (0.0%)
2	BBB	1.02	0/5997	1.50	0/8085
3	CCC	1.03	0/5958	1.46	0/8006
4	RRR	0.81	0/386	1.07	5/598 (0.8%)
5	VVV	0.82	0/345	1.02	1/535 (0.2%)
6	MaM	0.85	0/98	1.02	0/150
7	XXX	0.90	0/46	1.00	0/61
7	YYY	0.89	0/120	1.09	0/163
All	All	1.02	0/18821	1.47	9/25516 (0.0%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	VVV	5	G	P-O3'-C3'	-6.66	110.21	120.20
1	AAA	606	GLY	CA-C-O	-6.37	118.08	122.22
4	RRR	17	C	P-O3'-C3'	-5.74	111.59	120.20
4	RRR	8	C	P-O3'-C3'	-5.46	112.02	120.20
4	RRR	3	U	P-O3'-C3'	-5.33	112.21	120.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	5755	0	5730	11	0
2	BBB	5881	0	5904	21	0
3	CCC	5858	0	6031	17	0
4	RRR	349	0	177	3	0
5	VVV	307	0	153	2	0
6	MaM	89	0	43	0	0
7	XXX	55	0	42	0	0
7	YYY	135	0	108	0	0
8	BBB	5	0	0	0	0
9	MaM	29	0	0	0	0
10	YYY	1	0	0	0	0
All	All	18464	0	18188	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 46 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:BBB:340:ALA:HB3	2:BBB:341:PRO:HD3	1.74	0.70
2:BBB:282:LEU:HD22	2:BBB:440:LEU:HD13	1.76	0.67
2:BBB:370:CYS:HA	2:BBB:373:LEU:HD13	1.81	0.63
2:BBB:410:MET:O	4:RRR:16:G:N2	2.33	0.61
1:AAA:305:LEU:HD11	1:AAA:494:ASP:HB3	1.91	0.53

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	713/751 (95%)	659 (92%)	53 (7%)	1 (0%)	48 77
2	BBB	748/772 (97%)	707 (94%)	37 (5%)	4 (0%)	24 55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	CCC	728/798 (91%)	663 (91%)	63 (9%)	2 (0%)	36	65
7	XXX	4/28 (14%)	4 (100%)	0	0	100	100
7	YYY	13/28 (46%)	11 (85%)	1 (8%)	1 (8%)	1	4
All	All	2206/2377 (93%)	2044 (93%)	154 (7%)	8 (0%)	30	60

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	BBB	198	GLY
1	AAA	673	GLY
2	BBB	648	HIS
3	CCC	471	ALA
7	YYY	41	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	638/664 (96%)	624 (98%)	14 (2%)	45	68
2	BBB	643/657 (98%)	629 (98%)	14 (2%)	45	68
3	CCC	639/694 (92%)	620 (97%)	19 (3%)	36	63
7	XXX	6/24 (25%)	6 (100%)	0	100	100
7	YYY	15/24 (62%)	15 (100%)	0	100	100
All	All	1941/2063 (94%)	1894 (98%)	47 (2%)	43	67

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	CCC	34	ARG
3	CCC	191	LYS
3	CCC	61	LEU
3	CCC	89	LYS
3	CCC	271	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	RRR	15/18 (83%)	6 (40%)	1 (6%)
5	VVV	13/14 (92%)	8 (61%)	1 (7%)
6	MaM	3/13 (23%)	2 (66%)	0
All	All	31/45 (68%)	16 (51%)	2 (6%)

5 of 16 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
4	RRR	4	A
4	RRR	5	C
4	RRR	6	C
4	RRR	13	U
4	RRR	14	C

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
4	RRR	4	A
5	VVV	5	G

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	SEP	YYY	35	7	8,9,10	0.62	0	7,12,14	0.62	0
7	SEP	XXX	26	7	8,9,10	0.59	0	7,12,14	0.60	0
7	SEP	YYY	42	7	8,9,10	0.63	0	7,12,14	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	SEP	YYY	35	7	-	1/6/8/10	-
7	SEP	XXX	26	7	-	5/6/8/10	-
7	SEP	YYY	42	7	-	3/6/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	YYY	35	SEP	N-CA-CB-OG
7	XXX	26	SEP	N-CA-CB-OG
7	XXX	26	SEP	C-CA-CB-OG
7	XXX	26	SEP	CB-OG-P-O1P
7	XXX	26	SEP	CB-OG-P-O2P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	IC5	MaM	101	6	28,31,31	1.04	2 (7%)	42,49,49	0.65	0
8	PO4	BBB	801	-	4,4,4	0.70	0	6,6,6	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	IC5	MaM	101	6	-	6/14/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	MaM	101	IC5	PB-O3B	3.72	1.62	1.50
9	MaM	101	IC5	C2'-C1'	-2.66	1.50	1.53

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
9	MaM	101	IC5	C5'-O5'-PA-O1A
9	MaM	101	IC5	PB-O3A-PA-O5'
9	MaM	101	IC5	O4'-C4'-C5'-O5'
9	MaM	101	IC5	C3'-C4'-C5'-O5'
9	MaM	101	IC5	PB-O3A-PA-O1A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	717/751 (95%)	0.66	31 (4%) 40 22	59, 104, 208, 247	0
2	BBB	750/772 (97%)	0.52	30 (4%) 42 23	62, 106, 146, 186	0
3	CCC	732/798 (91%)	0.70	35 (4%) 35 19	71, 111, 187, 223	0
4	RRR	18/18 (100%)	0.70	4 (22%) 2 1	66, 94, 173, 175	0
5	VVV	14/14 (100%)	-0.49	0 100 100	72, 78, 81, 81	0
6	MaM	5/13 (38%)	1.69	1 (20%) 3 1	143, 146, 192, 212	0
7	XXX	6/28 (21%)	1.83	3 (50%) 0 0	137, 139, 150, 152	0
7	YYY	15/28 (53%)	0.88	1 (6%) 24 12	106, 126, 141, 142	0
All	All	2257/2422 (93%)	0.63	105 (4%) 36 19	59, 107, 192, 247	0

The worst 5 of 105 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	BBB	645	THR	6.5
1	AAA	723	ILE	5.2
6	MaM	5	U	5.2
1	AAA	189	VAL	5.1
2	BBB	636	ILE	5.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	SEP	YYY	42	10/11	0.87	0.11	141,149,154,155	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	SEP	YYY	35	10/11	0.88	0.10	121,123,125,128	0
7	SEP	XXX	26	10/11	0.90	0.11	132,136,147,149	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	PO4	BBB	801	5/5	0.86	0.09	136,137,138,141	0
9	IC5	MaM	101	29/29	0.87	0.13	156,181,183,186	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around IC5 MaM 101:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.