



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:30 AM UTC

PDB ID : 2Z56 / pdb_00002z56
Title : Crystal structure of G56S-propeptide:S324A-subtilisin complex
Authors : Pulido, M.A.; Tanaka, S.; Sringiew, C.; You, D.J.; Matsumura, H.; Koga, Y.;
Takano, K.; Kanaya, S.
Deposited on : 2007-06-29
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	318	17% 71% 25% .
2	B	65	20% 72% 25% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2978 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tk-subtilisin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	318	Total	C	N	O	S	0	0	0
			2314	1452	384	472	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	324	ALA	SER	engineered mutation	UNP P58502

- Molecule 2 is a protein called Tk-subtilisin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	65	Total	C	N	O	S	0	0	0
			502	330	85	86	1			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	56	SER	GLY	engineered mutation	UNP P58502

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	7	Total	Ca	0	0
			7	7		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	128	Total 128	O 128	0	0
5	B	26	Total 26	O 26	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tk-subtilisin



• Molecule 2: Tk-subtilisin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.52Å 67.46Å 74.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.90 49.87 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-1.90) 99.7 (49.87-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.189 , 0.209 (Not available) , 0.206	Depositor DCC
R_{free} test set	1310 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	26.3	Xtriage
Anisotropy	0.186	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 36.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2978	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.92	49/2361 (2.1%)	1.46	17/3235 (0.5%)
2	B	1.64	3/512 (0.6%)	1.23	1/693 (0.1%)
All	All	1.87	52/2873 (1.8%)	1.42	18/3928 (0.5%)

All (52) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	233	MET	SD-CE	-18.66	1.32	1.79
1	A	210	VAL	C-O	10.74	1.35	1.24
1	A	247	MET	SD-CE	-9.01	1.57	1.79
2	B	36	ILE	CA-C	8.59	1.59	1.52
1	A	327	THR	C-O	-8.44	1.16	1.24
1	A	236	GLY	C-O	7.84	1.32	1.24
1	A	196	ILE	CG1-CD1	-7.58	1.22	1.51
1	A	365	ILE	C-O	7.31	1.32	1.24
1	A	308	ILE	C-O	-7.07	1.16	1.24
2	B	58	GLU	CA-C	6.91	1.62	1.52
1	A	171	VAL	CA-CB	6.81	1.62	1.55
1	A	180	TYR	C-O	6.67	1.31	1.24
1	A	316	SER	CA-C	6.31	1.59	1.53
1	A	197	ALA	CA-CB	-6.26	1.43	1.53
1	A	353	THR	CA-C	-6.25	1.45	1.52
1	A	388	ARG	CD-NE	-6.16	1.37	1.46
1	A	101	ILE	CA-CB	-6.08	1.47	1.54
1	A	87	PRO	CA-C	6.08	1.60	1.52
1	A	228	ALA	CA-C	6.03	1.60	1.52
1	A	327	THR	CA-CB	5.99	1.60	1.53
1	A	372	ASP	CA-C	-5.93	1.45	1.52
1	A	269	SER	CA-C	5.80	1.57	1.52
1	A	86	ILE	CA-CB	-5.77	1.47	1.53
1	A	226	ASP	CG-OD2	5.76	1.36	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	280	ILE	N-CA	-5.72	1.39	1.46
1	A	315	ASP	N-CA	-5.70	1.38	1.46
1	A	230	VAL	C-O	5.69	1.29	1.24
1	A	156	HIS	C-O	5.62	1.30	1.24
1	A	210	VAL	CA-CB	5.61	1.61	1.54
1	A	335	ALA	N-CA	-5.61	1.39	1.46
1	A	340	ALA	C-O	5.60	1.30	1.24
1	A	157	VAL	CA-CB	-5.59	1.47	1.54
1	A	165	ASN	C-N	-5.56	1.28	1.33
1	A	158	ILE	N-CA	-5.44	1.40	1.46
1	A	183	ARG	CA-CB	5.44	1.59	1.52
1	A	326	ALA	CA-CB	5.42	1.61	1.53
1	A	119	ASP	N-CA	5.42	1.53	1.46
1	A	390	ALA	C-O	-5.38	1.17	1.24
1	A	131	TRP	NE1-CE2	5.38	1.43	1.37
1	A	345	TYR	C-N	-5.37	1.25	1.33
1	A	159	GLY	C-O	-5.32	1.17	1.23
1	A	96	PRO	C-O	-5.31	1.17	1.24
1	A	242	SER	C-O	-5.28	1.17	1.24
1	A	321	MET	SD-CE	-5.24	1.66	1.79
1	A	275	ALA	CA-C	-5.23	1.45	1.52
1	A	111	VAL	C-O	5.16	1.29	1.24
1	A	387	VAL	CA-C	5.12	1.59	1.52
2	B	36	ILE	CA-CB	-5.09	1.49	1.55
1	A	329	HIS	N-CA	-5.07	1.40	1.46
1	A	196	ILE	CA-C	-5.06	1.46	1.52
1	A	157	VAL	C-O	5.05	1.29	1.24
1	A	249	ILE	CA-C	-5.04	1.45	1.52

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	SER	N-CA-C	10.20	123.67	111.33
1	A	345	TYR	N-CA-C	-7.09	98.38	109.72
1	A	242	SER	N-CA-C	6.44	119.11	111.71
1	A	297	ARG	N-CA-C	5.95	117.73	110.41
1	A	294	PHE	N-CA-C	5.54	119.89	113.18
1	A	326	ALA	CA-C-N	5.43	129.00	120.97
1	A	326	ALA	C-N-CA	5.43	129.00	120.97
1	A	198	ILE	CA-C-N	5.36	125.89	119.94
1	A	198	ILE	C-N-CA	5.36	125.89	119.94
1	A	196	ILE	CA-CB-CG2	5.24	119.41	110.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	196	ILE	CB-CG1-CD1	-5.23	102.82	113.80
1	A	167	ASP	N-CA-C	5.19	119.59	113.16
1	A	282	VAL	N-CA-C	5.18	115.31	107.75
1	A	226	ASP	CB-CG-OD2	5.14	130.23	118.40
1	A	354	PHE	N-CA-C	5.14	116.88	111.28
2	B	30	VAL	N-CA-C	-5.08	106.85	111.67
1	A	388	ARG	NE-CZ-NH2	-5.06	114.65	119.20
1	A	303	ALA	N-CA-C	5.00	112.72	108.22

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2314	0	2239	45	1
2	B	502	0	536	23	0
3	A	7	0	0	1	0
4	B	1	0	0	0	0
5	A	128	0	0	12	1
5	B	26	0	0	6	1
All	All	2978	0	2775	64	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:147:CYS:HB3	5:A:1134:HOH:O	0.92	1.08
2:B:13:ASP:CG	2:B:16:LYS:NZ	2.15	1.05
2:B:13:ASP:CG	2:B:16:LYS:HZ3	1.66	1.02
2:B:5:THR:N	5:B:2009:HOH:O	1.92	1.01
1:A:314:ASP:OD2	5:A:1064:HOH:O	1.84	0.96
1:A:122:HIS:HD2	1:A:124:ASP:H	1.28	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:241:ASP:OD2	2:B:64:HIS:HD2	1.69	0.76
2:B:68:LEU:HD12	2:B:68:LEU:O	1.87	0.73
1:A:307:ASP:OD2	5:A:1037:HOH:O	2.06	0.73
1:A:357:ILE:HB	1:A:368:ILE:HD11	1.72	0.72
1:A:338:GLN:HE22	1:A:349:LEU:H	1.37	0.71
2:B:13:ASP:CG	2:B:16:LYS:HZ1	1.93	0.71
1:A:371:ASP:OD1	5:A:1117:HOH:O	2.08	0.71
2:B:10:VAL:HG13	2:B:57:VAL:HG13	1.74	0.67
2:B:13:ASP:OD2	2:B:16:LYS:NZ	2.27	0.67
1:A:356:ASP:O	1:A:367:HIS:HD2	1.78	0.67
2:B:13:ASP:CB	2:B:16:LYS:HZ1	2.09	0.65
2:B:10:VAL:CG1	2:B:57:VAL:HG13	2.30	0.61
1:A:300:GLU:OE2	1:A:367:HIS:HE1	1.83	0.61
1:A:226:ASP:OD2	3:A:1002:CA:CA	1.78	0.60
1:A:147:CYS:CB	5:A:1134:HOH:O	1.78	0.59
1:A:338:GLN:HA	1:A:338:GLN:HE21	1.68	0.59
1:A:318:GLU:OE2	5:A:1013:HOH:O	2.18	0.57
1:A:321:MET:HE2	5:A:1067:HOH:O	2.04	0.56
1:A:122:HIS:HE1	1:A:316:SER:O	1.88	0.56
1:A:267:ALA:H	1:A:296:ASN:ND2	2.05	0.55
1:A:208:ASP:OD2	1:A:213:LYS:HD2	2.07	0.53
1:A:144:LEU:HA	1:A:147:CYS:SG	2.48	0.53
1:A:324:ALA:HB2	2:B:69:LEU:C	2.34	0.52
2:B:55:PRO:HB3	5:B:2025:HOH:O	2.10	0.52
1:A:137:ARG:NH2	5:A:1099:HOH:O	2.43	0.51
1:A:342:TYR:O	1:A:345:TYR:O	2.28	0.51
2:B:13:ASP:CB	2:B:16:LYS:NZ	2.70	0.51
1:A:213:LYS:HD2	5:A:1119:HOH:O	2.10	0.51
1:A:269:SER:HB2	1:A:270:PRO:CD	2.42	0.50
1:A:143:LYS:NZ	5:A:1087:HOH:O	2.44	0.49
1:A:369:THR:O	1:A:388:ARG:HD3	2.13	0.49
1:A:327:THR:N	1:A:328:PRO:HD2	2.28	0.48
2:B:5:THR:CA	5:B:2009:HOH:O	2.54	0.48
1:A:357:ILE:CB	1:A:368:ILE:HD11	2.44	0.48
1:A:213:LYS:CE	5:A:1041:HOH:O	2.62	0.47
2:B:5:THR:HA	5:B:2009:HOH:O	2.16	0.46
1:A:241:ASP:OD2	2:B:64:HIS:CD2	2.59	0.46
2:B:20:HIS:CE1	5:B:2027:HOH:O	2.69	0.46
1:A:95:ALA:N	1:A:96:PRO:CD	2.79	0.46
1:A:344:LYS:HD3	1:A:345:TYR:CZ	2.51	0.45
2:B:52:LYS:NZ	5:B:2016:HOH:O	2.48	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:ILE:C	1:A:196:ILE:HD12	2.41	0.45
2:B:68:LEU:HD12	2:B:68:LEU:C	2.42	0.45
1:A:192:SER:O	1:A:196:ILE:HG23	2.17	0.45
1:A:371:ASP:N	1:A:388:ARG:HD2	2.31	0.45
2:B:6:ILE:HD11	2:B:48:VAL:CG2	2.48	0.44
1:A:338:GLN:NE2	1:A:349:LEU:H	2.12	0.44
1:A:248:ILE:HD13	1:A:276:TYR:CD2	2.52	0.43
2:B:23:LEU:C	2:B:23:LEU:HD13	2.43	0.43
2:B:68:LEU:C	2:B:68:LEU:CD1	2.92	0.43
1:A:156:HIS:CD2	1:A:310:SER:HB3	2.53	0.43
1:A:324:ALA:CB	2:B:69:LEU:C	2.92	0.43
1:A:224:ASP:HA	5:A:1133:HOH:O	2.19	0.42
1:A:371:ASP:HB2	1:A:388:ARG:HD2	2.01	0.42
1:A:156:HIS:HA	1:A:311:THR:O	2.19	0.42
1:A:337:ILE:HD11	1:A:392:ALA:HB1	2.02	0.42
1:A:294:PHE:CD1	1:A:294:PHE:C	2.98	0.41
1:A:300:GLU:OE2	1:A:367:HIS:CE1	2.68	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1034:HOH:O	5:A:1057:HOH:O[2_554]	2.12	0.08
1:A:265:GLU:OE2	5:B:2017:HOH:O[2_565]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/318 (99%)	302 (96%)	14 (4%)	0	100	100
2	B	63/65 (97%)	63 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	379/383 (99%)	365 (96%)	14 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	242/242 (100%)	240 (99%)	2 (1%)	73	75
2	B	56/56 (100%)	53 (95%)	3 (5%)	20	12
All	All	298/298 (100%)	293 (98%)	5 (2%)	53	52

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	196	ILE
1	A	338	GLN
2	B	5	THR
2	B	65	GLN
2	B	68	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	122	HIS
1	A	250	GLN
1	A	290	ASN
1	A	296	ASN
1	A	338	GLN
1	A	343	GLN
1	A	367	HIS
2	B	46	ASN
2	B	64	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	318/318 (100%)	1.04	53 (16%) 4 4	20, 27, 39, 56	0
2	B	65/65 (100%)	1.32	13 (20%) 3 2	25, 35, 48, 60	0
All	All	383/383 (100%)	1.08	66 (17%) 4 4	20, 28, 43, 60	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	398	GLY	7.1
1	A	99	TRP	6.1
1	A	174	ALA	5.8
1	A	153	HIS	5.7
1	A	132	CYS	5.3
1	A	140	VAL	5.2
2	B	48	VAL	5.1
1	A	387	VAL	4.6
1	A	354	PHE	4.5
2	B	55	PRO	4.4
1	A	144	LEU	4.0
1	A	370	ALA	3.9
1	A	210	VAL	3.7
1	A	298	GLN	3.7
2	B	17	PHE	3.5
1	A	313	PRO	3.4
1	A	306	VAL	3.3
2	B	68	LEU	3.3
1	A	248	ILE	3.2
2	B	38	ALA	3.2
1	A	104	GLY	3.2
1	A	167	ASP	3.0
1	A	245	TYR	3.0
1	A	155	THR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	218	ILE	3.0
1	A	246	ASP	2.9
1	A	100	SER	2.8
1	A	217	GLY	2.8
1	A	265	GLU	2.8
1	A	101	ILE	2.7
1	A	180	TYR	2.7
1	A	294	PHE	2.7
1	A	368	ILE	2.7
1	A	173	VAL	2.7
1	A	282	VAL	2.7
1	A	177	VAL	2.5
1	A	272	TYR	2.5
1	A	384	TYR	2.5
1	A	394	GLN	2.4
2	B	69	LEU	2.4
2	B	46	ASN	2.4
2	B	65	GLN	2.4
2	B	56	SER	2.4
2	B	53	LYS	2.4
1	A	102	THR	2.4
2	B	66	ALA	2.4
2	B	23	LEU	2.3
2	B	45	ALA	2.3
1	A	150	GLN	2.3
1	A	178	GLN	2.3
1	A	366	LEU	2.3
1	A	280	ILE	2.2
1	A	135	THR	2.2
1	A	355	ASP	2.2
1	A	396	ALA	2.2
1	A	249	ILE	2.2
1	A	136	LEU	2.2
1	A	238	PRO	2.2
1	A	143	LYS	2.2
1	A	125	LEU	2.1
1	A	118	VAL	2.1
1	A	147	CYS	2.1
1	A	139	LYS	2.0
1	A	252	TYR	2.0
1	A	145	ARG	2.0
1	A	390	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	A	1007	1/1	0.91	0.15	21,21,21,21	0
3	CA	A	1006	1/1	0.97	0.05	28,28,28,28	0
4	ZN	B	2001	1/1	0.98	0.04	37,37,37,37	0
3	CA	A	1001	1/1	0.99	0.02	24,24,24,24	0
3	CA	A	1003	1/1	0.99	0.05	22,22,22,22	0
3	CA	A	1005	1/1	0.99	0.03	22,22,22,22	0
3	CA	A	1002	1/1	1.00	0.10	14,14,14,14	0
3	CA	A	1004	1/1	1.00	0.11	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.