



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 04:37 PM UTC

PDB ID : 1Z6R / pdb_00001z6r
Title : Crystal structure of Mlc from Escherichia coli
Authors : Schiefner, A.; Gerber, K.; Seitz, S.; Welte, W.; Diederichs, K.; Boos, W.
Deposited on : 2005-03-23
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	 4% 61% 25% 7% 6%
1	B	406	 % 59% 28% 7% 6%
1	C	406	 3% 54% 32% 7% 6%
1	D	406	 6% 60% 30% 5% 6%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 11728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mlc protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	382	Total	C	N	O	S	Se	0	0	0
			2931	1855	514	549	4	9			
1	B	382	Total	C	N	O	S	Se	0	0	0
			2931	1855	514	549	4	9			
1	C	382	Total	C	N	O	S	Se	0	0	0
			2931	1855	514	549	4	9			
1	D	382	Total	C	N	O	S	Se	0	0	0
			2931	1855	514	549	4	9			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP P50456
A	52	HIS	ARG	engineered mutation	UNP P50456
A	54	MSE	MET	modified residue	UNP P50456
A	167	MSE	MET	modified residue	UNP P50456
A	176	MSE	MET	modified residue	UNP P50456
A	201	MSE	MET	modified residue	UNP P50456
A	286	MSE	MET	modified residue	UNP P50456
A	289	MSE	MET	modified residue	UNP P50456
A	329	MSE	MET	modified residue	UNP P50456
A	384	MSE	MET	modified residue	UNP P50456
A	394	MSE	MET	modified residue	UNP P50456
B	1	MSE	MET	modified residue	UNP P50456
B	52	HIS	ARG	engineered mutation	UNP P50456
B	54	MSE	MET	modified residue	UNP P50456
B	167	MSE	MET	modified residue	UNP P50456
B	176	MSE	MET	modified residue	UNP P50456
B	201	MSE	MET	modified residue	UNP P50456
B	286	MSE	MET	modified residue	UNP P50456
B	289	MSE	MET	modified residue	UNP P50456
B	329	MSE	MET	modified residue	UNP P50456
B	384	MSE	MET	modified residue	UNP P50456

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	394	MSE	MET	modified residue	UNP P50456
C	1	MSE	MET	modified residue	UNP P50456
C	52	HIS	ARG	engineered mutation	UNP P50456
C	54	MSE	MET	modified residue	UNP P50456
C	167	MSE	MET	modified residue	UNP P50456
C	176	MSE	MET	modified residue	UNP P50456
C	201	MSE	MET	modified residue	UNP P50456
C	286	MSE	MET	modified residue	UNP P50456
C	289	MSE	MET	modified residue	UNP P50456
C	329	MSE	MET	modified residue	UNP P50456
C	384	MSE	MET	modified residue	UNP P50456
C	394	MSE	MET	modified residue	UNP P50456
D	1	MSE	MET	modified residue	UNP P50456
D	52	HIS	ARG	engineered mutation	UNP P50456
D	54	MSE	MET	modified residue	UNP P50456
D	167	MSE	MET	modified residue	UNP P50456
D	176	MSE	MET	modified residue	UNP P50456
D	201	MSE	MET	modified residue	UNP P50456
D	286	MSE	MET	modified residue	UNP P50456
D	289	MSE	MET	modified residue	UNP P50456
D	329	MSE	MET	modified residue	UNP P50456
D	384	MSE	MET	modified residue	UNP P50456
D	394	MSE	MET	modified residue	UNP P50456

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0



• Molecule 1: Mlc protein



• Molecule 1: Mlc protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	235.95Å 74.71Å 154.95Å 90.00° 129.15° 90.00°	Depositor
Resolution (Å)	19.94 – 2.70 19.94 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.94-2.70) 98.7 (19.94-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 2.71Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.203 , 0.263 0.201 , 0.258	Depositor DCC
R_{free} test set	2856 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	67.0	Xtriage
Anisotropy	0.084	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 77.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.025 for -h-2*1,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11728	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	6/2969 (0.2%)	1.02	7/4010 (0.2%)
1	B	0.94	3/2969 (0.1%)	1.08	9/4010 (0.2%)
1	C	1.09	14/2969 (0.5%)	1.02	7/4010 (0.2%)
1	D	1.05	13/2969 (0.4%)	0.98	5/4010 (0.1%)
All	All	0.99	36/11876 (0.3%)	1.02	28/16040 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	3

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	139	HIS	CE1-NE2	22.08	1.54	1.32
1	C	38	ARG	CZ-NH1	21.54	1.62	1.32
1	C	285	SER	CB-OG	16.96	1.76	1.42
1	D	133	ASP	CG-OD1	14.96	1.53	1.25
1	D	139	HIS	CG-ND1	13.96	1.53	1.38

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	44	PRO	O-C-N	9.19	135.05	122.64
1	B	88	LEU	CA-C-N	-7.31	112.91	123.00
1	B	88	LEU	C-N-CA	-7.31	112.91	123.00
1	A	287	SER	N-CA-C	-6.96	105.72	114.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	D	153	LEU	N-CA-C	6.33	113.82	108.13

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	286	MSE	Mainchain
1	C	38	ARG	Sidechain
1	C	44	PRO	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2931	0	2988	76	0
1	B	2931	0	2988	93	0
1	C	2931	0	2988	93	0
1	D	2931	0	2988	78	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
All	All	11728	0	11952	326	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 326 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:38:ARG:CG	1:C:38:ARG:CD	1.76	1.57
1:B:384:MSE:CE	1:B:384:MSE:SE	2.16	1.44
1:D:289:MSE:SE	1:D:289:MSE:CE	2.15	1.43
1:C:285:SER:OG	1:C:285:SER:CB	1.76	1.33
1:D:91:ARG:CZ	1:D:380:ASN:HD21	1.59	1.13

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	378/406 (93%)	346 (92%)	25 (7%)	7 (2%)	6	17
1	B	378/406 (93%)	353 (93%)	25 (7%)	0	100	100
1	C	378/406 (93%)	344 (91%)	31 (8%)	3 (1%)	16	37
1	D	378/406 (93%)	351 (93%)	24 (6%)	3 (1%)	16	37
All	All	1512/1624 (93%)	1394 (92%)	105 (7%)	13 (1%)	14	35

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	290	LEU
1	C	106	SER
1	D	369	GLN
1	A	286	MSE
1	C	294	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/329 (98%)	271 (84%)	50 (16%)	2	7
1	B	321/329 (98%)	268 (84%)	53 (16%)	2	6
1	C	321/329 (98%)	262 (82%)	59 (18%)	1	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	321/329 (98%)	272 (85%)	49 (15%)	3	7
All	All	1284/1316 (98%)	1073 (84%)	211 (16%)	2	6

5 of 211 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	101	LEU
1	C	300	LEU
1	D	325	ILE
1	C	108	LEU
1	C	188	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	15	GLN
1	D	361	GLN
1	D	41	GLN
1	D	185	HIS
1	D	380	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	373/406 (91%)	0.50	17 (4%) 37 34	57, 71, 78, 87	0
1	B	373/406 (91%)	0.10	5 (1%) 75 73	61, 71, 78, 87	0
1	C	373/406 (91%)	0.45	11 (2%) 53 50	61, 71, 77, 84	0
1	D	373/406 (91%)	0.61	26 (6%) 22 19	60, 71, 77, 86	0
All	All	1492/1624 (91%)	0.41	59 (3%) 42 38	57, 71, 78, 87	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	77	GLY	6.0
1	D	381	GLN	5.9
1	D	382	GLY	5.5
1	D	194	HIS	4.9
1	A	77	GLY	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	A	501	1/1	0.99	0.16	55,55,55,55	0
2	ZN	B	502	1/1	0.99	0.16	46,46,46,46	0
2	ZN	C	503	1/1	0.99	0.12	71,71,71,71	0
2	ZN	D	504	1/1	0.99	0.17	86,86,86,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.