



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:58 PM UTC

PDB ID : 4Z63 / pdb_00004z63
Title : The plant peptide hormone receptor in arabidopsis
Authors : Chai, J.; Wang, J.; Han, Z.
Deposited on : 2015-04-03
Resolution : 2.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	631	 2% 76% 19% . .
2	P	5	 40% 40% 20%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5075 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phytosulfokine receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	610	Total	C	N	O	S	0	0	0
			4748	3003	829	897	19			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	649	HIS	-	expression tag	UNP Q9ZVR7
A	650	HIS	-	expression tag	UNP Q9ZVR7
A	651	HIS	-	expression tag	UNP Q9ZVR7
A	652	HIS	-	expression tag	UNP Q9ZVR7
A	653	HIS	-	expression tag	UNP Q9ZVR7
A	654	HIS	-	expression tag	UNP Q9ZVR7

- Molecule 2 is a protein called Phytosulfokine.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	5	Total	C	N	O	S	0	0	0
			57	33	6	16	2			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			13	8	1	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	143	Total	O	0	0
			143	143		
4	P	2	Total	O	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phytosulfokine receptor 1



• Molecule 2: Phytosulfokine



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	92.89Å 92.89Å 242.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.54 – 2.51 33.54 – 2.51	Depositor EDS
% Data completeness (in resolution range)	98.0 (33.54-2.51) 97.8 (33.54-2.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.62 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
R, R_{free}	0.229 , 0.276 (Not available) , 0.244	Depositor DCC
R_{free} test set	1821 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	37.2	Xtriage
Anisotropy	0.691	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 22.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5075	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TYS, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	6/4850 (0.1%)	1.04	21/6572 (0.3%)
2	P	0.55	0/23	1.13	0/27
All	All	0.65	6/4873 (0.1%)	1.04	21/6599 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	492	ARG	CA-C	10.79	1.68	1.52
1	A	492	ARG	N-CA	6.61	1.54	1.46
1	A	515	ALA	C-O	-5.88	1.15	1.23
1	A	615	GLY	C-O	-5.83	1.16	1.23
1	A	358	GLU	C-O	-5.81	1.16	1.24
1	A	359	SER	C-O	-5.35	1.17	1.24

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	491	SER	O-C-N	-11.35	107.13	122.23
1	A	492	ARG	N-CA-C	10.85	126.26	112.92
1	A	357	PRO	CA-C-O	7.23	129.53	121.43
1	A	188	PHE	N-CA-C	-7.03	99.26	109.59
1	A	327	ARG	N-CA-C	6.29	119.89	112.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	492	ARG	CA-C-N	6.11	133.50	121.58
1	A	492	ARG	C-N-CA	6.11	133.50	121.58
1	A	455	ILE	CA-C-N	6.06	126.03	120.03
1	A	455	ILE	C-N-CA	6.06	126.03	120.03
1	A	613	SER	N-CA-C	5.87	118.47	110.55
1	A	371	LEU	N-CA-C	5.72	120.25	112.88
1	A	491	SER	CA-C-N	-5.47	113.90	122.65
1	A	491	SER	C-N-CA	-5.47	113.90	122.65
1	A	351	THR	N-CA-C	5.46	118.94	112.72
1	A	374	SER	N-CA-C	5.45	120.04	113.17
1	A	358	GLU	CB-CG-CD	5.38	121.75	112.60
1	A	75	THR	N-CA-C	5.37	119.94	113.17
1	A	225	LEU	CA-C-N	-5.37	117.51	122.29
1	A	225	LEU	C-N-CA	-5.37	117.51	122.29
1	A	114	ASP	CB-CA-C	-5.30	108.89	115.89
1	A	56	SER	N-CA-C	5.09	117.49	111.33

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	491	SER	Peptide,Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4748	0	4710	61	1
2	P	57	0	44	1	0
3	A	125	0	115	5	0
4	A	143	0	0	7	0
4	P	2	0	0	0	0
All	All	5075	0	4869	63	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:484:THR:HG21	1:A:542:GLU:HB3	1.42	0.99
1:A:484:THR:HG21	1:A:542:GLU:CB	2.02	0.88
1:A:140:ILE:HG13	1:A:164:PRO:HG3	1.66	0.77
1:A:331:ARG:NH1	4:A:803:HOH:O	2.17	0.77
1:A:484:THR:HG23	1:A:542:GLU:O	1.87	0.74
1:A:484:THR:CG2	1:A:542:GLU:HB3	2.19	0.72
1:A:567:LEU:HD22	1:A:570:MET:HE2	1.73	0.70
1:A:371:LEU:HD13	1:A:376:LEU:HD11	1.75	0.68
1:A:69:THR:HB	1:A:80:ARG:HB2	1.79	0.65
3:A:705:NAG:O3	4:A:802:HOH:O	2.16	0.62
1:A:488:SER:HA	1:A:494:ILE:HG21	1.80	0.62
1:A:372:SER:HB2	2:P:31:THR:HG22	1.85	0.58
1:A:241:ARG:HB3	1:A:263:VAL:HB	1.84	0.57
1:A:30:ARG:HG2	1:A:60:THR:HG21	1.86	0.56
1:A:446:LEU:O	1:A:449:ASN:ND2	2.32	0.56
1:A:490:THR:OG1	1:A:491:SER:N	2.38	0.56
1:A:319:ASN:ND2	4:A:805:HOH:O	2.25	0.55
1:A:406:LEU:HD21	1:A:432:PRO:HD3	1.88	0.55
1:A:53:TRP:O	1:A:56:SER:HB3	2.08	0.54
1:A:109:ARG:NH1	4:A:813:HOH:O	2.33	0.54
1:A:406:LEU:HD12	1:A:407:PRO:HD2	1.90	0.54
1:A:191:GLY:H	1:A:215:ASP:CG	2.16	0.53
1:A:592:GLN:HG3	1:A:617:PHE:HD1	1.74	0.52
1:A:592:GLN:NE2	1:A:614:GLY:O	2.43	0.52
3:A:705:NAG:O3	4:A:801:HOH:O	2.10	0.52
1:A:379:ILE:HB	1:A:401:PHE:CG	2.46	0.51
1:A:484:THR:HG21	1:A:542:GLU:HB2	1.89	0.51
1:A:616:GLN:NE2	4:A:822:HOH:O	2.44	0.50
1:A:486:LEU:O	1:A:490:THR:HG23	2.12	0.49
1:A:289:ALA:HB1	1:A:310:LEU:HA	1.94	0.49
1:A:149:LEU:HD21	1:A:152:PHE:HB2	1.95	0.48
1:A:238:ARG:NH2	1:A:262:ASP:H	2.11	0.48
1:A:346:ASN:OD1	1:A:370:SER:OG	2.31	0.47
1:A:30:ARG:HG2	1:A:60:THR:CG2	2.45	0.47
1:A:240:ILE:HG22	1:A:263:VAL:HG21	1.97	0.47
1:A:459:ILE:HD12	1:A:462:PHE:HE2	1.81	0.46
1:A:398:THR:HG23	1:A:423:ALA:O	2.15	0.46
1:A:55:ASN:HB3	1:A:59:SER:HB2	1.98	0.46
1:A:526:PRO:HD2	1:A:550:HIS:CD2	2.50	0.46
1:A:417:LEU:HD21	1:A:420:LEU:HB2	1.99	0.45
1:A:112:ILE:HG22	1:A:136:LEU:HD21	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:531:GLY:HA2	1:A:555:LYS:O	2.17	0.45
1:A:239:GLU:OE1	3:A:709:NAG:H81	2.17	0.45
1:A:331:ARG:HE	1:A:331:ARG:HB3	1.59	0.44
1:A:498:GLU:HB2	1:A:522:PHE:CE2	2.51	0.44
1:A:242:ASN:ND2	3:A:709:NAG:C3	2.80	0.44
1:A:410:SER:HA	1:A:434:TRP:HB3	1.99	0.44
1:A:312:CYS:HB3	1:A:338:ASP:C	2.43	0.43
1:A:431:MET:HG2	1:A:475:PHE:CE1	2.53	0.43
1:A:80:ARG:CG	1:A:104:VAL:HB	2.48	0.43
1:A:117:PRO:HG2	1:A:120:ILE:HD12	1.99	0.43
1:A:220:LYS:HG3	3:A:709:NAG:O4	2.19	0.43
1:A:163:LEU:HA	1:A:164:PRO:HD3	1.83	0.43
1:A:459:ILE:HD12	1:A:462:PHE:CE2	2.54	0.43
1:A:265:ASP:OD1	4:A:804:HOH:O	2.22	0.42
1:A:80:ARG:HG3	1:A:104:VAL:HB	2.02	0.42
1:A:369:PHE:CZ	1:A:371:LEU:HD12	2.54	0.42
1:A:228:GLN:OE1	1:A:228:GLN:N	2.51	0.41
1:A:315:MET:HE2	1:A:315:MET:HB3	1.98	0.41
1:A:454:ALA:HA	1:A:476:THR:O	2.20	0.41
1:A:83:LEU:HD23	1:A:83:LEU:HA	1.94	0.41
1:A:397:LEU:HD23	1:A:397:LEU:HA	1.89	0.41
1:A:481:LYS:O	1:A:484:THR:HB	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:ASP:OD1	1:A:511:GLU:OE1[3_564]	2.06	0.14

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	608/631 (96%)	592 (97%)	15 (2%)	1 (0%)	43	62
2	P	2/5 (40%)	2 (100%)	0	0	100	100
All	All	610/636 (96%)	594 (97%)	15 (2%)	1 (0%)	43	62

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	615	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	548/567 (97%)	511 (93%)	37 (7%)	14	28
2	P	3/3 (100%)	2 (67%)	1 (33%)	0	0
All	All	551/570 (97%)	513 (93%)	38 (7%)	14	28

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	ASP
1	A	56	SER
1	A	94	GLU
1	A	96	LEU
1	A	109	ARG
1	A	118	LEU
1	A	140	ILE
1	A	162	SER
1	A	173	GLN
1	A	189	THR
1	A	207	ASP
1	A	240	ILE
1	A	265	ASP
1	A	266	GLU
1	A	284	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	291	SER
1	A	316	ILE
1	A	336	LEU
1	A	364	GLU
1	A	370	SER
1	A	375	SER
1	A	381	SER
1	A	440	GLU
1	A	468	LEU
1	A	485	LYS
1	A	492	ARG
1	A	507	MET
1	A	529	GLU
1	A	562	SER
1	A	565	SER
1	A	571	THR
1	A	582	ARG
1	A	591	LEU
1	A	599	LYS
1	A	610	VAL
1	A	627	SER
1	A	633	GLU
2	P	31	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	150	GLN
1	A	242	ASN
1	A	355	GLN
1	A	413	HIS
1	A	618	GLN
1	A	634	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TYS	P	28	2	15,16,17	2.08	2 (13%)	15,22,24	0.48	0
2	TYS	P	30	2	15,16,17	1.98	3 (20%)	15,22,24	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TYS	P	28	2	-	0/10/11/13	0/1/1/1
2	TYS	P	30	2	-	0/10/11/13	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	P	28	TYS	OH-CZ	-7.40	1.30	1.42
2	P	30	TYS	OH-CZ	-6.85	1.31	1.42
2	P	28	TYS	O2-S	2.18	1.54	1.45
2	P	30	TYS	O2-S	2.10	1.54	1.45
2	P	30	TYS	OH-S	2.08	1.62	1.58

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	702	1	14,14,15	0.59	0	17,19,21	1.23	2 (11%)
3	NAG	A	708	1	14,14,15	0.60	0	17,19,21	1.29	2 (11%)
3	NAG	A	707	1	14,14,15	0.55	0	17,19,21	0.81	0
3	NAG	A	703	1	14,14,15	0.48	0	17,19,21	0.95	0
3	NAG	A	706	1	14,14,15	0.65	0	17,19,21	1.41	2 (11%)
3	NAG	A	701	1	14,14,15	0.61	0	17,19,21	1.55	2 (11%)
3	NAG	A	709	-	13,13,15	0.94	1 (7%)	15,17,21	2.57	6 (40%)
3	NAG	A	705	1	14,14,15	0.47	0	17,19,21	0.98	1 (5%)
3	NAG	A	704	1	14,14,15	0.49	0	17,19,21	1.57	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	702	1	-	2/6/23/26	0/1/1/1
3	NAG	A	708	1	-	2/6/23/26	0/1/1/1
3	NAG	A	707	1	-	2/6/23/26	0/1/1/1
3	NAG	A	703	1	-	0/6/23/26	0/1/1/1
3	NAG	A	706	1	-	0/6/23/26	0/1/1/1
3	NAG	A	701	1	-	0/6/23/26	0/1/1/1
3	NAG	A	709	-	-	2/6/19/26	0/1/1/1
3	NAG	A	705	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	704	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	709	NAG	C3-C2	-2.37	1.49	1.53

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	709	NAG	C4-C3-C2	-5.88	105.26	112.50
3	A	701	NAG	C1-O5-C5	4.96	118.83	112.19
3	A	709	NAG	C3-C4-C5	-4.44	105.10	110.76
3	A	709	NAG	C3-C2-N2	-4.03	104.00	110.62
3	A	706	NAG	C1-O5-C5	3.70	117.14	112.19
3	A	708	NAG	C2-N2-C7	-3.48	118.24	122.90
3	A	704	NAG	C3-C4-C5	-3.27	104.31	110.23
3	A	706	NAG	C2-N2-C7	-3.26	118.53	122.90
3	A	702	NAG	C1-O5-C5	3.06	116.29	112.19
3	A	709	NAG	O5-C5-C6	3.00	113.50	107.66
3	A	702	NAG	C2-N2-C7	-2.97	118.92	122.90
3	A	709	NAG	C2-N2-C7	-2.75	119.34	122.82
3	A	701	NAG	C6-C5-C4	-2.52	106.83	113.02
3	A	704	NAG	C2-N2-C7	-2.46	119.60	122.90
3	A	708	NAG	O5-C1-C2	-2.36	107.65	111.29
3	A	704	NAG	O5-C5-C6	2.35	112.23	107.66
3	A	704	NAG	C1-O5-C5	-2.18	109.26	112.19
3	A	709	NAG	O4-C4-C5	2.15	114.54	110.08
3	A	705	NAG	C3-C4-C5	-2.02	106.56	110.23

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	709	NAG	C4-C5-C6-O6
3	A	709	NAG	O5-C5-C6-O6
3	A	704	NAG	C4-C5-C6-O6
3	A	708	NAG	O5-C5-C6-O6
3	A	708	NAG	C4-C5-C6-O6
3	A	704	NAG	O5-C5-C6-O6
3	A	707	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	707	NAG	O5-C5-C6-O6
3	A	702	NAG	C8-C7-N2-C2
3	A	702	NAG	O7-C7-N2-C2

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	709	NAG	3	0
3	A	705	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	610/631 (96%)	-0.05	13 (2%) 63 60	1, 10, 28, 52	0
2	P	3/5 (60%)	0.36	0 100 100	3, 3, 8, 20	0
All	All	613/636 (96%)	-0.05	13 (2%) 63 60	1, 10, 28, 52	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	411	SER	4.4
1	A	409	ASP	3.8
1	A	511	GLU	3.8
1	A	493	ASN	3.2
1	A	509	ARG	3.0
1	A	109	ARG	2.9
1	A	639	SER	2.8
1	A	492	ARG	2.7
1	A	514	ARG	2.6
1	A	74	ASN	2.6
1	A	494	ILE	2.2
1	A	180	ALA	2.1
1	A	635	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TYS	P	30	16/17	0.95	0.07	1,6,22,37	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	TYS	P	28	16/17	0.96	0.07	2,4,13,18	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	706	14/15	0.69	0.13	15,30,36,46	0
3	NAG	A	709	13/15	0.71	0.16	19,36,42,46	0
3	NAG	A	707	14/15	0.87	0.11	12,16,28,33	0
3	NAG	A	708	14/15	0.88	0.12	8,28,36,37	0
3	NAG	A	704	14/15	0.89	0.10	1,7,17,19	0
3	NAG	A	705	14/15	0.90	0.09	9,17,27,28	0
3	NAG	A	703	14/15	0.91	0.09	9,15,22,31	0
3	NAG	A	702	14/15	0.96	0.07	4,7,12,14	0
3	NAG	A	701	14/15	0.96	0.07	2,5,8,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.