



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:06 PM UTC

PDB ID : 6Z6A / pdb_00006z6a
Title : Keap1 macrocycle complex
Authors : Johansson, P.
Deposited on : 2020-05-28
Resolution : 2.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7164 (2.40-2.36)
Clashscore	190562	7722 (2.40-2.36)
Ramachandran outliers	187476	7626 (2.40-2.36)
Sidechain outliers	187428	7627 (2.40-2.36)
RSRZ outliers	180081	7170 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	
1	B	295	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4590 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Kelch-like ECH-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	285	Total	C	N	O	S	0	0	0
			2186	1359	397	415	15			
1	B	285	Total	C	N	O	S	0	0	0
			2186	1359	397	415	15			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	315	PRO	-	expression tag	UNP Q14145
A	316	ARG	-	expression tag	UNP Q14145
A	317	GLY	-	expression tag	UNP Q14145
A	318	SER	-	expression tag	UNP Q14145
A	319	ALA	-	expression tag	UNP Q14145
A	320	GLY	-	expression tag	UNP Q14145
A	540	ALA	GLU	conflict	UNP Q14145
A	542	ALA	GLU	conflict	UNP Q14145
B	315	PRO	-	expression tag	UNP Q14145
B	316	ARG	-	expression tag	UNP Q14145
B	317	GLY	-	expression tag	UNP Q14145
B	318	SER	-	expression tag	UNP Q14145
B	319	ALA	-	expression tag	UNP Q14145
B	320	GLY	-	expression tag	UNP Q14145
B	540	ALA	GLU	conflict	UNP Q14145
B	542	ALA	GLU	conflict	UNP Q14145

- Molecule 2 is (5S,8R)-8-[[[(2S)-1-ethanoylpyrrolidin-2-yl]carbonylamino]-N,N-dimethyl-7,11-bis(oxidanylidene)-10-oxa-3-thia-6-azabicyclo[10.4.0]hexadeca-1(16),12,14-triene-5-carboxamide (CCD ID: Q9E) (formula: C₂₃H₃₀N₄O₆S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			34	23	4	6	1		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	101	Total	O	0	0
			101	101		
5	B	81	Total	O	0	0
			81	81		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kelch-like ECH-associated protein 1



- Molecule 1: Kelch-like ECH-associated protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.65Å 75.80Å 203.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.37 – 2.37 47.37 – 2.37	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.37-2.37) 99.7 (47.37-2.37)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 2.37Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.192 , 0.209 0.192 , 0.209	Depositor DCC
R_{free} test set	2356 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	66.5	Xtriage
Anisotropy	0.207	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.024 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4590	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: Q9E, CL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	1/2239 (0.0%)	1.24	14/3048 (0.5%)
1	B	0.84	0/2239	1.26	14/3048 (0.5%)
All	All	0.85	1/4478 (0.0%)	1.25	28/6096 (0.5%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	360	VAL	CA-C	5.59	1.56	1.52

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	573	ASP	CA-CB-CG	7.08	119.68	112.60
1	B	438	ASN	N-CA-C	-6.95	104.69	113.72
1	A	438	ASN	N-CA-C	-6.70	105.02	113.72
1	B	483	ARG	N-CA-C	-6.43	101.43	110.50
1	A	479	ASP	CA-CB-CG	6.27	118.87	112.60
1	B	573	ASP	CA-CB-CG	6.23	118.83	112.60
1	B	526	ASP	CA-CB-CG	6.05	118.65	112.60
1	B	479	ASP	CA-CB-CG	6.05	118.65	112.60
1	B	438	ASN	CA-CB-CG	5.95	118.55	112.60
1	A	599	SER	CA-C-N	5.94	126.37	120.31
1	A	599	SER	C-N-CA	5.94	126.37	120.31
1	B	485	ASN	N-CA-C	-5.82	106.15	113.72
1	A	438	ASN	CA-CB-CG	5.69	118.29	112.60
1	A	485	ASN	N-CA-C	-5.62	106.42	113.72
1	B	524	GLY	CA-C-N	5.50	130.53	122.77
1	B	524	GLY	C-N-CA	5.50	130.53	122.77
1	A	385	ASP	CA-CB-CG	5.48	118.08	112.60
1	B	357	ASP	CA-CB-CG	5.42	118.02	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	415	ARG	CA-C-N	5.39	129.88	121.85
1	A	415	ARG	C-N-CA	5.39	129.88	121.85
1	A	579	ASP	N-CA-C	-5.38	106.72	113.72
1	B	579	ASP	N-CA-C	-5.26	106.88	113.72
1	A	579	ASP	CA-CB-CG	5.24	117.84	112.60
1	B	579	ASP	CA-CB-CG	5.16	117.76	112.60
1	A	357	ASP	CA-CB-CG	5.14	117.74	112.60
1	A	484	LEU	N-CA-C	5.05	117.71	109.59
1	B	415	ARG	CA-C-N	5.04	129.68	121.62
1	B	415	ARG	C-N-CA	5.04	129.68	121.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2186	0	2081	14	0
1	B	2186	0	2081	12	0
2	A	34	0	0	0	0
3	A	1	0	0	0	0
4	A	1	0	0	1	0
5	A	101	0	0	0	0
5	B	81	0	0	1	0
All	All	4590	0	4162	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:353:LEU:HD13	5:B:758:HOH:O	2.03	0.57
1:A:466:ALA:HB1	1:A:514:VAL:HG23	1.86	0.56
1:B:466:ALA:HB1	1:B:514:VAL:HG23	1.87	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:415:ARG:NH1	4:A:703:CL:CL	2.77	0.55
1:B:369:VAL:HG21	1:B:609:THR:HB	1.90	0.53
1:A:369:VAL:HG21	1:A:609:THR:HB	1.92	0.51
1:A:483:ARG:HB3	1:A:506:ILE:CG2	2.42	0.50
1:B:416:ILE:HD11	1:B:427:ALA:HB1	1.95	0.47
1:A:416:ILE:HD11	1:A:427:ALA:HB1	1.97	0.47
1:B:380:ARG:HD2	1:B:382:ASN:HB2	1.96	0.46
1:A:472:LEU:HB3	1:A:490:TYR:HB3	1.97	0.46
1:B:472:LEU:HB3	1:B:490:TYR:HB3	1.99	0.44
1:A:489:CYS:O	1:A:497:TRP:HA	2.17	0.44
1:B:515:LEU:HD22	1:B:566:ILE:HG13	2.00	0.44
1:B:464:GLY:HA3	1:B:512:VAL:CG2	2.48	0.44
1:A:515:LEU:HD22	1:A:566:ILE:HG13	2.01	0.43
1:B:489:CYS:O	1:B:497:TRP:HA	2.18	0.43
1:A:362:ARG:HD2	1:A:378:GLY:O	2.20	0.42
1:B:485:ASN:HB3	1:B:506:ILE:HG13	2.02	0.42
1:B:362:ARG:HD2	1:B:378:GLY:O	2.20	0.42
1:A:464:GLY:HA3	1:A:512:VAL:CG2	2.50	0.41
1:A:335:PHE:C	1:A:337:GLN:H	2.28	0.41
1:A:485:ASN:HB3	1:A:506:ILE:HG13	2.03	0.41
1:A:569:LEU:HD23	1:A:597:MET:HE3	2.03	0.41
1:A:380:ARG:HD3	1:A:382:ASN:OD1	2.21	0.41
1:B:569:LEU:HD23	1:B:597:MET:HE3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/295 (96%)	275 (97%)	8 (3%)	0	100	100
1	B	283/295 (96%)	275 (97%)	8 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	566/590 (96%)	550 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/235 (97%)	218 (95%)	11 (5%)	23	37
1	B	229/235 (97%)	220 (96%)	9 (4%)	28	45
All	All	458/470 (97%)	438 (96%)	20 (4%)	25	40

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	353	LEU
1	A	377	VAL
1	A	415	ARG
1	A	481	THR
1	A	483	ARG
1	A	506	ILE
1	A	528	GLN
1	A	529	ASP
1	A	576	THR
1	A	593	GLU
1	A	596	ARG
1	B	336	ARG
1	B	353	LEU
1	B	377	VAL
1	B	461	ILE
1	B	481	THR
1	B	483	ARG
1	B	506	ILE
1	B	529	ASP
1	B	593	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	359	GLN
1	A	482	ASN
1	B	359	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	Q9E	A	701	-	36,36,36	1.15	3 (8%)	50,50,50	1.48	10 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	Q9E	A	701	-	-	8/43/53/53	0/2/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	Q9E	C10-N27	4.60	1.40	1.34
2	A	701	Q9E	C11-N25	2.81	1.39	1.35
2	A	701	Q9E	C9-N26	2.31	1.39	1.34

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	Q9E	C16-O33-C7	-4.03	109.52	116.45
2	A	701	Q9E	O32-C11-N25	2.72	123.59	120.45
2	A	701	Q9E	C9-C19-N25	-2.70	105.13	112.50
2	A	701	Q9E	C19-N25-C11	2.66	125.70	120.73
2	A	701	Q9E	C12-C6-C5	2.34	125.73	122.73
2	A	701	Q9E	C19-C9-N26	-2.29	111.49	116.52
2	A	701	Q9E	C14-C19-N25	2.20	106.24	103.02
2	A	701	Q9E	C21-C11-N25	-2.16	115.66	117.99
2	A	701	Q9E	C12-C6-C4	-2.14	116.08	119.46
2	A	701	Q9E	O30-C9-N26	2.11	126.74	122.96

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	701	Q9E	C21-C11-N25-C15
2	A	701	Q9E	C21-C11-N25-C19
2	A	701	Q9E	O32-C11-N25-C15
2	A	701	Q9E	O32-C11-N25-C19
2	A	701	Q9E	O33-C16-C18-C8
2	A	701	Q9E	O33-C16-C18-N26
2	A	701	Q9E	C5-C7-O33-C16
2	A	701	Q9E	O28-C7-O33-C16

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	285/295 (96%)	0.18	6 (2%) 63 62	52, 66, 90, 110	0
1	B	285/295 (96%)	0.26	12 (4%) 40 40	55, 67, 93, 126	0
All	All	570/590 (96%)	0.22	18 (3%) 50 50	52, 66, 92, 126	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	325	GLY	4.9
1	B	525	TYR	4.7
1	B	527	GLY	4.1
1	A	325	GLY	3.9
1	B	529	ASP	3.9
1	B	528	GLN	3.7
1	B	526	ASP	3.6
1	A	575	HIS	2.8
1	B	384	PRO	2.8
1	B	575	HIS	2.7
1	B	483	ARG	2.7
1	A	434	CYS	2.4
1	B	489	CYS	2.3
1	B	383	SER	2.2
1	A	577	PHE	2.2
1	B	380	ARG	2.2
1	A	384	PRO	2.1
1	A	489	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	Q9E	A	701	34/34	0.93	0.13	66,75,90,91	0
3	NA	A	702	1/1	0.95	0.43	71,71,71,71	0
4	CL	A	703	1/1	0.98	0.12	78,78,78,78	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.