



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:18 AM UTC

PDB ID : 6Z6R / pdb_00006z6r
Title : Aspartyl/Asparaginyl beta-hydroxylase (AspH) oxygenase and TPR domains in complex with manganese, N-oxalyl-alpha-methylalanine, and factor X substrate peptide fragment(39mer-4Ser)
Authors : Nakashima, Y.; Brewitz, L.; Schofield, C.J.
Deposited on : 2020-05-29
Resolution : 2.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	8% 73% 26% .
2	B	39	10% 41% 5% 54%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	UQK	A	804	-	X	-	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7156 atoms, of which 3472 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartyl/asparaginyl beta-hydroxylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	429	Total	C	H	N	O	S	0	0	0
			6769	2174	3341	612	630	12			

- Molecule 2 is a protein called Coagulation factor X.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	18	Total	C	H	N	O	S	0	0	0
			243	79	116	20	26	2			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	90	SER	CYS	engineered mutation	UNP P00742
B	95	SER	CYS	engineered mutation	UNP P00742
B	112	SER	CYS	engineered mutation	UNP P00742
B	121	SER	CYS	engineered mutation	UNP P00742

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		

- Molecule 4 is BROMIDE ION (CCD ID: BR) (formula: Br).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Br	0	0
			1	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 6 is N-oxalyl-alpha-methylalanine (CCD ID: UQK) (formula: $C_6H_9NO_5$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	H	N	O	0	0
			19	6	7	1	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	106	Total 106	O 106	0	0
7	B	3	Total 3	O 3	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aspartyl/asparaginyl beta-hydroxylase



- Molecule 2: Coagulation factor X



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.85Å 90.59Å 122.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.70 – 2.13 50.70 – 2.13	Depositor EDS
% Data completeness (in resolution range)	66.3 (50.70-2.13) 66.3 (50.70-2.13)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 2.14Å)	Xtriage
Refinement program	PHENIX 1.17.1 _3660	Depositor
R, R_{free}	0.212 , 0.242 0.233 , 0.272	Depositor DCC
R_{free} test set	1011 reflections (3.19%)	wwPDB-VP
Wilson B-factor (Å ²)	35.7	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7156	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UQK, BR, GOL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/3510	0.93	12/4749 (0.3%)
2	B	0.94	0/128	1.10	0/169
All	All	0.72	0/3638	0.93	12/4918 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	484	THR	O-C-N	10.46	128.56	121.23
1	A	355	GLY	CA-C-N	-7.62	112.29	122.42
1	A	355	GLY	C-N-CA	-7.62	112.29	122.42
1	A	541	LYS	CA-C-N	-6.56	112.06	122.95
1	A	541	LYS	C-N-CA	-6.56	112.06	122.95
1	A	689	MET	CG-SD-CE	6.20	114.53	100.90
1	A	545	LYS	O-C-N	6.13	130.75	122.59
1	A	383	GLN	CA-C-N	-5.45	111.41	120.68
1	A	383	GLN	C-N-CA	-5.45	111.41	120.68
1	A	384	CYS	O-C-N	-5.26	115.28	122.33
1	A	483	VAL	CA-C-N	-5.10	116.49	122.00
1	A	483	VAL	C-N-CA	-5.10	116.49	122.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	396	GLU	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3428	3341	3340	87	0
2	B	127	116	116	1	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	6	8	8	0	0
6	A	12	7	0	0	0
7	A	106	0	0	4	1
7	B	3	0	0	0	0
All	All	3684	3472	3464	88	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:570:LEU:HD11	1:A:747:LEU:HD21	1.56	0.85
1:A:542:GLU:O	1:A:545:LYS:CB	2.31	0.79
1:A:586:LEU:HD23	1:A:689:MET:CE	2.18	0.73
1:A:492:VAL:HG13	1:A:514:GLY:HA3	1.71	0.72
1:A:406:GLN:OE1	1:A:426:ARG:NH1	2.23	0.71
1:A:352:ARG:HD3	1:A:384:CYS:SG	2.32	0.69
1:A:497:ILE:O	1:A:501:GLN:HG3	1.91	0.69
2:B:105:LEU:HD23	2:B:110:CYS:SG	2.32	0.69
1:A:748:THR:HG23	1:A:751:GLN:OE1	1.92	0.69
1:A:586:LEU:HD23	1:A:689:MET:HE1	1.73	0.69
1:A:352:ARG:CD	1:A:384:CYS:SG	2.83	0.66
1:A:352:ARG:HB2	1:A:384:CYS:SG	2.39	0.61
1:A:499:LYS:HD2	1:A:530:HIS:HB3	1.82	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:563:SER:OG	1:A:566:ASN:OD1	2.14	0.61
1:A:607:LYS:O	7:A:901:HOH:O	2.17	0.60
1:A:596:LEU:HD21	1:A:646:LYS:HD2	1.85	0.57
1:A:574:PRO:O	1:A:719:PHE:HB2	2.04	0.57
1:A:386:ASP:CG	1:A:425:ARG:HH21	2.12	0.57
1:A:389:ALA:N	1:A:397:VAL:HG11	2.21	0.56
1:A:540:ASN:OD1	1:A:540:ASN:C	2.49	0.55
1:A:483:VAL:O	1:A:483:VAL:HG12	2.06	0.55
1:A:540:ASN:OD1	1:A:541:LYS:N	2.40	0.54
1:A:661:ARG:HD2	7:A:902:HOH:O	2.07	0.54
1:A:352:ARG:CG	1:A:384:CYS:SG	2.96	0.54
1:A:369:ARG:HG3	1:A:370:LYS:N	2.23	0.54
1:A:602:LEU:HD23	1:A:605:MET:SD	2.48	0.54
1:A:484:THR:HG22	1:A:484:THR:O	2.07	0.53
1:A:375:PRO:HA	1:A:411:LEU:HD12	1.90	0.53
1:A:661:ARG:NH1	7:A:902:HOH:O	2.23	0.52
1:A:404:THR:O	1:A:407:GLU:HB2	2.10	0.52
1:A:524:ASP:HB3	1:A:527:PHE:CD2	2.44	0.52
1:A:691:LEU:HD23	1:A:693:LEU:HD21	1.92	0.52
1:A:504:ILE:O	1:A:508:ILE:HG13	2.10	0.51
1:A:628:PHE:HB3	1:A:667:TYR:HB2	1.93	0.50
1:A:366:GLU:OE2	1:A:369:ARG:NE	2.38	0.50
1:A:604:VAL:HG21	1:A:628:PHE:CE2	2.47	0.50
1:A:586:LEU:HD23	1:A:689:MET:HE3	1.93	0.49
1:A:454:THR:HG23	1:A:480:VAL:HG13	1.96	0.48
1:A:583:TYR:CD1	1:A:656:GLU:HG3	2.48	0.48
1:A:605:MET:HE3	1:A:669:ILE:HD12	1.95	0.48
1:A:483:VAL:O	1:A:483:VAL:CG1	2.61	0.48
1:A:352:ARG:NH1	1:A:387:ASP:OD2	2.44	0.48
1:A:528:TYR:CE2	1:A:555:HIS:CE1	3.02	0.47
1:A:364:PHE:CE1	1:A:380:GLY:HA3	2.50	0.47
1:A:586:LEU:CD2	1:A:689:MET:HE1	2.42	0.46
1:A:433:LEU:HB3	1:A:435:HIS:HE1	1.80	0.46
1:A:505:ALA:HA	1:A:508:ILE:HD12	1.97	0.46
1:A:748:THR:OG1	1:A:751:GLN:HG3	2.16	0.45
1:A:478:GLU:O	1:A:482:SER:HB2	2.16	0.45
1:A:594:TRP:HB2	1:A:691:LEU:HD21	1.98	0.45
1:A:662:ARG:HG3	1:A:743:TRP:CH2	2.51	0.45
1:A:420:LYS:O	1:A:424:LYS:HG3	2.17	0.45
1:A:433:LEU:HB3	1:A:435:HIS:CE1	2.51	0.45
1:A:680:THR:HG22	1:A:681:GLY:O	2.16	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:436:MET:SD	1:A:466:LEU:HB3	2.58	0.44
1:A:549:LEU:O	1:A:552:LYS:N	2.51	0.44
1:A:417:ASP:O	1:A:421:LEU:HD12	2.18	0.44
1:A:637:ASN:HB3	1:A:640:ALA:HB3	1.99	0.44
1:A:528:TYR:HE2	1:A:555:HIS:HE1	1.65	0.44
1:A:720:ASP:HB3	1:A:723:PHE:CD1	2.52	0.44
1:A:350:LYS:O	1:A:351:LEU:C	2.59	0.44
1:A:499:LYS:HE3	1:A:533:ASP:HB3	1.99	0.43
1:A:535:MET:HA	1:A:538:VAL:HG22	1.99	0.43
1:A:639:ASN:O	1:A:642:LYS:HB2	2.18	0.43
1:A:402:ILE:HG23	1:A:426:ARG:HG3	2.01	0.43
1:A:659:GLY:O	1:A:743:TRP:NE1	2.49	0.43
1:A:689:MET:HB2	1:A:740:VAL:O	2.18	0.43
1:A:453:ASP:OD1	1:A:455:SER:N	2.52	0.42
1:A:407:GLU:O	1:A:410:SER:HB2	2.19	0.42
1:A:702:ILE:HG23	1:A:702:ILE:O	2.19	0.42
1:A:755:LEU:HD23	7:A:917:HOH:O	2.19	0.42
1:A:594:TRP:CG	1:A:714:GLY:HA2	2.54	0.42
1:A:653:LYS:O	1:A:653:LYS:CG	2.66	0.42
1:A:743:TRP:CD1	1:A:752:ARG:HG2	2.55	0.42
1:A:395:ASN:C	1:A:395:ASN:OD1	2.62	0.42
1:A:506:GLU:O	1:A:509:PRO:HD2	2.19	0.42
1:A:479:GLU:O	1:A:480:VAL:C	2.62	0.42
1:A:352:ARG:CB	1:A:384:CYS:SG	3.07	0.41
1:A:605:MET:HE1	1:A:734:PHE:CE2	2.55	0.41
1:A:499:LYS:CD	1:A:530:HIS:HB3	2.49	0.41
1:A:583:TYR:CE1	1:A:656:GLU:HB2	2.55	0.41
1:A:431:GLN:HA	1:A:436:MET:HE2	2.02	0.41
1:A:378:ARG:HB3	1:A:408:VAL:HG22	2.02	0.41
1:A:378:ARG:HD2	1:A:407:GLU:OE1	2.21	0.41
1:A:385:GLU:HB2	1:A:401:ALA:HB2	2.02	0.41
1:A:615:GLU:O	1:A:620:ARG:NH2	2.52	0.40
1:A:528:TYR:CE2	1:A:555:HIS:HE1	2.39	0.40
1:A:723:PHE:CD1	1:A:723:PHE:N	2.90	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:971:HOH:O	7:A:1002:HOH:O[3_745]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	427/429 (100%)	407 (95%)	20 (5%)	0	100	100
2	B	16/39 (41%)	13 (81%)	3 (19%)	0	100	100
All	All	443/468 (95%)	420 (95%)	23 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/371 (96%)	356 (100%)	0	100	100
2	B	13/33 (39%)	13 (100%)	0	100	100
All	All	369/404 (91%)	369 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	UQK	A	804	3	11,11,11	3.48	7 (63%)	13,16,16	3.73	5 (38%)
5	GOL	A	803	-	5,5,5	0.78	0	5,5,5	1.27	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	UQK	A	804	3	-	7/14/15/15	-
5	GOL	A	803	-	-	2/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	804	UQK	C05-C07	-6.28	1.46	1.54
6	A	804	UQK	C05-N	5.26	1.41	1.34
6	A	804	UQK	CA-C	-4.33	1.44	1.53
6	A	804	UQK	CA-N	-4.02	1.40	1.47
6	A	804	UQK	O06-C05	-3.75	1.16	1.23
6	A	804	UQK	OXT-C	-2.54	1.21	1.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	804	UQK	O09-C07	-2.27	1.24	1.30

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	804	UQK	C03-CA-CB	10.15	125.33	110.40
6	A	804	UQK	OXT-C-O	-5.07	107.63	123.86
6	A	804	UQK	C03-CA-C	-4.66	95.36	108.65
6	A	804	UQK	OXT-C-CA	3.78	127.41	114.93
5	A	803	GOL	C3-C2-C1	-2.51	102.59	111.80
6	A	804	UQK	O09-C07-O08	-2.24	118.57	123.90

There are no chirality outliers.


All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	803	GOL	O1-C1-C2-C3
6	A	804	UQK	OXT-C-CA-C03
6	A	804	UQK	C03-CA-N-C05
6	A	804	UQK	C-CA-N-C05
6	A	804	UQK	CB-CA-N-C05
6	A	804	UQK	O-C-CA-C03
5	A	803	GOL	O1-C1-C2-O2
6	A	804	UQK	OXT-C-CA-N
6	A	804	UQK	O-C-CA-N

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	429/429 (100%)	0.67	33 (7%)	19 22	17, 48, 77, 102	0
2	B	18/39 (46%)	1.13	4 (22%)	2 3	31, 49, 78, 91	0
All	All	447/468 (95%)	0.69	37 (8%)	17 20	17, 48, 77, 102	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	113	LEU	4.5
1	A	549	LEU	4.3
1	A	397	VAL	3.6
1	A	569	GLY	3.5
1	A	388	LEU	3.4
1	A	384	CYS	3.4
1	A	546	TRP	3.4
2	B	116	PHE	3.4
1	A	481	LEU	3.3
1	A	398	LEU	3.2
1	A	719	PHE	3.2
1	A	543	ALA	2.9
1	A	508	ILE	2.9
1	A	395	ASN	2.9
1	A	560	TRP	2.9
1	A	557	ALA	2.7
1	A	345	LEU	2.5
1	A	361	VAL	2.5
1	A	433	LEU	2.5
1	A	705	ALA	2.4
1	A	441	LEU	2.4
1	A	570	LEU	2.4
1	A	531	LEU	2.4
1	A	432	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	516	GLU	2.3
1	A	544	TYR	2.3
1	A	357	ILE	2.3
1	A	343	ALA	2.3
1	A	528	TYR	2.3
1	A	484	THR	2.2
2	B	105	LEU	2.2
1	A	697	LYS	2.2
2	B	115	GLY	2.1
1	A	706	ASN	2.1
1	A	556	PHE	2.1
1	A	476	VAL	2.1
1	A	442	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	A	803	6/6	0.89	0.11	39,48,73,88	0
3	MN	A	801	1/1	0.90	0.08	46,46,46,46	0
6	UQK	A	804	12/12	0.92	0.14	24,55,67,78	0
4	BR	A	802	1/1	0.97	0.05	52,52,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.