



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:45 AM UTC

PDB ID : 7Z61 / pdb_00007z61
Title : Crystal structure of PI3Kgamma with a dihydropurinone inhibitor (compound 18)
Authors : Goldberg, F.W.; Ting, A.K.T.; Schimpl, M.
Deposited on : 2022-03-10
Resolution : 2.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1819 (2.76-2.72)
Clashscore	190562	1866 (2.76-2.72)
Ramachandran outliers	187476	1830 (2.76-2.72)
Sidechain outliers	187428	1831 (2.76-2.72)
RSRZ outliers	180081	1819 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	966	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6778 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit gamma isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	834	Total	C	N	O	S	0	0	0
			6749	4333	1148	1234	34			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	143	MET	-	initiating methionine	UNP P48736
A	1103	HIS	-	expression tag	UNP P48736
A	1104	HIS	-	expression tag	UNP P48736
A	1105	HIS	-	expression tag	UNP P48736
A	1106	HIS	-	expression tag	UNP P48736
A	1107	HIS	-	expression tag	UNP P48736
A	1108	HIS	-	expression tag	UNP P48736

- Molecule 2 is 9-[(3 {R},4 {R})-4-fluoranylpyrrolidin-3-yl]-7-methyl-2-[(7-methylquinolin-6-yl)amino]purin-8-one (CCD ID: IGJ) (formula: C₂₀H₂₀FN₇O) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
2	A	1	29	20	1	7	1	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit gamma isoform



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	142.89Å 67.95Å 107.46Å 90.00° 95.92° 90.00°	Depositor
Resolution (Å)	62.21 – 2.74 62.21 – 2.74	Depositor EDS
% Data completeness (in resolution range)	98.9 (62.21-2.74) 98.9 (62.21-2.74)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 2.73Å)	Xtriage
Refinement program	BUSTER 2.11.8 (24-FEB-2021)	Depositor
R, R_{free}	0.217 , 0.276 0.216 , 0.280	Depositor DCC
R_{free} test set	1317 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	111.9	Xtriage
Anisotropy	0.132	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 132.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6778	wwPDB-VP
Average B, all atoms (Å ²)	141.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IGJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	1/6894 (0.0%)	1.16	15/9328 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	899	THR	CA-C	6.18	1.60	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	578	PHE	CA-C-N	9.04	132.40	120.28
1	A	578	PHE	C-N-CA	9.04	132.40	120.28
1	A	898	ASN	CA-C-N	7.53	134.60	122.10
1	A	898	ASN	C-N-CA	7.53	134.60	122.10
1	A	513	SER	CA-C-N	7.34	131.92	121.42
1	A	513	SER	C-N-CA	7.34	131.92	121.42
1	A	593	PHE	CA-CB-CG	-6.74	107.06	113.80
1	A	470	ASP	CA-CB-CG	6.36	118.96	112.60
1	A	961	PHE	CA-CB-CG	5.82	119.62	113.80
1	A	157	LEU	CA-C-N	5.38	127.83	120.46
1	A	157	LEU	C-N-CA	5.38	127.83	120.46
1	A	991	PHE	CA-CB-CG	5.34	119.14	113.80
1	A	216	ALA	N-CA-C	5.17	116.73	111.14
1	A	905	GLU	N-CA-C	5.17	119.23	113.02
1	A	694	PHE	CA-CB-CG	-5.03	108.77	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6749	0	6765	106	0
2	A	29	0	0	0	0
All	All	6778	0	6765	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:899:THR:HB	1:A:901:ALA:H	1.19	1.03
1:A:1000:LYS:H	1:A:1076:ARG:HH12	1.11	0.99
1:A:214:LYS:NZ	1:A:300:GLY:HA2	1.83	0.92
1:A:939:THR:HG23	1:A:945:GLY:HA2	1.53	0.89
1:A:899:THR:HB	1:A:901:ALA:N	1.96	0.81
1:A:175:PHE:HD1	1:A:472:ARG:HH21	1.29	0.80
1:A:834:HIS:HB2	1:A:876:ILE:HD12	1.63	0.80
1:A:214:LYS:HZ2	1:A:300:GLY:HA2	1.47	0.78
1:A:181:VAL:O	1:A:185:MET:HG3	1.84	0.77
1:A:193:PRO:HB2	1:A:313:PRO:HB3	1.67	0.76
1:A:225:HIS:CE1	1:A:304:HIS:HD2	2.03	0.76
1:A:214:LYS:HZ1	1:A:300:GLY:HA2	1.54	0.72
1:A:1013:CYS:HB3	1:A:1068:PHE:HE2	1.55	0.72
1:A:750:LYS:HZ1	1:A:808:LYS:HE3	1.57	0.70
1:A:497:PHE:HE1	1:A:1045:LYS:NZ	1.90	0.70
1:A:274:VAL:HG21	1:A:292:TRP:CD1	2.28	0.69
1:A:1035:LEU:HD12	1:A:1048:ILE:HD13	1.75	0.69
1:A:225:HIS:HE1	1:A:304:HIS:HD2	1.40	0.67
1:A:150:PHE:HE2	1:A:722:ARG:NH1	1.93	0.66
1:A:1014:VAL:HG11	1:A:1065:LYS:HG3	1.78	0.65
1:A:629:GLN:HG3	1:A:1029:ILE:HG13	1.78	0.65
1:A:1008:LYS:O	1:A:1012:ILE:HG13	1.97	0.65
1:A:428:LEU:HD22	1:A:465:ASN:HB3	1.81	0.63
1:A:274:VAL:HG11	1:A:292:TRP:CE2	2.34	0.63
1:A:497:PHE:HE1	1:A:1045:LYS:HZ3	1.43	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:VAL:HB	1:A:310:PRO:HG3	1.81	0.61
1:A:1013:CYS:HB3	1:A:1068:PHE:CE2	2.37	0.60
1:A:583:LEU:HD22	1:A:610:LEU:HD22	1.84	0.59
1:A:432:GLN:NE2	1:A:434:TYR:OH	2.35	0.59
1:A:750:LYS:NZ	1:A:808:LYS:HE3	2.17	0.59
1:A:1035:LEU:HA	1:A:1039:MET:HG2	1.85	0.59
1:A:804:MET:HE2	1:A:812:TRP:HB2	1.86	0.58
1:A:750:LYS:HE3	1:A:808:LYS:HA	1.86	0.57
1:A:888:ILE:HD12	1:A:907:LEU:HD11	1.85	0.57
1:A:748:ASP:HB3	1:A:770:LYS:NZ	2.19	0.57
1:A:564:LEU:HB2	1:A:1052:ARG:HD2	1.87	0.57
1:A:481:VAL:HG22	1:A:517:SER:HB3	1.86	0.56
1:A:657:LEU:HD11	1:A:690:ARG:HG2	1.86	0.56
1:A:830:ILE:HG21	1:A:878:MET:HE2	1.87	0.56
1:A:890:LYS:HA	1:A:893:GLN:HG2	1.89	0.54
1:A:274:VAL:HG11	1:A:292:TRP:NE1	2.23	0.54
1:A:364:LYS:HB3	1:A:519:LEU:HB3	1.89	0.53
1:A:381:VAL:HB	1:A:404:PHE:HB2	1.90	0.53
1:A:833:LYS:HZ1	1:A:836:ASP:HB2	1.74	0.52
1:A:947:ARG:NH2	1:A:963:ILE:O	2.43	0.52
1:A:576:TRP:O	1:A:579:ARG:HB2	2.10	0.51
1:A:702:GLU:O	1:A:706:SER:HB3	2.11	0.51
1:A:274:VAL:CG2	1:A:292:TRP:CD1	2.94	0.51
1:A:362:ARG:NH2	1:A:413:TRP:CD1	2.79	0.51
1:A:1039:MET:HB3	1:A:1040:PRO:HD2	1.93	0.50
1:A:225:HIS:HE1	1:A:304:HIS:CD2	2.24	0.50
1:A:235:VAL:HG13	1:A:244:ILE:HG21	1.93	0.50
1:A:912:LYS:NZ	1:A:921:PHE:CE2	2.79	0.49
1:A:929:VAL:HG11	1:A:1008:LYS:HD3	1.94	0.49
1:A:838:LEU:O	1:A:842:MET:HG3	2.12	0.49
1:A:223:VAL:HG12	1:A:225:HIS:CE1	2.48	0.49
1:A:395:CYS:HB3	1:A:416:PHE:HD2	1.76	0.49
1:A:743:GLN:NE2	1:A:872:THR:OG1	2.46	0.49
1:A:1067:TYR:O	1:A:1071:GLN:HG2	2.12	0.48
1:A:825:ASN:ND2	1:A:825:ASN:H	2.11	0.48
1:A:833:LYS:NZ	1:A:836:ASP:HB2	2.29	0.48
1:A:395:CYS:SG	1:A:417:SER:OG	2.71	0.48
1:A:834:HIS:HD2	1:A:876:ILE:HB	1.78	0.47
1:A:847:ILE:HG21	1:A:942:LEU:HD21	1.97	0.47
1:A:701:SER:HB2	1:A:871:SER:HB3	1.97	0.46
1:A:362:ARG:NH2	1:A:413:TRP:CG	2.83	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:169:HIS:CE1	1:A:466:LEU:HD12	2.50	0.46
1:A:274:VAL:CG1	1:A:292:TRP:NE1	2.79	0.46
1:A:623:ASP:HB3	1:A:626:LEU:HB2	1.98	0.46
1:A:622:LEU:HD13	1:A:647:LYS:HB3	1.98	0.46
1:A:193:PRO:HB2	1:A:313:PRO:CB	2.40	0.45
1:A:235:VAL:HG11	1:A:244:ILE:HD13	1.98	0.45
1:A:734:GLN:O	1:A:738:VAL:HG23	2.17	0.45
1:A:767:LEU:HD22	1:A:803:VAL:HG23	1.99	0.44
1:A:1002:THR:HG23	1:A:1006:PHE:HD2	1.82	0.44
1:A:551:LEU:HD22	1:A:574:LEU:HD21	1.99	0.44
1:A:605:ALA:O	1:A:609:GLN:HG3	2.17	0.44
1:A:748:ASP:CB	1:A:770:LYS:NZ	2.80	0.43
1:A:661:LEU:O	1:A:665:GLN:HG2	2.19	0.43
1:A:298:LYS:NZ	1:A:299:ASN:HD21	2.16	0.43
1:A:583:LEU:HD13	1:A:610:LEU:HD13	2.00	0.43
1:A:671:PRO:HB2	1:A:672:TYR:CD1	2.54	0.43
1:A:747:LEU:CD2	1:A:834:HIS:CG	3.02	0.43
1:A:161:ASP:O	1:A:164:ASP:HB2	2.19	0.43
1:A:747:LEU:HD21	1:A:834:HIS:CG	2.53	0.43
1:A:911:LEU:HD22	1:A:924:ALA:HB1	2.00	0.43
1:A:176:THR:HG23	1:A:674:ASP:HB2	2.01	0.42
1:A:918:GLU:O	1:A:922:GLN:HG2	2.20	0.42
1:A:362:ARG:HH21	1:A:413:TRP:CD1	2.38	0.42
1:A:565:ASN:HA	1:A:566:PRO:HD3	1.93	0.42
1:A:225:HIS:CE1	1:A:304:HIS:CD2	2.93	0.42
1:A:705:GLN:HG3	1:A:839:ARG:CZ	2.49	0.42
1:A:154:LEU:O	1:A:158:ILE:HG13	2.19	0.42
1:A:365:ILE:HD13	1:A:518:ILE:HG22	2.02	0.42
1:A:221:PHE:HE1	1:A:234:LYS:HG2	1.86	0.41
1:A:680:PHE:O	1:A:684:ARG:HG2	2.20	0.41
1:A:935:TYR:O	1:A:939:THR:HB	2.20	0.41
1:A:362:ARG:HH21	1:A:413:TRP:NE1	2.19	0.41
1:A:555:LEU:HD11	1:A:575:LEU:HD23	2.02	0.41
1:A:235:VAL:HG12	1:A:236:SER:N	2.35	0.41
1:A:272:LEU:HB3	1:A:305:VAL:HG11	2.03	0.41
1:A:169:HIS:CE1	1:A:466:LEU:CD1	3.04	0.41
1:A:467:LEU:O	1:A:476:ARG:HD2	2.21	0.41
1:A:747:LEU:HD21	1:A:834:HIS:CD2	2.56	0.41
1:A:197:ALA:HB2	1:A:316:ASP:OD2	2.21	0.41
1:A:935:TYR:CE2	1:A:961:PHE:HA	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	820/966 (85%)	764 (93%)	49 (6%)	7 (1%)	14	26

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	777	SER
1	A	967	HIS
1	A	1044	SER
1	A	227	SER
1	A	999	GLY
1	A	798	ILE
1	A	809	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	746/864 (86%)	689 (92%)	57 (8%)	12	22

All (57) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	209	GLU
1	A	214	LYS
1	A	240	THR
1	A	244	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	252	MET
1	A	282	VAL
1	A	285	THR
1	A	287	ILE
1	A	305	VAL
1	A	381	VAL
1	A	395	CYS
1	A	415	GLU
1	A	472	ARG
1	A	475	LEU
1	A	488	SER
1	A	520	LEU
1	A	570	GLU
1	A	574	LEU
1	A	587	LYS
1	A	610	LEU
1	A	613	ARG
1	A	615	GLU
1	A	626	LEU
1	A	629	GLN
1	A	683	LYS
1	A	711	GLN
1	A	728	MET
1	A	749	ILE
1	A	752	LEU
1	A	779	LEU
1	A	825	ASN
1	A	832	PHE
1	A	838	LEU
1	A	848	LEU
1	A	870	ILE
1	A	875	LYS
1	A	883	LYS
1	A	886	THR
1	A	898	ASN
1	A	908	ASN
1	A	918	GLU
1	A	939	THR
1	A	947	ARG
1	A	951	ASN
1	A	954	ILE
1	A	959	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	960	LEU
1	A	983	VAL
1	A	997	THR
1	A	1012	ILE
1	A	1029	ILE
1	A	1039	MET
1	A	1045	LYS
1	A	1046	GLU
1	A	1048	ILE
1	A	1049	GLU
1	A	1081	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (24) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	169	HIS
1	A	225	HIS
1	A	291	GLN
1	A	299	ASN
1	A	304	HIS
1	A	386	ASN
1	A	396	GLN
1	A	432	GLN
1	A	550	GLN
1	A	600	GLN
1	A	646	GLN
1	A	705	GLN
1	A	730	HIS
1	A	743	GLN
1	A	766	GLN
1	A	775	GLN
1	A	776	ASN
1	A	825	ASN
1	A	834	HIS
1	A	846	GLN
1	A	962	HIS
1	A	1010	GLN
1	A	1022	HIS
1	A	1041	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	IGJ	A	1201	-	31,33,33	0.27	0	37,49,49	1.41	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IGJ	A	1201	-	-	4/8/18/18	0/5/5/5

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1201	IGJ	C19-C6-N3	-6.11	122.20	126.50
2	A	1201	IGJ	C3-C2-N1	-4.29	106.94	113.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1201	IGJ	C16-C8-N4	-2.07	115.10	118.68

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1201	IGJ	C3-C2-N1-C1
2	A	1201	IGJ	C5-C2-N1-C6
2	A	1201	IGJ	C16-C8-N4-C7
2	A	1201	IGJ	C5-C2-N1-C1

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	834/966 (86%)	-0.08	8 (0%) 79 79	82, 138, 196, 210	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1075	CYS	2.9
1	A	1044	SER	2.5
1	A	746	THR	2.4
1	A	252	MET	2.3
1	A	393	VAL	2.3
1	A	987	LEU	2.3
1	A	599	GLY	2.2
1	A	968	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	IGJ	A	1201	29/29	0.92	0.08	104,109,113,113	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.