



wwPDB EM Validation Summary Report ⓘ

Jun 27, 2026 – 09:03 AM EDT

PDB ID : 9Z6Y / pdb_00009z6y
EMDB ID : EMD-73864
Title : Structure of the elongating EcDRT3 reverse transcriptase in complex with its non-coding RNA
Authors : Deng, P.; Gao, A.
Deposited on : 2025-11-14
Resolution : 2.60 Å(reported)
Based on initial model : .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev133
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	8728 (2.10 - 3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	426	
1	B	426	
1	C	426	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	D	426	
1	E	426	
1	F	426	
2	H	650	
2	I	650	
2	J	650	
2	K	650	
2	L	650	
2	M	650	
3	R	138	
3	S	138	
3	T	138	
3	U	138	
3	V	138	
3	W	138	
4	X	17	
4	Z	17	
4	b	17	
4	d	17	
4	f	17	
4	h	17	
5	Y	18	
5	a	18	
5	c	18	
5	e	18	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	g	18	39% 22% 78%
5	i	18	39% 22% 78%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 70614 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Drt3a reverse transcriptase protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	415	Total	C	N	O	S	0	0
			3444	2225	587	622	10		
1	B	415	Total	C	N	O	S	0	0
			3444	2225	587	622	10		
1	C	415	Total	C	N	O	S	0	0
			3444	2225	587	622	10		
1	D	415	Total	C	N	O	S	0	0
			3444	2225	587	622	10		
1	E	415	Total	C	N	O	S	0	0
			3444	2225	587	622	10		
1	F	415	Total	C	N	O	S	0	0
			3444	2225	587	622	10		

- Molecule 2 is a protein called Drt3b reverse transcriptase protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	H	626	Total	C	N	O	P	S	0	0
			5190	3366	846	958	1	19		
2	I	626	Total	C	N	O	P	S	0	0
			5190	3366	846	958	1	19		
2	J	626	Total	C	N	O	P	S	0	0
			5190	3366	846	958	1	19		
2	K	626	Total	C	N	O	P	S	0	0
			5190	3366	846	958	1	19		
2	L	626	Total	C	N	O	P	S	0	0
			5190	3366	846	958	1	19		
2	M	626	Total	C	N	O	P	S	0	0
			5190	3366	846	958	1	19		

- Molecule 3 is a RNA chain called non-coding RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	R	113	Total	C	N	O	P	0	0
			2406	1077	431	785	113		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
3	S	113	Total	C	N	O	P	0	0
			2406	1077	431	785	113		
3	T	113	Total	C	N	O	P	0	0
			2406	1077	431	785	113		
3	U	113	Total	C	N	O	P	0	0
			2406	1077	431	785	113		
3	V	113	Total	C	N	O	P	0	0
			2406	1077	431	785	113		
3	W	113	Total	C	N	O	P	0	0
			2406	1077	431	785	113		

- Molecule 4 is a DNA chain called cDNA (5'-D(P*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*T)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
4	X	17	Total	C	N	O	P	0	0
			356	170	58	111	17		
4	Z	17	Total	C	N	O	P	0	0
			356	170	58	111	17		
4	b	17	Total	C	N	O	P	0	0
			356	170	58	111	17		
4	d	17	Total	C	N	O	P	0	0
			356	170	58	111	17		
4	f	17	Total	C	N	O	P	0	0
			356	170	58	111	17		
4	h	17	Total	C	N	O	P	0	0
			356	170	58	111	17		

- Molecule 5 is a DNA chain called cDNA (5'-D(P*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*A)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
5	Y	18	Total	C	N	O	P	0	0
			360	171	72	99	18		
5	a	18	Total	C	N	O	P	0	0
			360	171	72	99	18		
5	c	18	Total	C	N	O	P	0	0
			360	171	72	99	18		
5	e	18	Total	C	N	O	P	0	0
			360	171	72	99	18		
5	g	18	Total	C	N	O	P	0	0
			360	171	72	99	18		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
5	i	18	Total	C	N	O	P	0	0
			360	171	72	99	18		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
6	A	1	Total	Mg	0
			1	1	
6	B	1	Total	Mg	0
			1	1	
6	C	1	Total	Mg	0
			1	1	
6	D	1	Total	Mg	0
			1	1	
6	E	1	Total	Mg	0
			1	1	
6	F	1	Total	Mg	0
			1	1	
6	H	2	Total	Mg	0
			2	2	
6	I	2	Total	Mg	0
			2	2	
6	J	2	Total	Mg	0
			2	2	
6	K	2	Total	Mg	0
			2	2	
6	L	2	Total	Mg	0
			2	2	
6	M	2	Total	Mg	0
			2	2	

- Molecule 7 is PYROPHOSPHATE 2- (CCD ID: POP) (formula: H₂O₇P₂).



Mol	Chain	Residues	Atoms			AltConf
7	H	1	Total	O	P	0
			9	7	2	
7	I	1	Total	O	P	0
			9	7	2	
7	J	1	Total	O	P	0
			9	7	2	
7	K	1	Total	O	P	0
			9	7	2	
7	L	1	Total	O	P	0
			9	7	2	
7	M	1	Total	O	P	0
			9	7	2	


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		AltConf
8	H	1	Total	O	0
			1	1	
8	I	1	Total	O	0
			1	1	
8	J	1	Total	O	0
			1	1	
8	K	1	Total	O	0
			1	1	
8	L	1	Total	O	0
			1	1	
8	M	1	Total	O	0
			1	1	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Drt3a reverse transcriptase protein



• Molecule 1: Drt3a reverse transcriptase protein




• Molecule 1: Drt3a reverse transcriptase protein





- Molecule 1: Drt3a reverse transcriptase protein

Chain D: 79% 19% .




- Molecule 1: Drt3a reverse transcriptase protein

Chain E: 79% 18% .




- Molecule 1: Drt3a reverse transcriptase protein

Chain F: 78% 19% .




- Molecule 2: Drt3b reverse transcriptase protein

Chain H: 80% 16% .




• Molecule 2: Drt3b reverse transcriptase protein

Chain I: 79% 17% •




• Molecule 2: Drt3b reverse transcriptase protein

Chain J: 80% 16% •




• Molecule 2: Drt3b reverse transcriptase protein

Chain K: 79% 17% •




• Molecule 2: Drt3b reverse transcriptase protein

Chain L: 79% 16%




• Molecule 2: Drt3b reverse transcriptase protein

Chain M: 79% 17%





• Molecule 3: non-coding RNA



Chain R: 30% 18% 43% 20% 18%





- Molecule 3: non-coding RNA




- Molecule 3: non-coding RNA







- Molecule 3: non-coding RNA










• Molecule 3: non-coding RNA



• Molecule 3: non-coding RNA




• Molecule 4: cDNA (5'-D(P*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*T)-3')




• Molecule 4: cDNA (5'-D(P*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*T)-3')







- Molecule 4: cDNA (5'-D(P*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*T)-3')




- Molecule 4: cDNA (5'-D(P*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*T)-3')



- Molecule 4: cDNA (5'-D(P*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*T)-3')



- Molecule 4: cDNA (5'-D(P*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*TP*GP*T)-3')



- Molecule 5: cDNA (5'-D(P*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*A)-3')



- Molecule 5: cDNA (5'-D(P*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*A)-3')



● Molecule 5: cDNA (5'-D(P*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*A
P*CP*A)-3')



● Molecule 5: cDNA (5'-D(P*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*A
P*CP*A)-3')



● Molecule 5: cDNA (5'-D(P*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*A
P*CP*A)-3')



● Molecule 5: cDNA (5'-D(P*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*AP*CP*A
P*CP*A)-3')



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	239792	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.621	Depositor
Minimum map value	-1.983	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.040	Depositor
Recommended contour level	0.1	Depositor
Map size (Å)	470.904, 470.904, 470.904	wwPDB
Map dimensions	560, 560, 560	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.8409, 0.8409, 0.8409	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PTR, POP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	3/3519 (0.1%)	0.46	2/4730 (0.0%)
1	B	0.39	3/3519 (0.1%)	0.46	2/4730 (0.0%)
1	C	0.39	3/3519 (0.1%)	0.46	2/4730 (0.0%)
1	D	0.39	3/3519 (0.1%)	0.46	2/4730 (0.0%)
1	E	0.39	3/3519 (0.1%)	0.46	2/4730 (0.0%)
1	F	0.39	3/3519 (0.1%)	0.46	2/4730 (0.0%)
2	H	0.46	7/5287 (0.1%)	0.44	6/7139 (0.1%)
2	I	0.46	7/5287 (0.1%)	0.44	6/7139 (0.1%)
2	J	0.46	7/5287 (0.1%)	0.44	6/7139 (0.1%)
2	K	0.46	7/5287 (0.1%)	0.44	6/7139 (0.1%)
2	L	0.46	7/5287 (0.1%)	0.44	6/7139 (0.1%)
2	M	0.48	9/5287 (0.2%)	0.45	6/7139 (0.1%)
3	R	0.35	0/2690	0.56	1/4186 (0.0%)
3	S	0.35	0/2690	0.56	1/4186 (0.0%)
3	T	0.35	0/2690	0.56	1/4186 (0.0%)
3	U	0.35	0/2690	0.56	1/4186 (0.0%)
3	V	0.35	0/2690	0.56	1/4186 (0.0%)
3	W	0.35	0/2690	0.56	1/4186 (0.0%)
4	X	0.40	0/397	0.56	0/614
4	Z	0.41	0/397	0.56	0/614
4	b	0.40	0/397	0.56	0/614
4	d	0.40	0/397	0.56	0/614
4	f	0.40	0/397	0.56	0/614
4	h	0.40	0/397	0.56	0/614
5	Y	0.36	0/404	0.56	0/617
5	a	0.36	0/404	0.56	0/617
5	c	0.36	0/404	0.56	0/617
5	e	0.36	0/404	0.56	0/617
5	g	0.36	0/404	0.56	0/617
5	i	0.36	0/404	0.56	0/617
All	All	0.41	62/73782 (0.1%)	0.49	54/103716 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	M	0	1

The worst 5 of 62 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	M	23	ILE	C-O	-7.39	1.19	1.24
2	I	23	ILE	C-O	-7.30	1.19	1.24
2	H	23	ILE	C-O	-7.21	1.19	1.24
2	K	23	ILE	C-O	-7.21	1.19	1.24
2	L	23	ILE	C-O	-7.19	1.19	1.24

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	196	VAL	N-CA-C	7.86	125.68	109.34
1	E	196	VAL	N-CA-C	7.85	125.67	109.34
1	B	196	VAL	N-CA-C	7.84	125.66	109.34
1	A	196	VAL	N-CA-C	7.83	125.62	109.34
1	C	196	VAL	N-CA-C	7.83	125.61	109.34

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	M	371	ASN	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3444	0	3451	60	0
1	B	3444	0	3450	58	0
1	C	3444	0	3451	59	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3444	0	3451	58	0
1	E	3444	0	3451	57	0
1	F	3444	0	3451	58	0
2	H	5190	0	5185	105	0
2	I	5190	0	5187	89	0
2	J	5190	0	5185	86	0
2	K	5190	0	5185	97	0
2	L	5190	0	5185	90	0
2	M	5190	0	5187	92	0
3	R	2406	0	1217	72	0
3	S	2406	0	1217	70	0
3	T	2406	0	1217	71	0
3	U	2406	0	1217	68	0
3	V	2406	0	1217	69	0
3	W	2406	0	1217	68	0
4	X	356	0	197	14	0
4	Z	356	0	197	26	0
4	b	356	0	197	19	0
4	d	356	0	197	16	0
4	f	356	0	197	16	0
4	h	356	0	197	15	0
5	Y	360	0	199	24	0
5	a	360	0	199	27	0
5	c	360	0	199	27	0
5	e	360	0	199	28	0
5	g	360	0	199	28	0
5	i	360	0	199	28	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	E	1	0	0	0	0
6	F	1	0	0	0	0
6	H	2	0	0	0	0
6	I	2	0	0	0	0
6	J	2	0	0	0	0
6	K	2	0	0	0	0
6	L	2	0	0	0	0
6	M	2	0	0	0	0
7	H	9	0	0	1	0
7	I	9	0	0	1	0
7	J	9	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	K	9	0	0	1	0
7	L	9	0	0	1	0
7	M	9	0	0	1	0
8	H	1	0	0	0	0
8	I	1	0	0	0	0
8	J	1	0	0	0	0
8	K	1	0	0	0	0
8	L	1	0	0	0	0
8	M	1	0	0	0	0
All	All	70614	0	61497	1307	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 1307 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:371:ASN:CG	4:Z:3:DT:H5''	1.14	1.53
2:H:371:ASN:ND2	4:Z:3:DT:C4'	1.70	1.53
2:M:371:ASN:OD1	4:d:3:DT:H5''	1.27	1.33
1:D:9:ARG:NH2	2:M:468:SER:O	1.65	1.30
1:C:9:ARG:NH2	2:I:468:SER:O	1.61	1.29

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	411/426 (96%)	406 (99%)	5 (1%)	0	100	100
1	B	411/426 (96%)	406 (99%)	5 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	411/426 (96%)	406 (99%)	5 (1%)	0	100	100
1	D	411/426 (96%)	406 (99%)	5 (1%)	0	100	100
1	E	411/426 (96%)	406 (99%)	5 (1%)	0	100	100
1	F	411/426 (96%)	406 (99%)	5 (1%)	0	100	100
2	H	620/650 (95%)	607 (98%)	13 (2%)	0	100	100
2	I	620/650 (95%)	607 (98%)	13 (2%)	0	100	100
2	J	620/650 (95%)	607 (98%)	13 (2%)	0	100	100
2	K	620/650 (95%)	607 (98%)	13 (2%)	0	100	100
2	L	620/650 (95%)	607 (98%)	13 (2%)	0	100	100
2	M	620/650 (95%)	606 (98%)	14 (2%)	0	100	100
All	All	6186/6456 (96%)	6077 (98%)	109 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	377/391 (96%)	376 (100%)	1 (0%)	86	94
1	B	377/391 (96%)	376 (100%)	1 (0%)	86	94
1	C	377/391 (96%)	376 (100%)	1 (0%)	86	94
1	D	377/391 (96%)	377 (100%)	0	100	100
1	E	377/391 (96%)	376 (100%)	1 (0%)	86	94
1	F	377/391 (96%)	377 (100%)	0	100	100
2	H	582/606 (96%)	578 (99%)	4 (1%)	76	89
2	I	582/606 (96%)	579 (100%)	3 (0%)	81	92
2	J	582/606 (96%)	579 (100%)	3 (0%)	81	92
2	K	582/606 (96%)	576 (99%)	6 (1%)	68	86
2	L	582/606 (96%)	577 (99%)	5 (1%)	70	87

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	M	582/606 (96%)	578 (99%)	4 (1%)	76	89
All	All	5754/5982 (96%)	5725 (100%)	29 (0%)	78	92

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	K	346	GLN
2	M	417	ASN
2	K	471	ASP
2	L	471	ASP
2	K	411	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
2	K	105	GLN
2	L	268	GLN
2	K	221	ASN
2	K	417	ASN
2	L	349	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	R	111/138 (80%)	56 (50%)	5 (4%)
3	S	111/138 (80%)	56 (50%)	5 (4%)
3	T	111/138 (80%)	56 (50%)	5 (4%)
3	U	111/138 (80%)	56 (50%)	5 (4%)
3	V	111/138 (80%)	56 (50%)	5 (4%)
3	W	111/138 (80%)	56 (50%)	5 (4%)
All	All	666/828 (80%)	336 (50%)	30 (4%)

5 of 336 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	R	6	A
3	R	8	C
3	R	10	A
3	R	12	U
3	R	13	A

5 of 30 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
3	T	115	A
3	W	105	U
3	U	105	U
3	W	115	A
3	V	115	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PTR	L	650	2	15,16,17	2.62	2 (13%)	17,22,24	1.18	1 (5%)
2	PTR	K	650	2	15,16,17	2.63	2 (13%)	17,22,24	1.17	1 (5%)
2	PTR	H	650	2	15,16,17	2.62	2 (13%)	17,22,24	1.17	1 (5%)
2	PTR	M	650	2	15,16,17	2.62	2 (13%)	17,22,24	1.18	1 (5%)
2	PTR	I	650	2	15,16,17	2.62	2 (13%)	17,22,24	1.17	1 (5%)
2	PTR	J	650	2	15,16,17	2.63	2 (13%)	17,22,24	1.18	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTR	L	650	2	-	1/10/11/13	0/1/1/1
2	PTR	K	650	2	-	1/10/11/13	0/1/1/1
2	PTR	H	650	2	-	1/10/11/13	0/1/1/1
2	PTR	M	650	2	-	1/10/11/13	0/1/1/1
2	PTR	I	650	2	-	1/10/11/13	0/1/1/1
2	PTR	J	650	2	-	1/10/11/13	0/1/1/1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	650	PTR	P-OH	9.56	1.77	1.59
2	I	650	PTR	P-OH	9.53	1.77	1.59
2	H	650	PTR	P-OH	9.53	1.77	1.59
2	K	650	PTR	P-OH	9.53	1.77	1.59
2	L	650	PTR	P-OH	9.52	1.77	1.59

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	650	PTR	O3P-P-O2P	2.14	115.83	107.80
2	H	650	PTR	O3P-P-O2P	2.13	115.81	107.80
2	K	650	PTR	O3P-P-O2P	2.13	115.80	107.80
2	L	650	PTR	O3P-P-O2P	2.13	115.79	107.80
2	M	650	PTR	O3P-P-O2P	2.13	115.77	107.80

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	650	PTR	N-CA-CB-CG
2	I	650	PTR	N-CA-CB-CG
2	J	650	PTR	N-CA-CB-CG
2	K	650	PTR	N-CA-CB-CG
2	L	650	PTR	N-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 18 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	POP	I	703	6	6,8,8	0.71	0	12,13,13	1.44	1 (8%)
7	POP	L	703	6	6,8,8	0.73	0	12,13,13	1.42	1 (8%)
7	POP	K	703	6	6,8,8	0.71	0	12,13,13	1.44	1 (8%)
7	POP	M	703	6	6,8,8	0.73	0	12,13,13	1.43	1 (8%)
7	POP	H	703	6	6,8,8	0.73	0	12,13,13	1.41	1 (8%)
7	POP	J	703	6	6,8,8	0.73	0	12,13,13	1.41	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	POP	I	703	6	-	2/6/6/6	-
7	POP	L	703	6	-	2/6/6/6	-
7	POP	K	703	6	-	2/6/6/6	-
7	POP	M	703	6	-	2/6/6/6	-
7	POP	H	703	6	-	2/6/6/6	-
7	POP	J	703	6	-	2/6/6/6	-

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	I	703	POP	O6-P2-O5	3.01	119.09	107.80
7	K	703	POP	O6-P2-O5	2.99	119.02	107.80
7	M	703	POP	O6-P2-O5	2.95	118.86	107.80
7	J	703	POP	O6-P2-O5	2.93	118.78	107.80
7	L	703	POP	O6-P2-O5	2.91	118.71	107.80

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	H	703	POP	P2-O-P1-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
7	H	703	POP	P2-O-P1-O1
7	I	703	POP	P2-O-P1-O1
7	J	703	POP	P2-O-P1-O1
7	K	703	POP	P2-O-P1-O1

There are no ring outliers.

6 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	I	703	POP	1	0
7	L	703	POP	1	0
7	K	703	POP	1	0
7	M	703	POP	1	0
7	H	703	POP	1	0
7	J	703	POP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-73864. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 280




Y Index: 280




Z Index: 280


6.2.2 Raw map



X Index: 280



Y Index: 280




Z Index: 280


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 307




Y Index: 309




Z Index: 245


6.3.2 Raw map



X Index: 243



Y Index: 309




Z Index: 316


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.1. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 637 nm³; this corresponds to an approximate mass of 575 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of $0.385 \text{ \AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.385 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.60	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.09	3.59	3.12

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.09 differs from the reported value 2.6 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-73864 and PDB model 9Z6Y. Per-residue inclusion information can be found in section [3](#) on page [9](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.1 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.1).

9.4 Atom inclusion [i](#)



At the recommended contour level, 91% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.1) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8540	 0.5070
A	 0.9150	 0.5230
B	 0.9180	 0.5270
C	 0.9200	 0.5300
D	 0.9200	 0.5290
E	 0.9200	 0.5300
F	 0.9180	 0.5280
H	 0.9700	 0.6400
I	 0.9690	 0.6320
J	 0.9690	 0.6300
K	 0.9690	 0.6310
L	 0.9690	 0.6310
M	 0.9700	 0.6320
R	 0.5780	 0.2390
S	 0.5790	 0.2380
T	 0.5810	 0.2410
U	 0.5790	 0.2420
V	 0.5810	 0.2440
W	 0.5820	 0.2380
X	 0.7020	 0.3850
Y	 0.6330	 0.3890
Z	 0.6880	 0.3850
a	 0.6220	 0.3770
b	 0.7080	 0.3920
c	 0.6220	 0.3700
d	 0.6990	 0.3890
e	 0.6190	 0.3720
f	 0.7050	 0.3850
g	 0.6250	 0.3730
h	 0.6910	 0.3890
i	 0.6250	 0.3720

