



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 04:49 PM UTC

PDB ID : 4Z7V / pdb_00004z7v
Title : L3-12 complex
Authors : Petersen, J.; Rossjohn, J.; Reid, H.H.; Koning, F.
Deposited on : 2015-04-08
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	192	4% 88% 6% 6%
1	C	192	3% 88% 6% 6%
2	B	213	11% 70% 10% 19%
2	D	213	8% 66% 14% 18%
3	E	204	5% 85% 12% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	204	4% 91% 6%
4	F	244	% 93% 6%
4	H	244	% 93% 6%
5	I	18	72% 28%
5	J	18	61% 11% 28%
6	K	6	100%
7	L	4	25% 75%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 13077 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class II HLA-DQ-alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	180	Total	C	N	O	S	0	0	0
			1418	912	236	268	2			
1	C	180	Total	C	N	O	S	0	0	0
			1408	906	232	268	2			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	THR	-	expression tag	UNP Q30069
A	183	SER	-	expression tag	UNP Q30069
A	184	GLY	-	expression tag	UNP Q30069
A	185	ASP	-	expression tag	UNP Q30069
A	186	ASP	-	expression tag	UNP Q30069
A	187	ASP	-	expression tag	UNP Q30069
A	188	ASP	-	expression tag	UNP Q30069
A	189	LYS	-	expression tag	UNP Q30069
C	182	THR	-	expression tag	UNP Q30069
C	183	SER	-	expression tag	UNP Q30069
C	184	GLY	-	expression tag	UNP Q30069
C	185	ASP	-	expression tag	UNP Q30069
C	186	ASP	-	expression tag	UNP Q30069
C	187	ASP	-	expression tag	UNP Q30069
C	188	ASP	-	expression tag	UNP Q30069
C	189	LYS	-	expression tag	UNP Q30069

- Molecule 2 is a protein called MHC class II HLA-DQ-beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	172	Total	C	N	O	S	0	0	0
			1338	857	225	249	7			
2	D	175	Total	C	N	O	S	0	0	0
			1361	870	231	253	7			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-12	GLY	-	expression tag	UNP O19707
B	-11	GLY	-	expression tag	UNP O19707
B	-10	SER	-	expression tag	UNP O19707
B	-9	ILE	-	expression tag	UNP O19707
B	-8	GLU	-	expression tag	UNP O19707
B	-7	GLY	-	expression tag	UNP O19707
B	-6	ARG	-	expression tag	UNP O19707
B	-5	GLY	-	expression tag	UNP O19707
B	-4	GLY	-	expression tag	UNP O19707
B	-3	SER	-	expression tag	UNP O19707
B	-2	GLY	-	expression tag	UNP O19707
B	-1	ALA	-	expression tag	UNP O19707
B	0	SER	-	expression tag	UNP O19707
B	193	THR	-	expression tag	UNP O19707
B	194	GLY	-	expression tag	UNP O19707
B	195	GLY	-	expression tag	UNP O19707
B	196	ASP	-	expression tag	UNP O19707
B	197	ASP	-	expression tag	UNP O19707
B	198	ASP	-	expression tag	UNP O19707
B	199	ASP	-	expression tag	UNP O19707
B	200	LYS	-	expression tag	UNP O19707
D	-12	GLY	-	expression tag	UNP O19707
D	-11	GLY	-	expression tag	UNP O19707
D	-10	SER	-	expression tag	UNP O19707
D	-9	ILE	-	expression tag	UNP O19707
D	-8	GLU	-	expression tag	UNP O19707
D	-7	GLY	-	expression tag	UNP O19707
D	-6	ARG	-	expression tag	UNP O19707
D	-5	GLY	-	expression tag	UNP O19707
D	-4	GLY	-	expression tag	UNP O19707
D	-3	SER	-	expression tag	UNP O19707
D	-2	GLY	-	expression tag	UNP O19707
D	-1	ALA	-	expression tag	UNP O19707
D	0	SER	-	expression tag	UNP O19707
D	193	THR	-	expression tag	UNP O19707
D	194	GLY	-	expression tag	UNP O19707
D	195	GLY	-	expression tag	UNP O19707
D	196	ASP	-	expression tag	UNP O19707
D	197	ASP	-	expression tag	UNP O19707
D	198	ASP	-	expression tag	UNP O19707
D	199	ASP	-	expression tag	UNP O19707
D	200	LYS	-	expression tag	UNP O19707

- Molecule 3 is a protein called T-CELL RECEPTOR, L3-12 ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	199	Total	C	N	O	S	0	0	0
			1534	954	268	303	9			
3	G	198	Total	C	N	O	S	0	0	0
			1524	949	266	300	9			


- Molecule 4 is a protein called T-CELL RECEPTOR, L3-12 BETA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	243	Total	C	N	O	S	0	0	0
			1916	1206	332	373	5			
4	H	242	Total	C	N	O	S	0	0	0
			1907	1202	332	368	5			

- Molecule 5 is a protein called deamidated DQ8-glia-alpha1 peptide.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	I	13	Total	C	N	O	0	0	0
			95	56	16	23			
5	J	13	Total	C	N	O	0	0	0
			95	56	16	23			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	K	6	Total	C	N	O	0	0	0
			70	40	2	28			

- Molecule 7 is an oligosaccharide called alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	L	4	Total	C	N	O	0	0	0
			48	28	2	18			

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	12	Total	O	0	0
			12	12		
9	B	7	Total	O	0	0
			7	7		
9	C	17	Total	O	0	0
			17	17		
9	D	10	Total	O	0	0
			10	10		
9	E	63	Total	O	0	0
			63	63		
9	F	79	Total	O	0	0
			79	79		

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	G	59	Total 59	O 59	0	0
9	H	84	Total 84	O 84	0	0
9	I	2	Total 2	O 2	0	0
9	J	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class II HLA-DQ-alpha chain




- Molecule 1: MHC class II HLA-DQ-alpha chain




- Molecule 2: MHC class II HLA-DQ-beta-1




- Molecule 2: MHC class II HLA-DQ-beta-1





- Molecule 3: T-CELL RECEPTOR, L3-12 ALPHA CHAIN



- Molecule 3: T-CELL RECEPTOR, L3-12 ALPHA CHAIN



- Molecule 4: T-CELL RECEPTOR, L3-12 BETA CHAIN



- Molecule 4: T-CELL RECEPTOR, L3-12 BETA CHAIN



- Molecule 5: deamidated DQ8-glia-alpha1 peptide



- Molecule 5: deamidated DQ8-glia-alpha1 peptide



- Molecule 6: alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 7: α -L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)][α -L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  25% 75%

MAG1
FUC2
MAG3
FUC4

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	56.51Å 76.96Å 132.28Å 93.87° 89.47° 105.09°	Depositor
Resolution (Å)	74.14 – 2.65 74.14 – 2.65	Depositor EDS
% Data completeness (in resolution range)	98.3 (74.14-2.65) 98.3 (74.14-2.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 2.65Å)	Xtriage
Refinement program	BUSTER 2.10.1	Depositor
R, R_{free}	0.187 , 0.226 0.198 , 0.235	Depositor DCC
R_{free} test set	3114 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	46.2	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 69.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13077	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, MAN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.72	0/1462	1.12	0/2002
1	C	0.73	0/1451	1.11	0/1987
2	B	0.76	0/1371	1.09	6/1878 (0.3%)
2	D	0.85	1/1396 (0.1%)	1.11	7/1913 (0.4%)
3	E	0.72	0/1567	1.13	3/2134 (0.1%)
3	G	0.73	0/1557	1.11	5/2122 (0.2%)
4	F	0.68	0/1967	1.05	1/2681 (0.0%)
4	H	0.67	0/1958	1.04	0/2669
5	I	0.48	0/97	1.22	0/130
5	J	0.52	0/97	1.29	2/130 (1.5%)
All	All	0.73	1/12923 (0.0%)	1.09	24/17646 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	165	PRO	N-CD	5.33	1.55	1.47

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	76	PHE	CA-CB-CG	6.86	120.66	113.80
3	G	68	ASN	CA-CB-CG	6.55	119.16	112.60
2	D	22	GLU	N-CA-C	6.37	117.89	111.07
3	E	68	ASN	CA-CB-CG	6.18	118.78	112.60
3	E	179	ASP	CA-CB-CG	5.63	118.23	112.60
3	E	83	ASP	CA-CB-CG	5.46	118.06	112.60
2	B	41	ASP	CA-CB-CG	5.35	117.95	112.60
3	G	179	ASP	CA-CB-CG	5.33	117.93	112.60
3	G	83	ASP	CA-CB-CG	5.32	117.92	112.60
5	J	1	SER	CA-C-N	5.24	124.95	119.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	J	1	SER	C-N-CA	5.24	124.95	119.92
2	B	124	PRO	CA-C-N	5.21	127.79	120.28
2	B	124	PRO	C-N-CA	5.21	127.79	120.28
2	B	89	THR	CA-C-N	5.19	127.54	120.54
2	B	89	THR	C-N-CA	5.19	127.54	120.54
2	D	124	PRO	CA-C-N	5.17	127.73	120.28
2	D	124	PRO	C-N-CA	5.17	127.73	120.28
2	D	55	PRO	N-CA-C	5.16	116.99	110.70
2	B	55	PRO	N-CA-C	5.11	116.93	110.70
2	D	41	ASP	CA-CB-CG	5.10	117.70	112.60
2	D	164	THR	CA-C-N	-5.04	113.54	119.84
2	D	164	THR	C-N-CA	-5.04	113.54	119.84
3	G	130	ILE	CA-C-N	5.01	126.96	120.44
3	G	130	ILE	C-N-CA	5.01	126.96	120.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1418	0	1311	5	0
1	C	1408	0	1297	5	0
2	B	1338	0	1228	13	0
2	D	1361	0	1243	22	0
3	E	1534	0	1434	10	0
3	G	1524	0	1426	4	0
4	F	1916	0	1795	5	0
4	H	1907	0	1787	6	0
5	I	95	0	80	0	0
5	J	95	0	80	1	0
6	K	70	0	61	0	0
7	L	48	0	43	8	0
8	A	14	0	13	0	0
8	C	14	0	13	0	0
9	A	12	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	B	7	0	0	0	0
9	C	17	0	0	0	0
9	D	10	0	0	0	0
9	E	63	0	0	1	0
9	F	79	0	0	0	0
9	G	59	0	0	0	0
9	H	84	0	0	0	0
9	I	2	0	0	0	0
9	J	2	0	0	0	0
All	All	13077	0	11811	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:35:GLU:OE1	2:B:51:THR:HG21	1.43	1.14
2:D:166:GLN:O	2:D:169:ASP:OD1	1.85	0.95
2:D:35:GLU:OE2	2:D:51:THR:OG1	1.89	0.88
7:L:1:NAG:O4	7:L:3:NAG:O7	1.96	0.84
2:D:19:ASN:O	7:L:4:FUC:O4	1.96	0.82
2:D:162:GLU:HB2	2:D:163:MET:HA	1.62	0.80
7:L:2:FUC:H3	7:L:3:NAG:H61	1.70	0.74
3:G:178:LEU:HB3	4:H:184:CYS:HB3	1.70	0.72
2:D:35:GLU:OE2	2:D:51:THR:CB	2.40	0.69
2:B:35:GLU:OE1	2:B:51:THR:CG2	2.32	0.67
2:B:36:GLU:O	2:B:50:VAL:HG23	1.97	0.65
1:C:70:LEU:HD13	2:D:9:TYR:HB2	1.82	0.62
1:C:118:ASN:HB2	1:C:166:GLU:HB2	1.82	0.61
1:A:118:ASN:HB2	1:A:166:GLU:HB2	1.82	0.60
7:L:2:FUC:H5	7:L:3:NAG:O5	2.01	0.60
2:D:163:MET:HG3	2:D:171:TYR:CZ	2.36	0.60
4:H:55:GLN:OE1	4:H:110:THR:HG22	2.00	0.59
2:D:99:VAL:HG11	2:D:175:VAL:HG21	1.85	0.59
2:B:70:ARG:HD3	4:H:113:GLU:OE1	2.03	0.58
3:E:178:LEU:HB3	4:F:184:CYS:HB2	1.85	0.57
2:D:21:THR:OG1	7:L:4:FUC:H3	2.04	0.56
2:D:116:VAL:HG12	2:D:160:MET:HG3	1.88	0.56
1:A:70:LEU:HD13	2:B:9:TYR:HB2	1.88	0.55
2:D:162:GLU:HB2	2:D:163:MET:CA	2.35	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:26:LEU:HG	2:B:42:SER:HB3	1.89	0.54
2:D:26:LEU:HG	2:D:42:SER:HB3	1.90	0.54
2:D:20:GLY:HA3	7:L:4:FUC:O4	2.08	0.53
2:B:35:GLU:O	2:B:35:GLU:HG3	2.08	0.53
7:L:2:FUC:H3	7:L:3:NAG:C6	2.36	0.53
3:E:207:ASN:HA	3:E:210:ILE:HD11	1.91	0.52
1:A:8:SER:C	1:A:9(A):GLY:HA2	2.33	0.52
2:D:162:GLU:CB	2:D:163:MET:HA	2.31	0.52
1:C:8:SER:C	1:C:9(A):GLY:HA2	2.34	0.52
1:A:43:TRP:CE3	1:A:48:PHE:HB3	2.46	0.51
1:C:43:TRP:CE3	1:C:48:PHE:HB3	2.46	0.51
3:E:94:ALA:HB3	9:E:301:HOH:O	2.12	0.50
7:L:4:FUC:O3	7:L:4:FUC:H63	2.11	0.50
2:D:165:PRO:HG2	2:D:165:PRO:O	2.09	0.50
2:D:19:ASN:ND2	2:D:22:GLU:OE1	2.45	0.49
2:B:32:TYR:O	2:B:33:ASN:HB2	2.13	0.49
2:B:172:THR:HG22	2:B:187:GLU:HG2	1.93	0.49
2:D:70:ARG:HD3	4:F:113:GLU:OE1	2.12	0.49
2:D:172:THR:HG22	2:D:187:GLU:HG2	1.93	0.48
3:G:109:SER:O	4:H:112:GLY:HA3	2.12	0.48
3:E:165:SER:HB2	3:E:172:ILE:HD12	1.97	0.47
2:B:70:ARG:HD2	5:J:7:GLN:NE2	2.29	0.47
2:D:10:GLN:HB2	2:D:31:ILE:HB	1.96	0.46
2:D:166:GLN:C	2:D:169:ASP:OD1	2.56	0.46
4:H:40:TYR:HB2	4:H:105:ALA:HB3	1.98	0.45
2:B:37:TYR:HA	2:B:51:THR:HB	1.97	0.45
3:E:206:ASN:HA	3:E:207:ASN:HA	1.61	0.44
3:E:210:ILE:H	3:E:210:ILE:HG13	1.54	0.44
2:D:148:ILE:HB	2:D:156:GLN:HG3	1.99	0.43
1:C:50:ARG:HA	3:G:209:ILE:HD11	2.00	0.43
3:E:40:HIS:CD2	3:E:107:ARG:HB3	2.54	0.42
3:E:183:MET:HE1	4:F:210:SER:HB3	2.01	0.42
2:D:145:THR:HG23	2:D:160:MET:HE1	2.01	0.42
3:G:165:SER:HB2	3:G:172:ILE:HD12	2.02	0.42
3:E:39:ILE:O	3:E:55:HIS:HA	2.19	0.41
4:H:136:PRO:HB3	4:H:163:PHE:CD1	2.55	0.41
3:E:128:PRO:HG3	3:E:177:VAL:HG21	2.01	0.41
1:A:66:LEU:HG	2:B:9:TYR:CD1	2.57	0.40
4:F:63:GLY:HA2	4:F:80:GLN:HB3	2.03	0.40
2:B:30:TYR:CD1	2:B:30:TYR:N	2.89	0.40
4:F:25:PRO:HD3	4:F:86:HIS:HA	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/192 (93%)	173 (97%)	5 (3%)	0	100	100
1	C	178/192 (93%)	173 (97%)	5 (3%)	0	100	100
2	B	164/213 (77%)	156 (95%)	8 (5%)	0	100	100
2	D	169/213 (79%)	160 (95%)	9 (5%)	0	100	100
3	E	197/204 (97%)	189 (96%)	5 (2%)	3 (2%)	8	13
3	G	194/204 (95%)	185 (95%)	9 (5%)	0	100	100
4	F	241/244 (99%)	234 (97%)	7 (3%)	0	100	100
4	H	240/244 (98%)	234 (98%)	6 (2%)	0	100	100
5	I	11/18 (61%)	11 (100%)	0	0	100	100
5	J	11/18 (61%)	11 (100%)	0	0	100	100
All	All	1583/1742 (91%)	1526 (96%)	54 (3%)	3 (0%)	43	60

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	E	64	SER
3	E	145	SER
3	E	209	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/176 (88%)	151 (98%)	3 (2%)	50	71
1	C	152/176 (86%)	149 (98%)	3 (2%)	48	69
2	B	138/189 (73%)	132 (96%)	6 (4%)	26	44
2	D	139/189 (74%)	132 (95%)	7 (5%)	22	37
3	E	168/185 (91%)	162 (96%)	6 (4%)	31	51
3	G	168/185 (91%)	164 (98%)	4 (2%)	43	65
4	F	205/211 (97%)	198 (97%)	7 (3%)	32	53
4	H	203/211 (96%)	195 (96%)	8 (4%)	28	48
5	I	11/14 (79%)	11 (100%)	0	100	100
5	J	11/14 (79%)	11 (100%)	0	100	100
All	All	1349/1550 (87%)	1305 (97%)	44 (3%)	33	55

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ARG
1	A	154	LEU
1	A	175	LEU
2	B	22	GLU
2	B	50	VAL
2	B	68	LEU
2	B	99	VAL
2	B	161	LEU
2	B	185	ILE
1	C	151	LEU
1	C	156	SER
1	C	175	LEU
2	D	21	THR
2	D	68	LEU
2	D	160	MET
2	D	161	LEU
2	D	163	MET
2	D	165	PRO
2	D	185	ILE
3	E	17	GLU
3	E	19	VAL
3	E	150	VAL
3	E	159	GLN
3	E	176	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	E	210	ILE
4	F	13	THR
4	F	50	LEU
4	F	66	ARG
4	F	85	LEU
4	F	168	VAL
4	F	177	LYS
4	F	232	GLU
3	G	17	GLU
3	G	19	VAL
3	G	212	GLU
3	G	218	SER
4	H	13	THR
4	H	50	LEU
4	H	66	ARG
4	H	159	LEU
4	H	168	VAL
4	H	177	LYS
4	H	184	CYS
4	H	232	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (26) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	GLN
1	A	44	GLN
1	A	57	GLN
1	A	62	ASN
1	A	143	HIS
1	C	14	GLN
1	C	44	GLN
1	C	57	GLN
1	C	62	ASN
1	C	111	ASN
3	E	40	HIS
3	E	65	ASN
3	E	114	GLN
3	E	129	ASN
3	E	131	GLN
3	E	203	ASN
4	F	55	GLN
4	F	116	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	F	226	GLN
3	G	14	ASN
3	G	65	ASN
3	G	114	GLN
3	G	129	ASN
3	G	207	ASN
4	H	116	GLN
4	H	238	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	NAG	K	1	6,2	14,14,15	0.30	0	17,19,21	0.82	1 (5%)
6	NAG	K	2	6	14,14,15	0.32	0	17,19,21	0.94	2 (11%)
6	MAN	K	3	6	11,11,12	0.38	0	15,15,17	1.33	1 (6%)
6	MAN	K	4	6	11,11,12	0.34	0	15,15,17	0.97	1 (6%)
6	FUC	K	5	6	10,10,11	0.41	0	14,14,16	0.80	1 (7%)
6	FUC	K	6	6	10,10,11	0.50	0	14,14,16	1.20	1 (7%)
7	NAG	L	1	2,7	14,14,15	1.19	1 (7%)	17,19,21	1.89	4 (23%)
7	FUC	L	2	7	10,10,11	0.41	0	14,14,16	0.73	0
7	NAG	L	3	7	14,14,15	0.75	0	17,19,21	1.62	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	FUC	L	4	7	10,10,11	0.38	0	14,14,16	2.04	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	K	1	6,2	-	2/6/23/26	0/1/1/1
6	NAG	K	2	6	-	2/6/23/26	0/1/1/1
6	MAN	K	3	6	-	0/2/19/22	1/1/1/1
6	MAN	K	4	6	-	0/2/19/22	1/1/1/1
6	FUC	K	5	6	-	-	0/1/1/1
6	FUC	K	6	6	-	-	0/1/1/1
7	NAG	L	1	2,7	-	2/6/23/26	0/1/1/1
7	FUC	L	2	7	-	-	0/1/1/1
7	NAG	L	3	7	-	4/6/23/26	0/1/1/1
7	FUC	L	4	7	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	L	1	NAG	C1-C2	3.56	1.57	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	L	3	NAG	C2-N2-C7	-5.39	115.67	122.90
7	L	1	NAG	C1-O5-C5	5.27	119.25	112.19
6	K	3	MAN	C1-O5-C5	4.69	118.47	112.19
7	L	4	FUC	C1-C2-C3	3.85	115.24	109.64
6	K	4	MAN	C1-O5-C5	3.45	116.81	112.19
6	K	6	FUC	C1-O5-C5	3.39	120.95	112.97
7	L	4	FUC	O5-C5-C4	3.03	115.02	109.55
7	L	4	FUC	O5-C1-C2	-2.94	103.77	110.79
7	L	1	NAG	C6-C5-C4	-2.90	105.89	113.02
7	L	4	FUC	C1-O5-C5	2.81	119.59	112.97
6	K	2	NAG	O5-C1-C2	-2.65	107.19	111.29
7	L	1	NAG	C2-N2-C7	-2.47	119.59	122.90
6	K	5	FUC	C1-O5-C5	2.47	118.79	112.97
7	L	3	NAG	O5-C1-C2	-2.41	107.56	111.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	L	4	FUC	C2-C3-C4	2.40	115.08	110.86
6	K	1	NAG	C1-O5-C5	2.28	115.24	112.19
7	L	1	NAG	O3-C3-C2	-2.24	104.76	109.40
7	L	4	FUC	C3-C4-C5	-2.20	106.48	109.81
6	K	2	NAG	O4-C4-C5	-2.01	104.36	109.32

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	L	3	NAG	C8-C7-N2-C2
7	L	3	NAG	O7-C7-N2-C2
6	K	1	NAG	C4-C5-C6-O6
6	K	1	NAG	O5-C5-C6-O6
7	L	3	NAG	O5-C5-C6-O6
7	L	3	NAG	C4-C5-C6-O6
7	L	1	NAG	O5-C5-C6-O6
6	K	2	NAG	O5-C5-C6-O6
7	L	1	NAG	C4-C5-C6-O6
6	K	2	NAG	C4-C5-C6-O6


All (2) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	K	4	MAN	C1-C2-C3-C4-C5-O5
6	K	3	MAN	C1-C2-C3-C4-C5-O5

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	L	2	FUC	3	0
7	L	3	NAG	4	0
7	L	1	NAG	1	0
7	L	4	FUC	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	NAG	C	1000	1	14,14,15	0.31	0	17,19,21	0.75	1 (5%)
8	NAG	A	1000	1	14,14,15	0.29	0	17,19,21	0.84	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	C	1000	1	-	0/6/23/26	0/1/1/1
8	NAG	A	1000	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
8	A	1000	NAG	C1-O5-C5	2.74	115.86	112.19
8	C	1000	NAG	C1-O5-C5	2.38	115.37	112.19

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	1000	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	180/192 (93%)	0.58	8 (4%) 39 31	36, 79, 113, 127	0
1	C	180/192 (93%)	0.49	6 (3%) 49 40	40, 79, 118, 138	0
2	B	172/213 (80%)	0.80	23 (13%) 7 6	42, 82, 148, 164	0
2	D	175/213 (82%)	0.79	17 (9%) 13 10	41, 76, 148, 176	0
3	E	199/204 (97%)	0.14	10 (5%) 34 26	26, 44, 85, 100	0
3	G	198/204 (97%)	0.07	9 (4%) 38 31	28, 44, 80, 99	0
4	F	243/244 (99%)	-0.21	2 (0%) 82 79	27, 41, 65, 80	0
4	H	242/244 (99%)	-0.20	2 (0%) 82 79	29, 42, 61, 83	0
5	I	13/18 (72%)	0.15	0 100 100	40, 45, 70, 73	0
5	J	13/18 (72%)	0.22	0 100 100	44, 48, 76, 85	0
All	All	1615/1742 (92%)	0.26	77 (4%) 35 28	26, 54, 115, 176	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	143	SER	5.6
3	E	207	ASN	5.5
2	D	136	GLN	4.8
2	D	104	SER	4.8
2	D	115	LEU	4.6
2	B	136	GLN	4.3
2	B	137	GLU	4.2
3	E	209	ILE	3.8
2	B	139	THR	3.5
3	E	198	ASP	3.4
3	E	208	SER	3.4
2	B	184	ILE	3.2
2	D	184	ILE	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	115	LEU	3.1
3	G	209	ILE	3.1
2	B	104	SER	3.1
2	D	102	SER	3.0
1	C	180	PRO	3.0
2	D	188	TRP	3.0
4	F	197	ASN	3.0
1	A	160	ILE	3.0
1	C	160	ILE	3.0
2	D	161	LEU	2.9
1	C	97	VAL	2.8
1	C	174	LEU	2.8
1	A	159	GLU	2.8
4	H	27	SER	2.7
2	B	132	PHE	2.7
2	D	186	VAL	2.7
3	G	219	PRO	2.7
2	D	132	PHE	2.7
4	H	256	ALA	2.7
2	B	186	VAL	2.6
2	D	139	THR	2.6
2	B	163	MET	2.6
2	B	150	ASN	2.5
3	G	145	SER	2.5
2	B	185	ILE	2.5
2	B	130	ARG	2.5
3	E	210	ILE	2.5
1	A	2	VAL	2.5
3	G	47	SER	2.5
3	G	144	LYS	2.4
2	B	189	ARG	2.4
2	B	188	TRP	2.4
3	G	142	ASP	2.4
2	B	140	THR	2.4
2	D	164	THR	2.4
3	E	204	ALA	2.4
3	E	206	ASN	2.3
3	G	195	ASN	2.3
1	A	180	PRO	2.3
1	A	172	GLU	2.3
1	A	128	VAL	2.3
2	B	116	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	175	VAL	2.3
2	B	141	GLY	2.2
1	C	2	VAL	2.2
2	B	161	LEU	2.2
1	C	128	VAL	2.2
1	A	72	ILE	2.2
3	E	205	PHE	2.2
2	B	181	GLN	2.2
3	G	148	LYS	2.2
2	D	116	VAL	2.2
3	E	203	ASN	2.1
2	D	96	GLU	2.1
2	B	114	LEU	2.1
4	F	180	HIS	2.1
2	B	170	VAL	2.1
3	E	197	SER	2.0
2	D	185	ILE	2.0
2	B	171	TYR	2.0
2	D	170	VAL	2.0
2	D	103	PRO	2.0
1	A	99	LEU	2.0
2	D	142	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	FUC	K	6	10/11	0.38	0.15	120,124,127,128	0
6	MAN	K	3	11/12	0.41	0.18	141,144,149,149	0
7	NAG	L	3	14/15	0.60	0.14	114,117,123,124	0
7	FUC	L	4	10/11	0.63	0.12	99,102,104,106	0
6	NAG	K	2	14/15	0.69	0.13	124,131,137,138	0
7	FUC	L	2	10/11	0.69	0.13	107,110,115,115	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	MAN	K	4	11/12	0.71	0.16	145,150,157,159	0
6	FUC	K	5	10/11	0.71	0.13	127,131,136,137	0
6	NAG	K	1	14/15	0.85	0.10	114,117,124,125	0
7	NAG	L	1	14/15	0.86	0.11	95,100,105,110	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	NAG	C	1000	14/15	0.76	0.12	114,118,122,122	0
8	NAG	A	1000	14/15	0.78	0.12	113,117,123,123	0

6.5 Other polymers [i](#)

There are no such residues in this entry.