



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:19 PM UTC

PDB ID : 7Z74 / pdb_00007z74
Title : PI3KC2a core in complex with PITCOIN2
Authors : Lo, W.T.; Roske, Y.; Daumke, O.; Haucke, V.
Deposited on : 2022-03-15
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	910	5% 76% 14% 10%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6615 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol 4-phosphate 3-kinase C2 domain-containing subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	822	Total	C	N	O	S	0	2	0
			6485	4170	1079	1196	40			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	PHE	engineered mutation	UNP Q61194
A	2	ALA	GLU	engineered mutation	UNP Q61194
A	267	GLY	-	linker	UNP Q61194
A	268	SER	-	linker	UNP Q61194
A	269	GLY	-	linker	UNP Q61194
A	270	SER	-	linker	UNP Q61194
A	276	SER	-	linker	UNP Q61194
A	277	ALA	-	linker	UNP Q61194
A	278	GLY	-	linker	UNP Q61194
A	279	ALA	-	linker	UNP Q61194
A	280	GLY	-	linker	UNP Q61194
A	281	SER	-	linker	UNP Q61194
A	282	GLY	-	linker	UNP Q61194
A	283	ALA	-	linker	UNP Q61194
A	286	GLY	ALA	conflict	UNP Q61194
A	353	ALA	PHE	conflict	UNP Q61194
A	354	ALA	PHE	conflict	UNP Q61194
A	427	ALA	LEU	engineered mutation	UNP Q61194

- Molecule 2 is {N}-[4-(3-hydroxyphenyl)-1,3-thiazol-2-yl]-2-[4-oxidanylidene-3-(2-phenylethyl)pteridin-2-yl]sulfanyl-ethanamide (CCD ID: IKC) (formula: C₂₅H₂₀N₆O₃S₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			36	25	6	3	2		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	74	Total	O	0	0
			74	74		

- Molecule 1: Phosphatidylinositol 4-phosphate 3-kinase C2 domain-containing subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.31Å 135.15Å 151.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.26 – 2.50 43.26 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.26-2.50) 99.7 (43.26-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.01 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.226 , 0.270 (Not available) , 0.250	Depositor DCC
R_{free} test set	2049 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	69.5	Xtriage
Anisotropy	0.326	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6615	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: IKC, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	2/6622 (0.0%)	0.71	8/8974 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	839[A]	SER	C-O	10.04	1.36	1.23
1	A	839[B]	SER	C-O	10.04	1.36	1.23

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	839[A]	SER	CA-C-O	9.88	132.03	121.55
1	A	839[B]	SER	CA-C-O	9.88	132.03	121.55
1	A	839[A]	SER	O-C-N	-6.56	114.74	122.81
1	A	839[B]	SER	O-C-N	-6.56	114.74	122.81
1	A	839[A]	SER	N-CA-C	6.29	118.90	110.35
1	A	839[B]	SER	N-CA-C	6.29	118.90	110.35
1	A	791	ARG	N-CA-C	5.56	118.71	110.48
1	A	919	GLY	N-CA-C	-5.17	108.49	115.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6485	0	6480	87	0
2	A	36	0	0	0	0
3	A	20	0	30	0	0
4	A	74	0	0	0	0
All	All	6615	0	6510	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:812:ARG:HG2	1:A:871:ASN:HB2	1.40	1.03
1:A:873:ASN:ND2	1:A:886:ASP:HB2	1.91	0.84
1:A:910:THR:H	1:A:913:MET:HE2	1.48	0.78
1:A:619:LEU:HD21	1:A:655:VAL:HG22	1.69	0.75
1:A:714:LYS:HG3	1:A:791:ARG:NH1	2.04	0.73
1:A:668:MET:O	1:A:672:GLN:HG3	1.94	0.68
1:A:980:THR:HB	1:A:984:GLU:HG3	1.77	0.67
1:A:96:SER:HA	1:A:137:LEU:HD13	1.77	0.66
1:A:714:LYS:HG3	1:A:791:ARG:HH12	1.59	0.66
1:A:981:THR:HG22	1:A:984:GLU:HG2	1.77	0.66
1:A:72:SER:HB2	1:A:76:ILE:HD11	1.78	0.65
1:A:19:LYS:HG2	1:A:655:VAL:HG23	1.79	0.64
1:A:695:LYS:O	1:A:698:GLU:HG2	1.99	0.61
1:A:714:LYS:HE2	1:A:791:ARG:NH1	2.15	0.61
1:A:679:GLU:O	1:A:683:GLN:HG3	2.03	0.59
1:A:671:VAL:HG21	1:A:795:THR:HB	1.83	0.58
1:A:301:THR:HA	1:A:368:GLN:HE21	1.69	0.57
1:A:983:ALA:O	1:A:987:ILE:HG12	2.04	0.57
1:A:671:VAL:HG21	1:A:795:THR:CB	2.35	0.57
1:A:753:VAL:HG21	1:A:801:MET:HE3	1.88	0.56
1:A:140:MET:HE3	1:A:140:MET:HA	1.87	0.56
1:A:781:GLU:HG3	1:A:940:LEU:HD22	1.88	0.56
1:A:753:VAL:HG11	1:A:801:MET:HE3	1.88	0.56
1:A:909:LEU:HA	1:A:913:MET:CE	2.36	0.55
1:A:512:PHE:HE1	1:A:516:ASN:CG	2.14	0.54
1:A:11:PHE:O	1:A:15:ILE:HG12	2.07	0.54
1:A:497:ILE:HD13	1:A:523:HIS:CD2	2.43	0.54
1:A:673:LEU:O	1:A:677:VAL:HG23	2.07	0.54
1:A:989:PHE:HA	1:A:992:LEU:HD12	1.90	0.54
1:A:431:THR:OG1	1:A:450:GLU:HB2	2.09	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:677:VAL:HG13	1:A:696:SER:HB2	1.90	0.53
1:A:912:ASP:N	1:A:912:ASP:OD1	2.40	0.53
1:A:512:PHE:O	1:A:512:PHE:CD1	2.62	0.52
1:A:669:LYS:O	1:A:673:LEU:HG	2.10	0.52
1:A:499:ARG:NH2	1:A:503:PHE:O	2.38	0.52
1:A:870:HIS:CE1	1:A:872:ASP:HB2	2.45	0.52
1:A:730:PHE:HB2	1:A:736:PRO:HD2	1.93	0.51
1:A:753:VAL:HG11	1:A:801:MET:CE	2.41	0.51
1:A:566:ALA:HB2	1:A:602:TYR:CE1	2.46	0.51
1:A:766:LEU:HD23	1:A:890:PHE:HE2	1.76	0.50
1:A:873:ASN:HD22	1:A:886:ASP:HB2	1.73	0.50
1:A:413:LEU:O	1:A:420:LEU:HD12	2.12	0.50
1:A:876:LEU:HD12	1:A:881:HIS:O	2.12	0.50
1:A:330:SER:OG	1:A:379:HIS:HB2	2.13	0.49
1:A:720:LYS:HG3	1:A:721:GLU:HB3	1.95	0.48
1:A:778:TRP:CE2	1:A:937:ALA:HB1	2.49	0.48
1:A:508:GLU:O	1:A:511:VAL:HG22	2.14	0.48
1:A:671:VAL:HG23	1:A:755:PHE:CE2	2.49	0.47
1:A:713:LEU:HB2	1:A:801:MET:HE2	1.96	0.47
1:A:16:MET:HE2	1:A:155:VAL:H	1.80	0.47
1:A:345:LYS:HG2	1:A:361:GLU:OE1	2.15	0.47
1:A:497:ILE:HD13	1:A:523:HIS:HD2	1.80	0.46
1:A:566:ALA:HB2	1:A:602:TYR:CZ	2.50	0.46
1:A:79:MET:HE3	1:A:94:VAL:HG11	1.96	0.46
1:A:87:ASP:OD1	1:A:87:ASP:N	2.48	0.46
1:A:906:PRO:HG3	1:A:992:LEU:HD22	1.97	0.46
1:A:49:ALA:O	1:A:70:VAL:HG22	2.16	0.46
1:A:608:SER:O	1:A:612:ARG:HG3	2.16	0.46
1:A:508:GLU:H	1:A:508:GLU:CD	2.24	0.46
1:A:38:VAL:HG22	1:A:587:ASP:CG	2.41	0.45
1:A:770:MET:O	1:A:774:MET:HG3	2.16	0.45
1:A:988:PHE:CE2	1:A:992:LEU:HD11	2.52	0.45
1:A:589:LEU:HD22	1:A:624:ILE:HD12	1.97	0.45
1:A:935:CYS:SG	1:A:986:THR:HG23	2.57	0.45
1:A:295:LYS:HB3	1:A:465:ILE:HD13	1.99	0.45
1:A:619:LEU:HD23	1:A:619:LEU:HA	1.79	0.44
1:A:818:TYR:OH	1:A:823:SER:HB2	2.18	0.43
1:A:812:ARG:HG2	1:A:871:ASN:CB	2.29	0.43
1:A:140:MET:HE3	1:A:141:CYS:H	1.84	0.43
1:A:918:ASN:OD1	1:A:927:PHE:HB2	2.19	0.42
1:A:512:PHE:CE1	1:A:516:ASN:CG	2.95	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:382:LEU:HD12	1:A:383:PHE:H	1.83	0.42
1:A:700:VAL:O	1:A:703:PHE:HB3	2.19	0.42
1:A:315:SER:O	1:A:319:VAL:HG23	2.18	0.42
1:A:325:TYR:HB3	1:A:382:LEU:HD11	2.02	0.42
1:A:827:LYS:N	1:A:828:PRO:HD3	2.34	0.42
1:A:79:MET:HA	1:A:82:LEU:HD12	2.02	0.42
1:A:778:TRP:CE3	1:A:785:LEU:HD12	2.55	0.42
1:A:593:LEU:HA	1:A:593:LEU:HD23	1.83	0.41
1:A:971:TYR:CD1	1:A:971:TYR:C	2.98	0.41
1:A:487:SER:C	1:A:489:ILE:H	2.28	0.41
1:A:302:GLU:O	1:A:369:ILE:HG12	2.20	0.41
1:A:989:PHE:O	1:A:993:ILE:HG13	2.21	0.41
1:A:16:MET:O	1:A:20:THR:HG23	2.21	0.41
1:A:111:ASN:OD1	1:A:112:HIS:N	2.54	0.41
1:A:714:LYS:HE2	1:A:791:ARG:HH12	1.85	0.40
1:A:827:LYS:N	1:A:828:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	810/910 (89%)	796 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	715/800 (89%)	715 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	119	HIS
1	A	303	GLN
1	A	352	ASN
1	A	368	GLN
1	A	523	HIS
1	A	694	GLN
1	A	936	GLN
1	A	950	ASN
1	A	967	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	1102	-	3,3,3	0.52	0	2,2,2	0.16	0
2	IKC	A	1101	-	40,40,40	2.12	9 (22%)	50,55,55	2.81	13 (26%)
3	EDO	A	1105	-	3,3,3	0.46	0	2,2,2	0.35	0
3	EDO	A	1106	-	3,3,3	0.41	0	2,2,2	0.38	0
3	EDO	A	1104	-	3,3,3	0.46	0	2,2,2	0.34	0
3	EDO	A	1103	-	3,3,3	0.44	0	2,2,2	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1102	-	-	1/1/1/1	-
2	IKC	A	1101	-	-	9/18/18/18	0/5/5/5
3	EDO	A	1105	-	-	0/1/1/1	-
3	EDO	A	1106	-	-	0/1/1/1	-
3	EDO	A	1104	-	-	0/1/1/1	-
3	EDO	A	1103	-	-	0/1/1/1	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	IKC	C25-N24	5.48	1.46	1.38
2	A	1101	IKC	C28-C27	-5.39	1.39	1.47
2	A	1101	IKC	C02-N24	4.82	1.46	1.37
2	A	1101	IKC	C05-S04	4.82	1.82	1.74
2	A	1101	IKC	C05-N06	3.93	1.35	1.29
2	A	1101	IKC	C27-N26	-3.63	1.32	1.39
2	A	1101	IKC	C13-N15	-3.31	1.32	1.40
2	A	1101	IKC	C35-C27	2.56	1.39	1.36
2	A	1101	IKC	O01-C02	-2.04	1.19	1.23

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	IKC	C25-N26-C27	9.90	117.95	109.96
2	A	1101	IKC	S36-C25-N26	-8.76	106.49	115.69
2	A	1101	IKC	C35-S36-C25	8.16	94.75	88.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	IKC	C27-C35-S36	-4.92	105.20	111.16
2	A	1101	IKC	C03-S04-C05	4.25	107.69	99.04
2	A	1101	IKC	N15-C05-N06	-4.22	120.90	124.93
2	A	1101	IKC	C28-C27-C35	-3.27	122.67	126.54
2	A	1101	IKC	C03-C02-N24	3.24	119.30	114.53
2	A	1101	IKC	C12-C07-N06	-3.08	118.87	123.19
2	A	1101	IKC	N24-C25-N26	3.01	126.37	121.18
2	A	1101	IKC	O14-C13-C12	-2.81	119.55	124.18
2	A	1101	IKC	C13-C12-N11	2.35	121.06	116.23
2	A	1101	IKC	S36-C25-N24	2.16	127.14	123.29

There are no chirality outliers.


All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1101	IKC	N24-C02-C03-S04
2	A	1101	IKC	C02-C03-S04-C05
2	A	1101	IKC	N26-C25-N24-C02
2	A	1101	IKC	S36-C25-N24-C02
2	A	1101	IKC	N26-C27-C28-C29
2	A	1101	IKC	O01-C02-C03-S04
2	A	1101	IKC	N26-C27-C28-C34
2	A	1101	IKC	C35-C27-C28-C29
3	A	1102	EDO	O1-C1-C2-O2
2	A	1101	IKC	C35-C27-C28-C34

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	822/910 (90%)	0.47	46 (5%) 30 26	39, 81, 138, 197	2 (0%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	40	VAL	4.5
1	A	1006	PHE	4.2
1	A	388	GLN	4.0
1	A	3	VAL	3.9
1	A	476	ILE	3.5
1	A	41	GLN	3.5
1	A	292	ARG	3.4
1	A	894	ALA	3.4
1	A	480	ASP	3.2
1	A	489	ILE	3.2
1	A	966	ILE	3.1
1	A	432	SER	2.9
1	A	841	GLU	2.9
1	A	734	ALA	2.9
1	A	521	LEU	2.9
1	A	512	PHE	2.8
1	A	487	SER	2.8
1	A	485	LEU	2.8
1	A	826	ASP	2.7
1	A	520	CYS	2.6
1	A	293	ASN	2.6
1	A	484	THR	2.6
1	A	39	THR	2.6
1	A	92	VAL	2.5
1	A	89	LEU	2.5
1	A	821	THR	2.5
1	A	689	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	526	CYS	2.5
1	A	824	PHE	2.4
1	A	866	ILE	2.4
1	A	523	HIS	2.4
1	A	871	ASN	2.3
1	A	872	ASP	2.3
1	A	729	PHE	2.3
1	A	136	THR	2.2
1	A	429	LEU	2.2
1	A	353	ALA	2.2
1	A	829	LEU	2.2
1	A	819	GLY	2.1
1	A	869	ARG	2.1
1	A	682	ARG	2.1
1	A	735	MET	2.1
1	A	908	VAL	2.1
1	A	137	LEU	2.1
1	A	874	ILE	2.1
1	A	820	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	A	1104	4/4	0.68	0.13	100,103,104,108	0
3	EDO	A	1106	4/4	0.83	0.10	105,107,115,120	0
2	IKC	A	1101	36/36	0.85	0.15	71,98,115,117	0
3	EDO	A	1105	4/4	0.87	0.20	80,81,84,94	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	A	1103	4/4	0.89	0.17	72,73,73,75	0
3	EDO	A	1102	4/4	0.89	0.21	82,83,86,87	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.