



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 06:09 AM UTC

PDB ID : 7Z84 / pdb_00007z84
EMDB ID : EMD-14542
Title : Complex I from E. coli, DDM/LMNG-purified, under Turnover at pH 8, Open-ready state
Authors : Kravchuk, V.; Kampjut, D.; Sazanov, L.
Deposited on : 2022-03-16
Resolution : 2.87 Å (reported)
Based on initial models : 3RKO, 4HEA

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	12062 (2.37 - 3.37)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	F	445	
2	E	166	
3	G	908	
4	C	600	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	B	220	
6	I	180	
7	H	325	
8	A	147	
9	L	613	
10	M	509	
11	N	485	
12	K	100	
13	J	184	

2 Entry composition

There are 20 unique types of molecules in this entry. The entry contains 37772 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH-quinone oxidoreductase subunit F.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	F	439	Total	C	N	O	S	0	0
			3407	2162	596	629	20		

- Molecule 2 is a protein called NADH dehydrogenase I subunit E.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	E	156	Total	C	N	O	S	0	0
			1220	768	215	229	8		

- Molecule 3 is a protein called NADH-quinone oxidoreductase.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	905	Total	C	N	O	S	0	0
			7022	4388	1269	1328	37		

- Molecule 4 is a protein called NADH-quinone oxidoreductase subunit CD.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	C	586	Total	C	N	O	S	0	0
			4738	3037	825	852	24		

- Molecule 5 is a protein called NADH-quinone oxidoreductase subunit B.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	B	208	Total	C	N	O	S	0	0
			1656	1051	288	301	16		

- Molecule 6 is a protein called NADH-quinone oxidoreductase subunit I.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	I	180	Total	C	N	O	S	0	0
			1436	915	242	264	15		

- Molecule 7 is a protein called NADH-quinone oxidoreductase subunit H.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	H	323	Total	C	N	O	S	0	0
			2544	1707	400	419	18		

- Molecule 8 is a protein called NADH-quinone oxidoreductase subunit A.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	A	113	Total	C	N	O	S	0	0
			894	606	143	141	4		

- Molecule 9 is a protein called NADH-quinone oxidoreductase subunit L.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	L	605	Total	C	N	O	S	0	0
			4629	3077	740	780	32		

- Molecule 10 is a protein called NADH dehydrogenase I subunit M.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	M	504	Total	C	N	O	S	0	0
			3953	2661	617	646	29		

- Molecule 11 is a protein called NADH-quinone oxidoreductase subunit N.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	N	470	Total	C	N	O	S	0	0
			3563	2382	563	598	20		


- Molecule 12 is a protein called NADH-quinone oxidoreductase subunit K.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	K	100	Total	C	N	O	S	0	0
			760	494	132	129	5		

- Molecule 13 is a protein called NADH-quinone oxidoreductase subunit J.


Mol	Chain	Residues	Atoms					AltConf	Trace
13	J	162	Total	C	N	O	S	0	0
			1226	824	188	207	7		

- Molecule 14 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			AltConf
14	F	1	Total	Fe	S	0
			8	4	4	
14	G	1	Total	Fe	S	0
			8	4	4	
14	G	1	Total	Fe	S	0
			8	4	4	
14	G	1	Total	Fe	S	0
			8	4	4	
14	B	1	Total	Fe	S	0
			8	4	4	
14	I	1	Total	Fe	S	0
			8	4	4	
14	I	1	Total	Fe	S	0
			8	4	4	

- Molecule 15 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula: $C_{17}H_{21}N_4O_9P$).




Mol	Chain	Residues	Atoms					AltConf
15	F	1	Total	C	N	O	P	0
			31	17	4	9	1	

- Molecule 16 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (CCD ID: NAI) (formula: C₂₁H₂₉N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					AltConf
16	F	1	Total	C	N	O	P	0
			44	21	7	14	2	

- Molecule 17 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			AltConf
17	E	1	Total	Fe	S	0
			4	2	2	
17	G	1	Total	Fe	S	0
			4	2	2	

- Molecule 18 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		AltConf
18	G	1	Total	Ca	0
			1	1	

- Molecule 19 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (CCD ID: 3PE) (formula: C₄₁H₈₂NO₈P).



Mol	Chain	Residues	Atoms					AltConf
19	I	1	Total	C	N	O	P	0
			51	41	1	8	1	
19	H	1	Total	C	N	O	P	0
			51	41	1	8	1	
19	L	1	Total	C	N	O	P	0
			51	41	1	8	1	
19	L	1	Total	C	N	O	P	0
			51	41	1	8	1	
19	L	1	Total	C	N	O	P	0
			51	41	1	8	1	
19	L	1	Total	C	N	O	P	0
			51	41	1	8	1	
19	M	1	Total	C	N	O	P	0
			51	41	1	8	1	
19	M	1	Total	C	N	O	P	0
			51	41	1	8	1	
19	J	1	Total	C	N	O	P	0
			51	41	1	8	1	
19	J	1	Total	C	N	O	P	0
			51	41	1	8	1	

- Molecule 20 is EICOSANE (CCD ID: LFA) (formula: C₂₀H₄₂).




Mol	Chain	Residues	Atoms		AltConf
20	H	1	Total	C	0
			20	20	
20	L	1	Total	C	0
			20	20	
20	M	1	Total	C	0
			20	20	
20	N	1	Total	C	0
			14	14	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: NADH-quinone oxidoreductase subunit F



• Molecule 2: NADH dehydrogenase I subunit E




• Molecule 3: NADH-quinone oxidoreductase






• Molecule 4: NADH-quinone oxidoreductase subunit CD

Chain C: 84% 14%



• Molecule 5: NADH-quinone oxidoreductase subunit B

Chain B: 76% 19% 5%




• Molecule 6: NADH-quinone oxidoreductase subunit I


Chain I: 83% 17%



• Molecule 7: NADH-quinone oxidoreductase subunit H

Chain H: 79% 20%






• Molecule 8: NADH-quinone oxidoreductase subunit A

Chain A: 65% 11% 23%




• Molecule 9: NADH-quinone oxidoreductase subunit L

Chain L: 84% 15%



• Molecule 10: NADH dehydrogenase I subunit M


Chain M: 85% 14%





• Molecule 11: NADH-quinone oxidoreductase subunit N

Chain N: 80% 16%






• Molecule 12: NADH-quinone oxidoreductase subunit K



• Molecule 13: NADH-quinone oxidoreductase subunit J



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	40884	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	80	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.495	Depositor
Minimum map value	0.000	Depositor
Average map value	0.004	Depositor
Map value standard deviation	0.018	Depositor
Recommended contour level	0.045	Depositor
Map size (Å)	156.87999, 211.99998, 240.61998	wwPDB
Map dimensions	148, 200, 227	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.06, 1.06, 1.06	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAI, LFA, 3PE, FMN, SF4, FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	F	0.17	0/3486	0.37	0/4713
2	E	0.16	0/1248	0.38	0/1691
3	G	0.16	0/7173	0.37	2/9726 (0.0%)
4	C	0.18	0/4868	0.39	2/6605 (0.0%)
5	B	0.17	0/1692	0.38	0/2292
6	I	0.15	0/1470	0.36	0/1985
7	H	0.21	0/2620	0.44	0/3565
8	A	0.19	0/921	0.41	0/1251
9	L	0.18	0/4747	0.41	1/6468 (0.0%)
10	M	0.19	0/4074	0.43	2/5546 (0.0%)
11	N	0.20	0/3649	0.46	2/4977 (0.0%)
12	K	0.19	0/769	0.46	1/1040 (0.1%)
13	J	0.18	0/1252	0.45	0/1708
All	All	0.18	0/37969	0.40	10/51567 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	G	0	1
9	L	0	1
11	N	0	2
All	All	0	4

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	268	TYR	CA-C-N	7.05	124.68	120.24
4	C	268	TYR	C-N-CA	7.05	124.68	120.24
11	N	64	THR	N-CA-C	-5.78	97.03	109.81
10	M	136	MET	CB-CG-SD	5.66	129.67	112.70
9	L	515	LYS	CB-CG-CD	5.59	124.17	111.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	G	669	THR	Peptide
9	L	257	THR	Peptide
11	N	63	VAL	Peptide
11	N	64	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	3407	0	3374	41	0
2	E	1220	0	1187	10	0
3	G	7022	0	6824	68	0
4	C	4738	0	4658	53	0
5	B	1656	0	1644	29	0
6	I	1436	0	1415	24	0
7	H	2544	0	2591	50	0
8	A	894	0	904	13	0
9	L	4629	0	4772	53	0
10	M	3953	0	4053	44	0
11	N	3563	0	3741	56	0
12	K	760	0	817	16	0
13	J	1226	0	1297	25	0
14	B	8	0	0	0	0
14	F	8	0	0	0	0
14	G	24	0	0	1	0
14	I	16	0	0	0	0
15	F	31	0	19	2	0
16	F	44	0	27	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
17	E	4	0	0	1	0
17	G	4	0	0	0	0
18	G	1	0	0	0	0
19	H	51	0	82	8	0
19	I	51	0	82	1	0
19	J	102	0	164	5	0
19	L	204	0	328	11	0
19	M	102	0	164	1	0
20	H	20	0	42	2	0
20	L	20	0	42	1	0
20	M	20	0	42	0	0
20	N	14	0	27	0	0
All	All	37772	0	38296	428	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 428 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:N:63:VAL:O	11:N:67:MET:HB2	1.74	0.87
10:M:181:SER:HB2	10:M:230:ALA:HA	1.66	0.78
9:L:223:LEU:HD13	9:L:283:VAL:HG22	1.72	0.71
1:F:95:GLU:HB2	16:F:503:NAI:H42N	1.74	0.70
11:N:65:PRO:HG2	13:J:136:PRO:HB3	1.75	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	437/445 (98%)	432 (99%)	5 (1%)	0	100	100
2	E	154/166 (93%)	150 (97%)	4 (3%)	0	100	100
3	G	903/908 (99%)	877 (97%)	26 (3%)	0	100	100
4	C	582/600 (97%)	571 (98%)	11 (2%)	0	100	100
5	B	202/220 (92%)	196 (97%)	6 (3%)	0	100	100
6	I	178/180 (99%)	174 (98%)	4 (2%)	0	100	100
7	H	321/325 (99%)	313 (98%)	8 (2%)	0	100	100
8	A	109/147 (74%)	109 (100%)	0	0	100	100
9	L	601/613 (98%)	590 (98%)	11 (2%)	0	100	100
10	M	502/509 (99%)	491 (98%)	11 (2%)	0	100	100
11	N	464/485 (96%)	456 (98%)	7 (2%)	1 (0%)	43	70
12	K	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
13	J	160/184 (87%)	160 (100%)	0	0	100	100
All	All	4711/4882 (96%)	4616 (98%)	94 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
11	N	64	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	353/359 (98%)	343 (97%)	10 (3%)	38	70
2	E	129/139 (93%)	126 (98%)	3 (2%)	44	74
3	G	732/735 (100%)	722 (99%)	10 (1%)	59	83
4	C	506/519 (98%)	500 (99%)	6 (1%)	63	84
5	B	181/192 (94%)	177 (98%)	4 (2%)	45	75
6	I	154/154 (100%)	153 (99%)	1 (1%)	78	92

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
7	H	267/269 (99%)	263 (98%)	4 (2%)	57	82
8	A	89/119 (75%)	87 (98%)	2 (2%)	45	75
9	L	481/486 (99%)	473 (98%)	8 (2%)	53	80
10	M	413/418 (99%)	409 (99%)	4 (1%)	68	87
11	N	374/385 (97%)	368 (98%)	6 (2%)	55	81
12	K	79/79 (100%)	76 (96%)	3 (4%)	29	61
13	J	128/146 (88%)	125 (98%)	3 (2%)	44	74
All	All	3886/4000 (97%)	3822 (98%)	64 (2%)	54	81

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
11	N	480	LEU
12	K	46	PHE
4	C	133	ASN
4	C	106	ASP
12	K	51	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
7	H	152	GLN
9	L	409	ASN
7	H	312	ASN
9	L	74	ASN
10	M	206	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 1 is monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
19	3PE	J	201	-	50,50,50	0.30	0	53,55,55	0.27	0
14	SF4	F	501	1	0,12,12	-	-	-		
16	NAI	F	503	-	47,48,48	0.26	0	64,73,73	0.35	0
19	3PE	L	801	-	50,50,50	0.30	0	53,55,55	0.28	0
20	LFA	N	601	-	13,13,19	0.14	0	12,12,18	0.11	0
17	FES	G	1004	3	0,4,4	-	-	-		
19	3PE	M	1202	-	50,50,50	0.30	0	53,55,55	0.29	0
14	SF4	I	202	6	0,12,12	-	-	-		
14	SF4	I	201	6	0,12,12	-	-	-		
14	SF4	G	1001	3	0,12,12	-	-	-		
19	3PE	L	803	-	50,50,50	0.30	0	53,55,55	0.28	0
19	3PE	L	804	-	50,50,50	0.30	0	53,55,55	0.30	0
14	SF4	G	1002	3	0,12,12	-	-	-		
19	3PE	I	203	-	50,50,50	0.30	0	53,55,55	0.28	0
19	3PE	L	805	-	50,50,50	0.30	0	53,55,55	0.29	0
20	LFA	M	1201	-	19,19,19	0.13	0	18,18,18	0.10	0
20	LFA	L	802	-	19,19,19	0.12	0	18,18,18	0.12	0
17	FES	E	201	2	0,4,4	-	-	-		
20	LFA	H	402	-	19,19,19	0.14	0	18,18,18	0.10	0
19	3PE	M	1203	-	50,50,50	0.29	0	53,55,55	0.27	0
19	3PE	J	202	-	50,50,50	0.30	0	53,55,55	0.29	0
14	SF4	G	1003	3	0,12,12	-	-	-		
19	3PE	H	401	-	50,50,50	0.30	0	53,55,55	0.30	0
15	FMN	F	502	-	33,33,33	1.04	2 (6%)	48,50,50	1.26	6 (12%)
14	SF4	B	301	5	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
19	3PE	J	201	-	-	10/54/54/54	-
14	SF4	F	501	1	-	-	0/6/5/5
19	3PE	L	801	-	-	13/54/54/54	-
16	NAI	F	503	-	-	5/29/72/72	0/5/5/5
20	LFA	N	601	-	-	0/11/11/17	-
17	FES	G	1004	3	-	-	0/1/1/1
19	3PE	M	1202	-	-	9/54/54/54	-
14	SF4	I	202	6	-	-	0/6/5/5
14	SF4	I	201	6	-	-	0/6/5/5
14	SF4	G	1001	3	-	-	0/6/5/5
19	3PE	L	803	-	-	4/54/54/54	-
19	3PE	L	804	-	-	15/54/54/54	-
14	SF4	G	1002	3	-	-	0/6/5/5
19	3PE	I	203	-	-	7/54/54/54	-
19	3PE	L	805	-	-	12/54/54/54	-
20	LFA	M	1201	-	-	1/17/17/17	-
20	LFA	L	802	-	-	0/17/17/17	-
17	FES	E	201	2	-	-	0/1/1/1
20	LFA	H	402	-	-	1/17/17/17	-
19	3PE	M	1203	-	-	10/54/54/54	-
19	3PE	J	202	-	-	5/54/54/54	-
19	3PE	H	401	-	-	9/54/54/54	-
14	SF4	G	1003	3	-	-	0/6/5/5
15	FMN	F	502	-	-	7/18/18/18	0/3/3/3
14	SF4	B	301	5	-	-	0/6/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	F	502	FMN	C4A-N5	3.30	1.37	1.30
15	F	502	FMN	C10-N1	2.29	1.37	1.33

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	F	502	FMN	C4-N3-C2	-3.35	119.69	125.64
15	F	502	FMN	C4A-C10-N10	2.87	120.60	116.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	F	502	FMN	C4A-C4-N3	2.64	119.97	113.25
15	F	502	FMN	O4-C4-C4A	-2.44	120.09	126.53
15	F	502	FMN	C10-C4A-N5	-2.42	119.87	124.81

There are no chirality outliers.

5 of 108 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
15	F	502	FMN	N10-C1'-C2'-O2'
15	F	502	FMN	N10-C1'-C2'-C3'
15	F	502	FMN	C5'-O5'-P-O2P
15	F	502	FMN	C5'-O5'-P-O3P
19	I	203	3PE	C1-O11-P-O12


There are no ring outliers.


15 monomers are involved in 35 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
19	J	201	3PE	3	0
16	F	503	NAI	2	0
19	L	801	3PE	1	0
19	M	1202	3PE	1	0
19	L	803	3PE	5	0
19	L	804	3PE	2	0
19	I	203	3PE	1	0
19	L	805	3PE	4	0
20	L	802	LFA	1	0
17	E	201	FES	1	0
20	H	402	LFA	2	0
19	J	202	3PE	5	0
14	G	1003	SF4	1	0
19	H	401	3PE	8	0
15	F	502	FMN	2	0





The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the




average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















Ligand LFA M 1201	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand LFA H 402	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand 3PE M 1203	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand 3PE J 202	
 Bond lengths	 Bond angles
 Torsions	 Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-14542. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 74



Y Index: 100



Z Index: 113

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 76



Y Index: 53




Z Index: 152


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.045. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate [i](#)



The volume at the recommended contour level is 231 nm³; this corresponds to an approximate mass of 209 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.


7.3 Rotationally averaged power spectrum [i](#)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.348 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.87	-	-
Author-provided FSC curve	2.87	3.49	2.93
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-14542 and PDB model 7Z84. Per-residue inclusion information can be found in section 3 on page 11.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.045 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.045).

9.4 Atom inclusion [i](#)



At the recommended contour level, 90% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.045) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8700	 0.5910
A	 0.7920	 0.5480
B	 0.8850	 0.6100
C	 0.9140	 0.6170
E	 0.9180	 0.6080
F	 0.9290	 0.6200
G	 0.9370	 0.6360
H	 0.8330	 0.5560
I	 0.9210	 0.6290
J	 0.7590	 0.5220
K	 0.8390	 0.5590
L	 0.7910	 0.5420
M	 0.8020	 0.5650
N	 0.8620	 0.5700

