Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:15 PM UTC

PDB ID	:	2Z9W / pdb_00002z9w
Title	:	Crystal structure of pyridoxamine-pyruvate aminotransferase complexed with pyridoxal
Authors	:	Yoshikane, Y.; Yokochi, N.; Yamasaki, M.; Mizutani, K.; Ohnishi, K.; Mikami, B.; Hayashi, H.; Yagi, T.
Deposited on	:	2007-09-26
Resolution	:	1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	% 92% 8%
1	B	392	93% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	953	-	-	X	-

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 6837 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate aminotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	10	0
			2991	1900	517	559	15			
1	B	392	Total	C	N	O	S	0	8	0
			2969	1892	510	552	15			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	O	S		0	0
			5	4	1			
2	A	1	Total	O	S		0	0
			5	4	1			
2	A	1	Total	O	S		0	0
			5	4	1			
2	B	1	Total	O	S		0	0
			5	4	1			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is 3-HYDROXY-5-(HYDROXYMETHYL)-2-METHYLISONICOTINALDEHYDE (CCD ID: PXL) (formula: $C_8H_9NO_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 11 8 1 2	0	0
3	B	1	Total C N O 11 8 1 2	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	387	Total O 387 387	0	0
5	B	391	Total O 391 391	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aspartate aminotransferase



• Molecule 1: Aspartate aminotransferase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α, β, γ	68.73Å 68.73Å 311.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.84 – 1.70 14.84 – 1.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (14.84-1.70) 94.8 (14.84-1.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	6.39 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.154 , 0.181 (Not available) , 0.170	Depositor DCC
R_{free} test set	4988 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å^2)	11.6	Xtriage
Anisotropy	0.401	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.42 , 50.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	6837	wwPDB-VP
Average B, all atoms (Å^2)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL, PXL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.62	2/3076 (0.1%)	0.77	0/4192
1	B	0.60	0/3064	0.76	0/4176
All	All	0.61	2/6140 (0.0%)	0.77	0/8368

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	197	LYS	CA-C	8.13	1.56	1.52
1	A	197	LYS	C-O	-5.83	1.17	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2991	0	3004	29	0
1	B	2969	0	2996	21	0
2	A	15	0	0	1	0
2	B	20	0	0	0	0
3	A	11	0	7	0	0
3	B	11	0	8	1	0
4	A	18	0	24	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	24	0	32	1	0
5	A	387	0	0	1	0
5	B	391	0	0	4	0
All	All	6837	0	6071	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:ASP:H	4:A:953:GOL:H32	1.08	1.06
1:A:43:LEU:HG	1:A:47[A]:LYS:HE2	1.44	0.97
1:A:299:ASP:H	4:A:953:GOL:C3	1.83	0.90
1:A:299:ASP:N	4:A:953:GOL:H32	1.93	0.80
1:B:370:ASN:HD21	1:B:376:LEU:H	1.29	0.79
1:A:370:ASN:HD21	1:A:376:LEU:H	1.30	0.78
1:B:47[B]:LYS:HE3	5:B:2893:HOH:O	1.83	0.77
1:B:47[B]:LYS:CE	5:B:2893:HOH:O	2.35	0.71
1:A:338[A]:GLU:O	1:A:342[A]:LYS:HD2	1.97	0.65
1:B:196:ASN:HD22	1:B:196:ASN:H	1.45	0.63
1:A:229:MET:HE1	1:B:70:VAL:HG23	1.79	0.63
1:A:144:HIS:HE1	1:A:149:GLY:HA2	1.63	0.62
1:B:53:GLN:NE2	5:B:2838:HOH:O	2.32	0.62
1:A:298:SER:HA	4:A:953:GOL:H31	1.82	0.61
1:A:70:VAL:HG23	1:B:229:MET:HE1	1.87	0.56
1:B:196:ASN:H	1:B:196:ASN:ND2	2.03	0.56
1:B:370:ASN:ND2	1:B:376:LEU:H	2.02	0.56
1:A:338[A]:GLU:OE2	1:A:342[A]:LYS:HE2	2.08	0.54
1:A:27[A]:ARG:NH1	2:A:1605:SO4:O4	2.42	0.53
1:B:144:HIS:HE1	1:B:149:GLY:HA2	1.74	0.52
1:A:98:GLY:C	4:A:955:GOL:H11	2.35	0.51
1:A:370:ASN:ND2	1:A:376:LEU:H	2.05	0.51
1:A:109:HIS:HE1	5:A:1831:HOH:O	1.95	0.50
1:B:167:TYR:CZ	1:B:217:LYS:HD2	2.47	0.48
1:A:44:LEU:HA	1:A:47[A]:LYS:HE3	1.94	0.48
1:A:20[B]:ASN:HB3	5:B:2927:HOH:O	2.13	0.47
1:A:195:PRO:HB2	1:A:201:ALA:HB3	1.96	0.46
1:B:197:LYS:NZ	3:B:1197:PXL:O3	2.48	0.46
1:A:102:TRP:CE2	4:A:955:GOL:H32	2.51	0.45
1:B:195:PRO:HB2	1:B:201:ALA:HB3	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:THR:HG21	1:B:35:TYR:CG	2.51	0.45
1:A:370:ASN:HD21	1:A:376:LEU:HB2	1.83	0.44
1:A:98:GLY:O	4:A:955:GOL:H31	2.18	0.44
1:A:167:TYR:CZ	1:A:217:LYS:HD2	2.53	0.44
1:B:60:ASN:H	1:B:60:ASN:HD22	1.67	0.43
1:B:17:GLY:HA3	1:B:197:LYS:HE2	1.99	0.43
1:B:17:GLY:HA2	1:B:18:PRO:C	2.43	0.43
1:A:60:ASN:H	1:A:60:ASN:HD22	1.67	0.43
1:A:133:HIS:HD2	1:A:135:GLU:OE1	2.02	0.43
1:B:12:ILE:HG23	4:B:960:GOL:H12	2.00	0.43
1:A:196:ASN:HA	1:A:201:ALA:O	2.19	0.42
1:A:338[A]:GLU:O	1:A:342[A]:LYS:NZ	2.46	0.42
1:B:65:LEU:HB2	1:B:206:THR:HB	2.02	0.42
1:B:2:MET:O	1:B:3[A]:ARG:HD3	2.20	0.41
1:B:196:ASN:HA	1:B:201:ALA:O	2.20	0.41
1:A:17:GLY:HA2	1:A:18:PRO:C	2.45	0.41
1:A:229:MET:HE3	1:A:229:MET:HB2	1.77	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/392 (102%)	394 (98%)	6 (2%)	0	100	100
1	B	398/392 (102%)	391 (98%)	7 (2%)	0	100	100
All	All	798/784 (102%)	785 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	309/299 (103%)	303 (98%)	6 (2%)	50 34
1	B	307/299 (103%)	301 (98%)	6 (2%)	48 32
All	All	616/598 (103%)	604 (98%)	12 (2%)	48 34

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	ASN
1	A	68	GLU
1	A	146	THR
1	A	229	MET
1	A	376	LEU
1	A	380	LYS
1	B	20	ASN
1	B	60	ASN
1	B	68	GLU
1	B	146	THR
1	B	196	ASN
1	B	229	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	GLN
1	A	60	ASN
1	A	109	HIS
1	A	124	GLN
1	A	133	HIS
1	A	144	HIS
1	A	255	ASN
1	A	322	GLN
1	A	354	GLN
1	A	370	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	53	GLN
1	B	60	ASN
1	B	144	HIS
1	B	196	ASN
1	B	255	ASN
1	B	322	GLN
1	B	370	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	PXL	B	1197	1	11,11,12	0.85	1 (9%)	15,15,16	1.53	4 (26%)
4	GOL	A	955	-	5,5,5	0.38	0	5,5,5	0.27	0
2	SO4	A	1603	-	4,4,4	0.24	0	6,6,6	0.25	0
2	SO4	A	1601	-	4,4,4	0.17	0	6,6,6	0.41	0
4	GOL	B	956	-	5,5,5	0.38	0	5,5,5	0.30	0
4	GOL	A	958	-	5,5,5	0.25	0	5,5,5	0.34	0
4	GOL	B	960	-	5,5,5	0.36	0	5,5,5	0.31	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
2	SO4	B	2604	-	4,4,4	0.26	0	6,6,6	0.20	0
4	GOL	A	953	-	5,5,5	0.46	0	5,5,5	0.76	0
4	GOL	B	954	-	5,5,5	0.39	0	5,5,5	0.28	0
2	SO4	B	2602	-	4,4,4	0.26	0	6,6,6	0.10	0
3	PXL	A	1197	1	11,11,12	0.91	0	15,15,16	1.13	2 (13%)
2	SO4	A	1605	-	4,4,4	0.24	0	6,6,6	0.25	0
4	GOL	B	957	-	5,5,5	0.26	0	5,5,5	0.40	0
2	SO4	B	2605	-	4,4,4	0.29	0	6,6,6	0.14	0
2	SO4	B	2601	-	4,4,4	0.28	0	6,6,6	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PXL	B	1197	1	-	0/2/2/4	0/1/1/1
4	GOL	A	955	-	-	0/4/4/4	-
4	GOL	B	956	-	-	4/4/4/4	-
4	GOL	A	958	-	-	1/4/4/4	-
4	GOL	B	960	-	-	2/4/4/4	-
4	GOL	A	953	-	-	0/4/4/4	-
4	GOL	B	954	-	-	4/4/4/4	-
3	PXL	A	1197	1	-	0/2/2/4	0/1/1/1
4	GOL	B	957	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1197	PXL	C2A-C2	2.15	1.53	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	1197	PXL	C5A-C5-C6	-2.94	114.61	119.36
3	B	1197	PXL	C6-C5-C4	2.75	120.35	118.10
3	B	1197	PXL	O3-C3-C2	2.30	122.36	117.58
3	A	1197	PXL	C5-C6-N1	-2.08	120.44	123.83
3	B	1197	PXL	C5A-C5-C4	2.07	125.04	121.93
3	A	1197	PXL	C6-N1-C2	2.02	122.86	119.20

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	954	GOL	O1-C1-C2-C3
4	B	954	GOL	C1-C2-C3-O3
4	B	954	GOL	O2-C2-C3-O3
4	B	956	GOL	O1-C1-C2-C3
4	B	956	GOL	C1-C2-C3-O3
4	B	956	GOL	O1-C1-C2-O2
4	B	956	GOL	O2-C2-C3-O3
4	B	960	GOL	C1-C2-C3-O3
4	B	954	GOL	O1-C1-C2-O2
4	B	960	GOL	O2-C2-C3-O3
4	A	958	GOL	O1-C1-C2-C3

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1197	PXL	1	0
4	A	955	GOL	3	0
4	B	960	GOL	1	0
4	A	953	GOL	4	0
2	A	1605	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	392/392 (100%)	-0.58	3 (0%)	82	85	7, 10, 17, 26	10 (2%)
1	B	392/392 (100%)	-0.70	1 (0%)	90	91	4, 9, 16, 24	8 (2%)
All	All	784/784 (100%)	-0.64	4 (0%)	87	89	4, 10, 17, 26	18 (2%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	393	ALA	4.8
1	B	393	ALA	4.0
1	A	313	ASP	2.2
1	A	20[A]	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	SO4	A	1605	5/5	0.75	0.17	71,71,71,71	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	GOL	B	954	6/6	0.77	0.17	27,33,34,37	0
4	GOL	A	953	6/6	0.78	0.14	26,31,33,34	0
4	GOL	B	956	6/6	0.80	0.15	38,40,40,42	0
2	SO4	B	2602	5/5	0.84	0.11	56,56,57,57	0
4	GOL	B	960	6/6	0.84	0.14	32,36,38,38	0
4	GOL	A	955	6/6	0.88	0.10	31,32,33,34	0
4	GOL	A	958	6/6	0.91	0.09	12,19,20,20	0
4	GOL	B	957	6/6	0.92	0.10	15,22,24,25	0
2	SO4	A	1603	5/5	0.92	0.13	31,31,33,34	0
2	SO4	B	2605	5/5	0.93	0.15	39,39,40,40	0
2	SO4	B	2604	5/5	0.93	0.10	33,34,35,36	0
2	SO4	A	1601	5/5	0.94	0.09	27,27,30,30	0
3	PXL	B	1197	11/12	0.95	0.06	8,8,10,11	0
3	PXL	A	1197	11/12	0.96	0.06	9,10,13,14	0
2	SO4	B	2601	5/5	0.97	0.07	22,23,25,28	0

6.5 Other polymers (i)

There are no such residues in this entry.