



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2026 – 06:49 PM UTC

PDB ID : 6Z98 / pdb_00006z98
BMRB ID : 27687
Title : NMR solution structure of the peach allergen Pru p 1.0101
Authors : Eidelpes, R.; Fuehrer, S.; Hofer, F.; Kamenik, A.S.; Liedl, K.R.; Tollinger, M.
Deposited on : 2020-06-03

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 82%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	159	 77% 16% 6% •

2 Ensemble composition and analysis

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:159 (158)	0.42	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 9 single-model clusters were found.

Cluster number	Models
1	3, 4, 14
2	8, 15
3	9, 20
4	1, 2
5	6, 12
Single-model clusters	5; 7; 10; 11; 13; 16; 17; 18; 19

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2469 atoms, of which 1229 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Major allergen Pru p 1.


Mol	Chain	Residues	Atoms					Trace
1	A	159	Total	C	H	N	O	0
			2469	792	1229	201	247	

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Major allergen Pru p 1




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Major allergen Pru p 1



4.2.2 Score per residue for model 2

- Molecule 1: Major allergen Pru p 1






4.2.3 Score per residue for model 3

- Molecule 1: Major allergen Pru p 1


Chain A: 74% 18% 8% ..



4.2.4 Score per residue for model 4

- Molecule 1: Major allergen Pru p 1

Chain A: 75% 18% 7% .



4.2.5 Score per residue for model 5

- Molecule 1: Major allergen Pru p 1

Chain A: 77% 15% 7% .



4.2.6 Score per residue for model 6

- Molecule 1: Major allergen Pru p 1


Chain A: 73% 19% 7% ..






4.2.7 Score per residue for model 7

- Molecule 1: Major allergen Pru p 1




4.2.8 Score per residue for model 8 (medoid)

- Molecule 1: Major allergen Pru p 1




4.2.9 Score per residue for model 9

- Molecule 1: Major allergen Pru p 1



4.2.10 Score per residue for model 10


- Molecule 1: Major allergen Pru p 1






4.2.11 Score per residue for model 11

- Molecule 1: Major allergen Pru p 1




4.2.12 Score per residue for model 12

- Molecule 1: Major allergen Pru p 1



4.2.13 Score per residue for model 13


- Molecule 1: Major allergen Pru p 1



4.2.14 Score per residue for model 14


- Molecule 1: Major allergen Pru p 1






4.2.15 Score per residue for model 15

- Molecule 1: Major allergen Pru p 1




4.2.16 Score per residue for model 16

- Molecule 1: Major allergen Pru p 1




4.2.17 Score per residue for model 17

- Molecule 1: Major allergen Pru p 1




4.2.18 Score per residue for model 18

- Molecule 1: Major allergen Pru p 1




4.2.19 Score per residue for model 19

- Molecule 1: Major allergen Pru p 1



4.2.20 Score per residue for model 20

- Molecule 1: Major allergen Pru p 1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 40 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure calculation	
Amber	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1714
Number of shifts mapped to atoms	1714
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	82%

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.91±0.01	0±0/1265 (0.0± 0.0%)	1.72±0.02	28±3/1706 (1.6± 0.2%)
All	All	0.91	7/25300 (0.0%)	1.72	555/34120 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	7.3±1.7
All	All	0	147

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	14	PRO	CA-C	6.25	1.55	1.51	3	7

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	154	HIS	CA-C-N	10.84	132.16	120.12	7	20
1	A	154	HIS	C-N-CA	10.84	132.16	120.12	7	20
1	A	42	GLU	N-CA-C	9.36	124.66	109.59	6	20
1	A	43	ILE	CB-CA-C	8.94	122.87	112.68	6	20
1	A	53	ILE	N-CA-C	-8.59	93.06	106.72	12	20
1	A	42	GLU	CA-C-O	8.23	129.36	120.38	8	20
1	A	144	PHE	CA-CB-CG	7.91	121.71	113.80	1	5
1	A	53	ILE	O-C-N	7.88	130.36	122.71	20	17
1	A	53	ILE	N-CA-CB	7.74	121.51	112.15	7	20
1	A	147	ILE	CA-C-O	-7.69	112.26	120.64	17	20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	146	LEU	CA-C-N	7.68	133.22	120.62	20	20
1	A	146	LEU	C-N-CA	7.68	133.22	120.62	20	20
1	A	139	LYS	N-CA-C	-7.41	102.36	111.40	7	20
1	A	148	GLU	N-CA-CB	7.40	120.72	109.91	11	19
1	A	12	GLU	N-CA-CB	-7.40	99.92	110.58	20	20
1	A	54	LYS	CB-CA-C	7.26	122.78	110.81	11	20
1	A	27	ASP	CA-CB-CG	7.22	119.82	112.60	14	1
1	A	133	VAL	CA-C-N	7.20	130.16	120.65	13	10
1	A	133	VAL	C-N-CA	7.20	130.16	120.65	13	10
1	A	69	HIS	CA-CB-CG	6.80	120.61	113.80	11	2
1	A	52	THR	CA-C-N	-6.71	113.97	123.10	12	20
1	A	52	THR	C-N-CA	-6.71	113.97	123.10	12	20
1	A	13	ILE	CA-C-N	6.60	124.42	119.66	1	9
1	A	13	ILE	C-N-CA	6.60	124.42	119.66	1	9
1	A	43	ILE	N-CA-CB	-6.40	105.85	112.06	7	3
1	A	2	VAL	N-CA-C	6.35	117.03	107.75	20	3
1	A	14	PRO	N-CA-CB	6.25	106.47	103.22	12	4
1	A	55	LYS	CB-CA-C	6.23	120.94	110.79	6	18
1	A	146	LEU	N-CA-C	-6.19	103.85	111.40	16	8
1	A	14	PRO	N-CA-C	6.01	116.24	110.47	17	2
1	A	142	ASN	CA-CB-CG	6.01	118.61	112.60	7	3
1	A	29	LEU	CA-C-N	5.87	125.03	120.33	2	5
1	A	29	LEU	C-N-CA	5.87	125.03	120.33	2	5
1	A	155	PRO	CA-N-CD	-5.85	103.81	112.00	19	10
1	A	34	ALA	CA-C-N	5.80	125.47	119.56	8	3
1	A	34	ALA	C-N-CA	5.80	125.47	119.56	8	3
1	A	140	ALA	CB-CA-C	5.78	120.68	110.85	15	7
1	A	15	PRO	N-CA-C	5.72	117.68	110.70	3	10
1	A	151	LEU	CA-C-N	5.64	133.66	121.64	15	19
1	A	151	LEU	C-N-CA	5.64	133.66	121.64	15	19
1	A	156	ASP	CA-C-N	5.52	127.62	120.44	4	5
1	A	156	ASP	C-N-CA	5.52	127.62	120.44	4	5
1	A	79	HIS	CB-CG-CD2	-5.52	124.02	131.20	16	13
1	A	94	ASN	CA-CB-CG	5.47	118.07	112.60	11	3
1	A	42	GLU	O-C-N	-5.45	116.85	123.29	19	6
1	A	132	HIS	CB-CG-CD2	-5.42	124.16	131.20	10	5
1	A	75	ASP	CA-C-N	5.41	128.07	120.28	7	1
1	A	75	ASP	C-N-CA	5.41	128.07	120.28	7	1
1	A	63	GLN	OE1-CD-NE2	-5.40	117.20	122.60	14	1
1	A	148	GLU	N-CA-C	-5.22	105.28	110.97	11	1
1	A	137	LYS	CA-C-N	5.17	129.35	120.72	10	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	137	LYS	C-N-CA	5.17	129.35	120.72	10	1
1	A	40	HIS	CB-CG-CD2	-5.09	124.58	131.20	8	1
1	A	59	GLY	CA-C-N	5.07	127.78	120.38	8	1
1	A	59	GLY	C-N-CA	5.07	127.78	120.38	8	1
1	A	156	ASP	CA-CB-CG	5.07	117.67	112.60	19	1
1	A	85	LEU	CB-CA-C	5.06	118.23	110.14	10	2
1	A	154	HIS	CA-C-O	-5.06	116.22	121.07	19	1
1	A	23	VAL	CA-CB-CG1	5.04	118.96	110.40	17	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	2	VAL	Peptide	20
1	A	155	PRO	Peptide	20
1	A	158	TYR	Sidechain	20
1	A	17	ARG	Sidechain	19
1	A	44	LEU	Peptide	16
1	A	5	TYR	Sidechain	13
1	A	156	ASP	Peptide	7
1	A	46	GLY	Peptide	6
1	A	55	LYS	Peptide	5
1	A	10	THR	Peptide	5
1	A	81	TYR	Sidechain	3
1	A	83	TYR	Sidechain	3
1	A	52	THR	Peptide	2
1	A	23	VAL	Peptide	2
1	A	42	GLU	Peptide	1
1	A	64	TYR	Sidechain	1
1	A	91	LEU	Peptide	1
1	A	125	ASP	Peptide	1
1	A	100	TYR	Peptide	1
1	A	120	TYR	Sidechain	1

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1236	1224	1217	6±1
All	All	24720	24480	24340	113

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:11:SER:HB3	1:A:148:GLU:HA	0.61	1.72	9	20
1:A:11:SER:CB	1:A:148:GLU:HA	0.58	2.29	13	20
1:A:151:LEU:HD22	1:A:157:ALA:HB1	0.54	1.79	9	1
1:A:18:LEU:HD12	1:A:112:SER:HB3	0.54	1.80	18	2
1:A:143:LEU:O	1:A:147:ILE:HG22	0.50	2.06	14	16
1:A:34:ALA:N	1:A:35:PRO:CD	0.48	2.76	4	20
1:A:42:GLU:O	1:A:54:LYS:HA	0.46	2.11	14	20
1:A:5:TYR:CD1	1:A:120:TYR:CE1	0.44	3.05	2	5
1:A:140:ALA:HA	1:A:143:LEU:HD12	0.42	1.92	12	1
1:A:11:SER:HB2	1:A:148:GLU:HG3	0.41	1.91	2	1
1:A:147:ILE:O	1:A:148:GLU:C	0.41	2.64	14	7

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	157/159 (99%)	143±2 (91±1%)	12±2 (8±1%)	2±1 (1±0%)	14	63
All	All	3140/3180 (99%)	2861 (91%)	245 (8%)	34 (1%)	14	63

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	156	ASP	20
1	A	92	GLY	10
1	A	34	ALA	3
1	A	65	GLY	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/135 (100%)	119±2 (88±1%)	16±2 (12±1%)	7	50
All	All	2700/2700 (100%)	2385 (88%)	315 (12%)	7	50

All 43 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	29	LEU	20
1	A	53	ILE	20
1	A	78	ASN	20
1	A	86	ILE	20
1	A	91	LEU	20
1	A	151	LEU	20
1	A	97	LYS	19
1	A	146	LEU	19
1	A	24	LEU	16
1	A	43	ILE	15
1	A	23	VAL	14
1	A	13	ILE	13
1	A	129	LYS	12
1	A	149	THR	11
1	A	17	ARG	10
1	A	102	THR	7
1	A	152	LYS	6
1	A	105	VAL	6
1	A	143	LEU	5
1	A	147	ILE	5
1	A	155	PRO	4
1	A	84	THR	4
1	A	85	LEU	2
1	A	25	ASP	2
1	A	156	ASP	2
1	A	79	HIS	2
1	A	33	ILE	2
1	A	81	TYR	2
1	A	94	ASN	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	83	TYR	2
1	A	60	GLU	1
1	A	109	SER	1
1	A	142	ASN	1
1	A	148	GLU	1
1	A	12	GLU	1
1	A	30	VAL	1
1	A	123	LYS	1
1	A	130	GLU	1
1	A	131	GLU	1
1	A	27	ASP	1
1	A	55	LYS	1
1	A	116	SER	1
1	A	121	HIS	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 82% for the well-defined parts and 81% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1714
Number of shifts mapped to atoms	1714
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	152	-0.30 ± 0.09	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	138	-0.11 ± 0.08	None needed (< 0.5 ppm)
$^{13}\text{C}'$	150	0.02 ± 0.07	None needed (< 0.5 ppm)
^{15}N	142	0.01 ± 0.26	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 82%, i.e. 1710 atoms were assigned a chemical shift out of a possible 2098. 0 out of 18 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	745/788 (95%)	303/322 (94%)	300/316 (95%)	142/150 (95%)
Sidechain	965/1113 (87%)	644/722 (89%)	313/363 (86%)	8/28 (29%)

Continued on next page...

Continued from previous page...

	Total	^1H	^{13}C	^{15}N
Aromatic	0/197 (0%)	0/94 (0%)	0/89 (0%)	0/14 (0%)
Overall	1710/2098 (82%)	947/1138 (83%)	613/768 (80%)	150/192 (78%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 81%, i.e. 1714 atoms were assigned a chemical shift out of a possible 2104. 0 out of 18 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	749/794 (94%)	305/325 (94%)	302/318 (95%)	142/151 (94%)
Sidechain	965/1113 (87%)	644/722 (89%)	313/363 (86%)	8/28 (29%)
Aromatic	0/197 (0%)	0/94 (0%)	0/89 (0%)	0/14 (0%)
Overall	1714/2104 (81%)	949/1141 (83%)	615/770 (80%)	150/193 (78%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

