



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 05:49 PM UTC

PDB ID : 2ZAT / pdb_00002zat
Title : Crystal structure of a mammalian reductase
Authors : Tanaka, N.; Aoki, K.; Nakamura, K.T.
Deposited on : 2007-10-10
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	260	90% 5% . .
1	B	260	90% . . .
1	C	260	89% 7% .
1	D	260	90% 6% .

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8480 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dehydrogenase/reductase SDR family member 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	0	0	0
			1879	1179	337	354	9			
1	B	251	Total	C	N	O	S	0	0	0
			1879	1179	337	354	9			
1	C	251	Total	C	N	O	S	0	0	0
			1879	1179	337	354	9			
1	D	251	Total	C	N	O	S	0	0	0
			1879	1179	337	354	9			

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	178	Total	O	0	0
			178	178		
3	B	197	Total	O	0	0
			197	197		
3	C	196	Total	O	0	0
			196	196		
3	D	201	Total	O	0	0
			201	201		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Dehydrogenase/reductase SDR family member 4

Chain A:  90% 5% . .



- Molecule 1: Dehydrogenase/reductase SDR family member 4

Chain B:  90% . . .



- Molecule 1: Dehydrogenase/reductase SDR family member 4

Chain C:  89% 7% .



- Molecule 1: Dehydrogenase/reductase SDR family member 4

Chain D:  90% 6% .



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 42	Depositor
Cell constants a, b, c, α , β , γ	109.61Å 109.61Å 94.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.50	Depositor
% Data completeness (in resolution range)	99.0 (40.00-1.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.161 , 0.187	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	8480	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	2/1905 (0.1%)	0.85	1/2582 (0.0%)
1	B	0.79	1/1905 (0.1%)	0.84	2/2582 (0.1%)
1	C	0.80	2/1905 (0.1%)	0.81	2/2582 (0.1%)
1	D	0.79	1/1905 (0.1%)	0.85	0/2582
All	All	0.80	6/7620 (0.1%)	0.84	5/10328 (0.0%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	130	MET	SD-CE	-8.50	1.58	1.79
1	C	130	MET	SD-CE	-8.10	1.59	1.79
1	A	130	MET	SD-CE	-7.35	1.61	1.79
1	C	135	VAL	CA-CB	6.85	1.57	1.54
1	D	130	MET	SD-CE	-6.80	1.62	1.79
1	A	135	VAL	CA-CB	6.63	1.57	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	157	PRO	N-CA-C	5.76	119.91	111.03
1	A	204	VAL	CB-CA-C	-5.61	103.46	112.16
1	B	108	ILE	N-CA-C	5.47	116.85	110.62
1	C	102	ASN	O-C-N	5.28	124.92	121.23
1	C	108	ILE	N-CA-C	5.16	116.51	110.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1879	0	1937	7	0
1	B	1879	0	1937	12	0
1	C	1879	0	1937	11	0
1	D	1879	0	1937	13	0
2	A	48	0	25	2	0
2	B	48	0	25	5	0
2	C	48	0	25	3	0
2	D	48	0	25	2	0
3	A	178	0	0	3	0
3	B	197	0	0	9	0
3	C	196	0	0	4	0
3	D	201	0	0	2	0
All	All	8480	0	7848	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:3261:NAP:O3D	3:C:3448:HOH:O	1.78	0.99
2:B:2261:NAP:C5N	3:B:2392:HOH:O	2.37	0.71
1:B:104:PHE:O	3:B:2447:HOH:O	2.11	0.69
1:C:196:LEU:HD21	1:C:219:LEU:HG	1.80	0.64
1:D:196:LEU:HD13	1:D:224:LEU:HD22	1.78	0.64
1:A:139:GLU:OE1	1:A:185:ARG:NE	2.31	0.63
1:A:104:PHE:O	3:A:1437:HOH:O	2.15	0.63
1:B:161:LEU:HA	3:B:2447:HOH:O	2.01	0.60
2:B:2261:NAP:H5N	3:B:2444:HOH:O	2.00	0.60
2:B:2261:NAP:H5N	3:B:2392:HOH:O	1.97	0.59
2:C:3261:NAP:C5N	3:C:3447:HOH:O	2.51	0.58
1:A:98:ASN:OD1	2:A:1261:NAP:H4D	2.09	0.52
1:B:156:HIS:NE2	1:D:156:HIS:CE1	2.79	0.50
1:C:125:LYS:HE3	1:D:113:GLU:HG3	1.94	0.50
1:D:158:PHE:CB	1:D:161:LEU:HD12	2.42	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:125:LYS:CE	1:D:113:GLU:HG3	2.43	0.49
2:A:1261:NAP:C5N	3:A:1426:HOH:O	2.60	0.48
1:B:156:HIS:NE2	1:D:156:HIS:HE1	2.12	0.48
1:B:107:ASN:HD22	1:B:109:ILE:H	1.62	0.48
1:D:158:PHE:HB2	1:D:161:LEU:HD12	1.95	0.48
1:C:248:GLU:OE1	3:C:3454:HOH:O	2.20	0.47
1:B:107:ASN:CG	3:B:2407:HOH:O	2.57	0.47
1:B:196:LEU:HD13	1:B:224:LEU:HD22	1.98	0.46
1:D:98:ASN:OD1	2:D:4261:NAP:H4D	2.15	0.46
1:A:181:GLU:OE1	1:B:108:ILE:HG22	2.16	0.46
2:D:4261:NAP:C5N	3:D:4445:HOH:O	2.64	0.45
1:C:98:ASN:OD1	2:C:3261:NAP:H4D	2.16	0.45
1:A:161:LEU:HA	3:A:1437:HOH:O	2.15	0.45
1:C:152:VAL:HG22	1:C:251:VAL:HG13	1.98	0.45
1:B:132:LYS:NZ	3:B:2451:HOH:O	2.48	0.45
1:C:137:GLU:HG2	3:C:3450:HOH:O	2.15	0.45
1:A:196:LEU:HD13	1:A:224:LEU:HD22	1.99	0.45
1:B:18:LEU:HD21	1:B:85:ALA:HB2	1.99	0.44
1:A:159:PRO:O	1:A:160:ASN:HB2	2.18	0.43
1:B:107:ASN:ND2	3:B:2407:HOH:O	2.50	0.43
1:C:113:GLU:HG2	1:D:129:LEU:HD11	2.01	0.42
2:B:2261:NAP:C5N	3:B:2444:HOH:O	2.63	0.42
1:C:28:LEU:HD13	1:C:54:THR:HG23	2.00	0.42
1:D:205:LEU:HD22	3:D:4362:HOH:O	2.19	0.41
1:D:18:LEU:HD21	1:D:85:ALA:HB2	2.02	0.41
1:B:199:THR:OG1	2:B:2261:NAP:N7N	2.53	0.41
1:C:177:ASN:OD1	1:D:157:PRO:HB3	2.21	0.41
1:C:181:GLU:OE1	1:D:107:ASN:HA	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/260 (96%)	243 (98%)	6 (2%)	0	100	100
1	B	249/260 (96%)	244 (98%)	5 (2%)	0	100	100
1	C	249/260 (96%)	241 (97%)	7 (3%)	1 (0%)	30	12
1	D	249/260 (96%)	243 (98%)	6 (2%)	0	100	100
All	All	996/1040 (96%)	971 (98%)	24 (2%)	1 (0%)	48	24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	207	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/211 (97%)	199 (98%)	5 (2%)	42	14
1	B	204/211 (97%)	198 (97%)	6 (3%)	37	10
1	C	204/211 (97%)	202 (99%)	2 (1%)	68	45
1	D	204/211 (97%)	200 (98%)	4 (2%)	48	20
All	All	816/844 (97%)	799 (98%)	17 (2%)	47	19

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	185	ARG
1	A	196	LEU
1	A	207	MET
1	A	208	ASP
1	A	220	ARG
1	B	107	ASN
1	B	109	ILE
1	B	132	LYS
1	B	196	LEU
1	B	211	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	213	GLU
1	C	207	MET
1	C	209	LYS
1	D	114	GLU
1	D	196	LEU
1	D	207	MET
1	D	213	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	102	ASN
1	B	48	GLN
1	B	107	ASN
1	C	160	ASN
1	C	226	ASN
1	D	102	ASN
1	D	156	HIS
1	D	160	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	A	1261	-	50,52,52	1.84	7 (14%)	71,80,80	1.53	11 (15%)
2	NAP	B	2261	-	50,52,52	1.92	8 (16%)	71,80,80	1.78	18 (25%)
2	NAP	C	3261	-	50,52,52	1.87	7 (14%)	71,80,80	1.55	16 (22%)
2	NAP	D	4261	-	50,52,52	1.73	6 (12%)	71,80,80	1.57	14 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	1261	-	-	6/35/67/67	0/5/5/5
2	NAP	B	2261	-	-	5/35/67/67	0/5/5/5
2	NAP	C	3261	-	-	6/35/67/67	0/5/5/5
2	NAP	D	4261	-	-	6/35/67/67	0/5/5/5

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2261	NAP	O7N-C7N	9.91	1.42	1.24
2	A	1261	NAP	O7N-C7N	9.87	1.42	1.24
2	D	4261	NAP	O7N-C7N	9.50	1.41	1.24
2	C	3261	NAP	O7N-C7N	9.38	1.41	1.24
2	C	3261	NAP	PA-O3	4.47	1.64	1.59
2	B	2261	NAP	PA-O3	4.22	1.64	1.59
2	A	1261	NAP	PA-O3	3.79	1.63	1.59
2	C	3261	NAP	PN-O3	3.42	1.63	1.59
2	B	2261	NAP	PN-O3	3.23	1.63	1.59
2	A	1261	NAP	PN-O3	3.08	1.62	1.59
2	A	1261	NAP	C2A-N1A	2.90	1.39	1.33
2	D	4261	NAP	PN-O3	2.82	1.62	1.59
2	B	2261	NAP	C2A-N1A	2.72	1.38	1.33
2	D	4261	NAP	PA-O3	2.72	1.62	1.59
2	C	3261	NAP	C2A-N1A	2.64	1.38	1.33
2	C	3261	NAP	C2A-N3A	2.61	1.38	1.33
2	D	4261	NAP	C2N-N1N	2.47	1.37	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2261	NAP	C2N-N1N	2.47	1.37	1.35
2	D	4261	NAP	C2A-N3A	2.36	1.38	1.33
2	A	1261	NAP	C2A-N3A	2.34	1.38	1.33
2	B	2261	NAP	C3N-C7N	-2.34	1.47	1.50
2	A	1261	NAP	P2B-O2B	2.33	1.63	1.59
2	C	3261	NAP	P2B-O2B	2.27	1.63	1.59
2	B	2261	NAP	PN-O2N	-2.26	1.44	1.55
2	A	1261	NAP	C8A-N7A	2.13	1.35	1.31
2	B	2261	NAP	C2A-N3A	2.12	1.37	1.33
2	C	3261	NAP	PN-O2N	-2.04	1.45	1.55
2	D	4261	NAP	C8A-N7A	2.01	1.35	1.31

All (59) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4261	NAP	N3A-C2A-N1A	-4.76	121.38	128.58
2	A	1261	NAP	N3A-C2A-N1A	-4.61	121.60	128.58
2	B	2261	NAP	C2B-C1B-N9A	4.48	121.12	113.75
2	A	1261	NAP	C2B-C1B-N9A	4.22	120.70	113.75
2	D	4261	NAP	C5A-C4A-N3A	-4.19	120.94	126.72
2	B	2261	NAP	O7N-C7N-C3N	-4.14	114.53	119.60
2	C	3261	NAP	O7N-C7N-C3N	-3.51	115.30	119.60
2	B	2261	NAP	C4A-C5A-N7A	-3.39	106.70	110.58
2	B	2261	NAP	C4D-O4D-C1D	-3.33	106.87	109.92
2	B	2261	NAP	N9A-C8A-N7A	-3.33	109.21	113.94
2	B	2261	NAP	C5A-N7A-C8A	3.33	108.68	103.45
2	C	3261	NAP	C5A-C4A-N3A	-3.29	122.19	126.72
2	D	4261	NAP	O7N-C7N-C3N	-3.26	115.61	119.60
2	A	1261	NAP	O2A-PA-O3	3.25	116.06	107.27
2	C	3261	NAP	N9A-C8A-N7A	-3.24	109.34	113.94
2	B	2261	NAP	C2N-C3N-C4N	3.20	121.98	118.26
2	A	1261	NAP	C5A-C4A-N3A	-3.20	122.31	126.72
2	C	3261	NAP	C2B-C1B-N9A	3.18	118.99	113.75
2	D	4261	NAP	C2A-N3A-C4A	3.18	119.60	111.83
2	B	2261	NAP	C4N-C3N-C7N	-3.17	112.42	121.06
2	C	3261	NAP	N3A-C2A-N1A	-3.13	123.84	128.58
2	A	1261	NAP	C4D-O4D-C1D	-3.12	107.07	109.92
2	B	2261	NAP	C5A-C4A-N3A	-3.11	122.43	126.72
2	C	3261	NAP	C5A-N7A-C8A	3.10	108.32	103.45
2	D	4261	NAP	C2B-C1B-N9A	3.09	118.84	113.75
2	D	4261	NAP	C2N-C3N-C4N	3.07	121.83	118.26
2	B	2261	NAP	N3A-C2A-N1A	-3.06	123.96	128.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4261	NAP	N9A-C8A-N7A	-3.00	109.68	113.94
2	A	1261	NAP	N9A-C8A-N7A	-2.99	109.69	113.94
2	B	2261	NAP	O7N-C7N-N7N	2.95	126.87	122.62
2	C	3261	NAP	O2A-PA-O3	2.90	115.10	107.27
2	C	3261	NAP	C4A-C5A-N7A	-2.85	107.32	110.58
2	B	2261	NAP	C6N-N1N-C1D	-2.82	114.19	119.73
2	C	3261	NAP	C2N-C3N-C4N	2.81	121.53	118.26
2	A	1261	NAP	C4A-C5A-N7A	-2.77	107.41	110.58
2	A	1261	NAP	C5A-N7A-C8A	2.71	107.71	103.45
2	B	2261	NAP	O2A-PA-O3	2.66	114.47	107.27
2	A	1261	NAP	C2A-N3A-C4A	2.65	118.29	111.83
2	A	1261	NAP	C2N-C3N-C4N	2.56	121.24	118.26
2	D	4261	NAP	O2A-PA-O3	2.56	114.18	107.27
2	B	2261	NAP	O2N-PN-O3	2.46	113.93	107.27
2	C	3261	NAP	O3B-C3B-C4B	-2.42	104.14	111.08
2	B	2261	NAP	C5A-C4A-N9A	2.34	108.36	105.81
2	C	3261	NAP	O2N-PN-O3	2.28	113.44	107.27
2	D	4261	NAP	C5A-N7A-C8A	2.28	107.03	103.45
2	C	3261	NAP	C5A-C4A-N9A	2.23	108.24	105.81
2	C	3261	NAP	O4B-C1B-C2B	-2.21	102.78	106.59
2	D	4261	NAP	C5A-C4A-N9A	2.20	108.20	105.81
2	D	4261	NAP	C4A-C5A-N7A	-2.18	108.09	110.58
2	C	3261	NAP	C2A-N3A-C4A	2.16	117.10	111.83
2	B	2261	NAP	O4D-C4D-C3D	2.15	109.43	105.15
2	D	4261	NAP	N3A-C4A-N9A	2.13	130.79	127.17
2	B	2261	NAP	O3B-C3B-C4B	-2.12	105.00	111.08
2	D	4261	NAP	C4N-C3N-C7N	-2.07	115.41	121.06
2	A	1261	NAP	O3B-C3B-C4B	-2.02	105.28	111.08
2	C	3261	NAP	C4D-O4D-C1D	-2.01	108.08	109.92
2	B	2261	NAP	O4B-C1B-C2B	-2.01	103.13	106.59
2	D	4261	NAP	C3B-C2B-C1B	-2.01	98.96	102.81
2	C	3261	NAP	C6N-N1N-C1D	-2.01	115.79	119.73

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1261	NAP	C5D-O5D-PN-O3
2	A	1261	NAP	C5D-O5D-PN-O1N
2	A	1261	NAP	C5D-O5D-PN-O2N
2	B	2261	NAP	C5D-O5D-PN-O3
2	B	2261	NAP	C5D-O5D-PN-O1N

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	2261	NAP	C5D-O5D-PN-O2N
2	C	3261	NAP	C5D-O5D-PN-O3
2	C	3261	NAP	C5D-O5D-PN-O1N
2	C	3261	NAP	C5D-O5D-PN-O2N
2	D	4261	NAP	C5D-O5D-PN-O3
2	D	4261	NAP	C5D-O5D-PN-O1N
2	D	4261	NAP	C5D-O5D-PN-O2N
2	A	1261	NAP	PN-O3-PA-O2A
2	B	2261	NAP	PN-O3-PA-O2A
2	C	3261	NAP	PN-O3-PA-O2A
2	D	4261	NAP	PN-O3-PA-O2A
2	D	4261	NAP	PN-O3-PA-O1A
2	A	1261	NAP	O4B-C4B-C5B-O5B
2	A	1261	NAP	PN-O3-PA-O1A
2	B	2261	NAP	PN-O3-PA-O1A
2	C	3261	NAP	PN-O3-PA-O1A
2	C	3261	NAP	O4B-C4B-C5B-O5B
2	D	4261	NAP	O4B-C4B-C5B-O5B


There are no ring outliers.


4 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1261	NAP	2	0
2	B	2261	NAP	5	0
2	C	3261	NAP	3	0
2	D	4261	NAP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.