



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:46 AM UTC

PDB ID : 6ZA7 / pdb_00006za7
Title : Structure of the apo transcriptional repressor Atu1419 (VanR) from agrobacterium fabrum
Authors : Morera, S.; Vigouroux, A.; Legrand, P.
Deposited on : 2020-06-04
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	2% 86% 10% .
1	B	250	10% 82% 11% 6%
1	C	250	4% 84% 10% 5%
1	D	250	9% 86% 8% 6%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7347 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator, GntR family.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	240	Total	C	N	O	S	0	0	0
			1826	1135	342	343	6			
1	B	234	Total	C	N	O	S	0	0	0
			1780	1104	334	336	6			
1	C	238	Total	C	N	O	S	0	0	0
			1819	1131	342	340	6			
1	D	236	Total	C	N	O	S	0	0	0
			1801	1121	338	336	6			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	245	HIS	-	expression tag	UNP A9CJ36
A	246	HIS	-	expression tag	UNP A9CJ36
A	247	HIS	-	expression tag	UNP A9CJ36
A	248	HIS	-	expression tag	UNP A9CJ36
A	249	HIS	-	expression tag	UNP A9CJ36
A	250	HIS	-	expression tag	UNP A9CJ36
B	245	HIS	-	expression tag	UNP A9CJ36
B	246	HIS	-	expression tag	UNP A9CJ36
B	247	HIS	-	expression tag	UNP A9CJ36
B	248	HIS	-	expression tag	UNP A9CJ36
B	249	HIS	-	expression tag	UNP A9CJ36
B	250	HIS	-	expression tag	UNP A9CJ36
C	245	HIS	-	expression tag	UNP A9CJ36
C	246	HIS	-	expression tag	UNP A9CJ36
C	247	HIS	-	expression tag	UNP A9CJ36
C	248	HIS	-	expression tag	UNP A9CJ36
C	249	HIS	-	expression tag	UNP A9CJ36
C	250	HIS	-	expression tag	UNP A9CJ36
D	245	HIS	-	expression tag	UNP A9CJ36
D	246	HIS	-	expression tag	UNP A9CJ36
D	247	HIS	-	expression tag	UNP A9CJ36

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	248	HIS	-	expression tag	UNP A9CJ36
D	249	HIS	-	expression tag	UNP A9CJ36
D	250	HIS	-	expression tag	UNP A9CJ36

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	28	Total	O	0	0
			28	28		
6	B	16	Total	O	0	0
			16	16		
6	C	20	Total	O	0	0
			20	20		
6	D	17	Total	O	0	0
			17	17		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcriptional regulator, GntR family




- Molecule 1: Transcriptional regulator, GntR family




- Molecule 1: Transcriptional regulator, GntR family



- Molecule 1: Transcriptional regulator, GntR family





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.11Å 79.01Å 140.61Å 90.00° 95.08° 90.00°	Depositor
Resolution (Å)	43.38 – 2.34 43.38 – 2.34	Depositor EDS
% Data completeness (in resolution range)	71.5 (43.38-2.34) 71.5 (43.38-2.34)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 2.34Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.194 , 0.223 0.204 , 0.233	Depositor DCC
R_{free} test set	1718 reflections (3.58%)	wwPDB-VP
Wilson B-factor (Å ²)	54.0	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 43.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7347	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, SO4, GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/1850	1.49	5/2496 (0.2%)
1	B	0.81	0/1801	1.45	5/2426 (0.2%)
1	C	0.84	1/1844 (0.1%)	1.50	7/2487 (0.3%)
1	D	0.81	0/1825	1.44	4/2461 (0.2%)
All	All	0.82	1/7320 (0.0%)	1.47	21/9870 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	236	VAL	CA-C	6.30	1.58	1.53

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	42	ASP	CA-CB-CG	10.64	123.24	112.60
1	C	243	SER	CA-C-N	8.34	131.28	120.44
1	C	243	SER	C-N-CA	8.34	131.28	120.44
1	D	158	GLU	CB-CG-CD	6.00	122.81	112.60
1	C	158	GLU	CB-CG-CD	5.72	122.32	112.60
1	C	244	ASP	CA-C-N	5.50	132.04	121.54
1	C	244	ASP	C-N-CA	5.50	132.04	121.54
1	B	79	VAL	CA-C-N	5.46	127.44	120.56
1	B	79	VAL	C-N-CA	5.46	127.44	120.56
1	C	9	HIS	CA-C-N	5.44	125.98	120.00
1	C	9	HIS	C-N-CA	5.44	125.98	120.00
1	B	158	GLU	CB-CG-CD	5.41	121.80	112.60
1	B	129	ASP	CA-C-N	5.33	127.42	120.28
1	B	129	ASP	C-N-CA	5.33	127.42	120.28
1	D	162	SER	N-CA-C	5.32	117.99	111.82
1	A	142	HIS	CA-C-N	5.18	127.23	120.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	142	HIS	C-N-CA	5.18	127.23	120.28
1	A	129	ASP	CA-C-N	5.08	127.09	120.28
1	A	129	ASP	C-N-CA	5.08	127.09	120.28
1	D	129	ASP	CA-C-N	5.08	127.09	120.28
1	D	129	ASP	C-N-CA	5.08	127.09	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1826	0	1839	10	0
1	B	1780	0	1795	14	0
1	C	1819	0	1829	13	0
1	D	1801	0	1818	9	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
3	C	6	0	8	0	0
3	D	6	0	8	0	0
4	B	7	0	10	2	0
5	C	5	0	0	0	0
6	A	28	0	0	0	0
6	B	16	0	0	0	0
6	C	20	0	0	0	0
6	D	17	0	0	0	0
All	All	7347	0	7323	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:17:LEU:HD23	1:B:55:LEU:HD12	1.77	0.66
1:B:137:ASN:O	1:B:141:HIS:HD2	1.83	0.61
1:A:22:LEU:HD22	1:A:164:PRO:HG3	1.83	0.60
1:A:223:GLU:O	1:A:227:ARG:HG2	2.03	0.58
1:C:167:SER:HB3	1:C:170:ALA:HB2	1.85	0.57
1:B:125:VAL:HG13	1:B:186:ARG:HE	1.70	0.57
1:B:107:LEU:HD21	1:B:200:VAL:HG22	1.86	0.56
1:D:167:SER:HB3	1:D:170:ALA:HB2	1.89	0.55
1:A:167:SER:HB3	1:A:170:ALA:HB2	1.89	0.55
1:A:89:MET:HE2	1:A:156:GLU:HB3	1.88	0.54
1:C:10:GLY:O	1:C:14:VAL:HG23	2.07	0.54
1:B:212:ARG:HD2	4:B:303:PEG:H21	1.90	0.54
1:C:14:VAL:HG22	1:C:51:ALA:HA	1.91	0.53
1:D:14:VAL:HG22	1:D:51:ALA:HA	1.91	0.53
1:B:14:VAL:HG22	1:B:51:ALA:HA	1.91	0.52
1:A:229:ALA:HB1	1:A:232:LEU:HD13	1.92	0.51
1:C:49:ARG:NE	1:D:49:ARG:HD2	2.28	0.48
1:D:41:LEU:HB3	1:D:43:ILE:HD12	1.95	0.48
1:C:229:ALA:HB1	1:C:232:LEU:HD13	1.96	0.47
1:B:175:LYS:C	1:B:177:ASN:H	2.23	0.47
1:A:14:VAL:HG21	1:A:54:ARG:HD2	1.96	0.47
1:C:225:MET:HE1	1:C:236:VAL:HG21	1.95	0.47
1:A:41:LEU:HB3	1:A:43:ILE:HD12	1.97	0.46
1:C:154:ARG:O	1:C:158:GLU:HG2	2.15	0.46
1:B:173:PRO:HB2	1:B:176:ALA:HB3	1.97	0.45
1:B:10:GLY:O	1:B:14:VAL:HG23	2.17	0.45
1:D:154:ARG:O	1:D:158:GLU:HG2	2.16	0.45
1:B:14:VAL:HG21	1:B:54:ARG:HD2	1.99	0.45
1:A:137:ASN:O	1:A:141:HIS:HD2	1.99	0.44
1:D:14:VAL:HG21	1:D:54:ARG:HD2	2.00	0.44
1:B:41:LEU:HB3	1:B:43:ILE:HD12	1.98	0.44
1:C:152:MET:HG2	1:D:81:ASP:HB2	2.00	0.43
1:B:212:ARG:HD2	4:B:303:PEG:C2	2.48	0.43
1:A:164:PRO:HB2	1:A:238:GLY:HA2	2.00	0.43
1:C:81:ASP:HB2	1:D:152:MET:HG2	2.00	0.43
1:C:41:LEU:HB3	1:C:43:ILE:HD12	1.99	0.43
1:A:152:MET:HE1	1:B:163:LEU:HD13	2.03	0.41
1:D:38:ALA:HA	1:D:48:VAL:HG21	2.01	0.41
1:C:14:VAL:HG21	1:C:54:ARG:HD2	2.02	0.41
1:C:147:LEU:HD23	1:C:199:ILE:HD13	2.03	0.41
1:B:38:ALA:HA	1:B:48:VAL:HG21	2.02	0.40
1:C:38:ALA:HA	1:C:48:VAL:HG21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	238/250 (95%)	234 (98%)	4 (2%)	0	100	100
1	B	230/250 (92%)	225 (98%)	4 (2%)	1 (0%)	30	32
1	C	236/250 (94%)	231 (98%)	4 (2%)	1 (0%)	30	32
1	D	234/250 (94%)	231 (99%)	3 (1%)	0	100	100
All	All	938/1000 (94%)	921 (98%)	15 (2%)	2 (0%)	43	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	245	HIS
1	B	176	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/195 (94%)	178 (97%)	6 (3%)	33	43
1	B	180/195 (92%)	175 (97%)	5 (3%)	38	50
1	C	184/195 (94%)	181 (98%)	3 (2%)	55	67
1	D	182/195 (93%)	178 (98%)	4 (2%)	45	58
All	All	730/780 (94%)	712 (98%)	18 (2%)	42	54

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	ARG
1	A	31	ARG
1	A	35	VAL
1	A	40	LEU
1	A	152	MET
1	A	216	ARG
1	B	16	GLU
1	B	40	LEU
1	B	44	SER
1	B	125	VAL
1	B	216	ARG
1	C	40	LEU
1	C	216	ARG
1	C	245	HIS
1	D	40	LEU
1	D	86	ARG
1	D	216	ARG
1	D	242	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	115	GLN
1	A	116	GLN
1	A	143	GLN
1	B	116	GLN
1	B	177	ASN
1	C	143	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PEG	B	303	-	6,6,6	0.35	0	5,5,5	0.42	0
3	GOL	C	302	2	5,5,5	0.10	0	5,5,5	0.19	0
5	SO4	C	303	-	4,4,4	0.26	0	6,6,6	0.14	0
3	GOL	D	302	-	5,5,5	0.16	0	5,5,5	0.37	0
3	GOL	B	302	2	5,5,5	0.10	0	5,5,5	0.15	0
3	GOL	A	302	2	5,5,5	0.22	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	B	303	-	-	2/4/4/4	-
3	GOL	C	302	2	-	0/4/4/4	-
3	GOL	D	302	-	-	0/4/4/4	-
3	GOL	B	302	2	-	0/4/4/4	-
3	GOL	A	302	2	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	303	PEG	O2-C3-C4-O4
4	B	303	PEG	C4-C3-O2-C2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	303	PEG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/250 (96%)	0.31	5 (2%) 63 68	33, 59, 110, 128	0
1	B	234/250 (93%)	0.62	24 (10%) 12 14	39, 67, 134, 159	0
1	C	238/250 (95%)	0.35	9 (3%) 44 50	40, 68, 112, 155	0
1	D	236/250 (94%)	0.62	22 (9%) 14 17	43, 74, 126, 164	0
All	All	948/1000 (94%)	0.47	60 (6%) 26 31	33, 67, 121, 164	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	170	ALA	8.5
1	B	173	PRO	5.8
1	B	176	ALA	4.7
1	A	244	ASP	4.4
1	B	178	ILE	4.4
1	B	222	LEU	3.9
1	D	241	LEU	3.9
1	C	125	VAL	3.8
1	C	9	HIS	3.8
1	B	125	VAL	3.7
1	D	242	ILE	3.6
1	B	168	PRO	3.6
1	B	244	ASP	3.4
1	D	239	LEU	3.4
1	C	176	ALA	3.4
1	B	9	HIS	3.3
1	D	237	PRO	3.3
1	D	76	PHE	3.2
1	A	238	GLY	3.2
1	D	125	VAL	3.1
1	B	169	SER	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	43	ILE	3.0
1	C	172	LEU	2.9
1	B	130	PHE	2.9
1	D	40	LEU	2.9
1	B	181	PHE	2.9
1	C	178	ILE	2.8
1	B	64	LEU	2.8
1	C	246	HIS	2.7
1	B	165	PHE	2.7
1	D	133	TYR	2.7
1	D	243	SER	2.5
1	D	235	GLN	2.5
1	D	119	LEU	2.5
1	D	236	VAL	2.5
1	B	67	GLY	2.4
1	D	233	ILE	2.4
1	D	181	PHE	2.4
1	D	176	ALA	2.4
1	A	226	ILE	2.4
1	C	43	ILE	2.3
1	D	124	ARG	2.3
1	A	237	PRO	2.3
1	B	243	SER	2.3
1	B	128	VAL	2.3
1	A	230	PRO	2.3
1	D	225	MET	2.3
1	D	230	PRO	2.2
1	D	240	ALA	2.2
1	D	234	ALA	2.2
1	B	119	LEU	2.2
1	C	40	LEU	2.2
1	D	8	THR	2.2
1	B	179	GLY	2.1
1	B	124	ARG	2.1
1	C	41	LEU	2.1
1	B	102	VAL	2.1
1	B	226	ILE	2.1
1	B	121	PHE	2.1
1	B	166	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SO4	C	303	5/5	0.61	0.11	147,147,147,148	0
4	PEG	B	303	7/7	0.83	0.18	51,54,59,61	0
3	GOL	C	302	6/6	0.89	0.15	58,61,64,65	0
3	GOL	D	302	6/6	0.90	0.15	52,55,61,64	0
3	GOL	A	302	6/6	0.91	0.15	38,44,49,51	0
3	GOL	B	302	6/6	0.93	0.12	62,66,69,70	0
2	ZN	A	301	1/1	0.99	0.04	46,46,46,46	0
2	ZN	B	301	1/1	0.99	0.06	63,63,63,63	0
2	ZN	C	301	1/1	0.99	0.03	69,69,69,69	0
2	ZN	D	301	1/1	1.00	0.04	66,66,66,66	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.