



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 03:05 AM UTC

PDB ID : 6ZAR / pdb_00006zar
Title : As-isolated copper nitrite reductase from Bradyrhizobium sp. ORS 375 (two-domain) at 1.1 Å resolution (unrestrained, full matrix refinement by SHELX)
Authors : Rose, S.L.; Antonyuk, S.V.; Sasaki, D.; Yamashita, K.; Hirata, K.; Ueno, G.; Ago, H.; Eady, R.R.; Tosha, T.; Yamamoto, M.; Hasnain, S.S.
Deposited on : 2020-06-05
Resolution : 1.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1625 (1.12-1.08)
Clashscore	190562	1656 (1.12-1.08)
Ramachandran outliers	187476	1616 (1.12-1.08)
Sidechain outliers	187428	1612 (1.12-1.08)
RSRZ outliers	180081	1625 (1.12-1.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	415[A]	-	X	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	416[A]	-	X	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3491 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Copper-containing nitrite reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	345	Total	C	N	O	S	0	22	0
			2739	1743	476	507	13			


There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP H0SLX7
A	342	GLU	-	expression tag	UNP H0SLX7
A	343	ASN	-	expression tag	UNP H0SLX7
A	344	LEU	-	expression tag	UNP H0SLX7
A	345	TYR	-	expression tag	UNP H0SLX7
A	346	PHE	-	expression tag	UNP H0SLX7
A	347	GLN	-	expression tag	UNP H0SLX7

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cu	0	0
			2	2		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	1
			5	4	1		
3	A	1	Total	O	S	0	1
			5	4	1		
3	A	1	Total	O	S	0	1
			5	4	1		
3	A	1	Total	O	S	0	1
			5	4	1		
3	A	1	Total	O	S	0	1
			5	4	1		
3	A	1	Total	O	S	0	1
			5	4	1		
3	A	1	Total	O	S	0	1
			5	4	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	1
			6	3	3		
4	A	1	Total	C	O	0	1
			6	3	3		

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	1
			15	8	2	4	1		
5	A	1	Total	C	N	O	S	0	1
			15	8	2	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	638	Total	O	0	144
			657	657		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Copper-containing nitrite reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	106.72Å 106.72Å 106.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.10 30.00 – 1.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-1.10) 99.7 (30.00-1.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 1.10Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.125 , 0.149 0.133 , 0.134	Depositor DCC
R_{free} test set	8257 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	9.9	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 119.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.021 for l,-k,h	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	3491	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, CU, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	2/2884 (0.1%)	1.25	20/3914 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	240	HIS	ND1-CE1	-7.31	1.25	1.32
1	A	117	ARG	CD-NE	-5.82	1.38	1.46

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	ARG	NE-CZ-NH1	-9.11	112.39	121.50
1	A	341	ASN	OD1-CG-ND2	-8.56	114.04	122.60
1	A	154	ARG	CG-CD-NE	-7.87	94.69	112.00
1	A	154	ARG	CD-NE-CZ	7.10	134.34	124.40
1	A	4	LEU	CA-C-O	-6.79	113.31	120.58
1	A	4	LEU	O-C-N	6.78	131.07	122.93
1	A	240	HIS	ND1-CG-CD2	-6.74	99.36	106.10
1	A	117	ARG	NE-CZ-NH2	6.69	125.22	119.20
1	A	229	ASN	CA-CB-CG	-6.47	106.13	112.60
1	A	9	GLN	OE1-CD-NE2	6.38	128.98	122.60
1	A	166	ARG	NE-CZ-NH1	-6.34	115.16	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	337	THR	CA-C-O	5.95	126.81	119.97
1	A	177	ASP	CA-CB-CG	5.80	118.40	112.60
1	A	182	ARG	NH1-CZ-NH2	-5.52	112.12	119.30
1	A	240	HIS	CG-ND1-CE1	5.50	118.64	109.30
1	A	166	ARG	NH1-CZ-NH2	5.17	126.02	119.30
1	A	12	ASP	CA-CB-CG	-5.16	107.44	112.60
1	A	4	LEU	CA-C-N	-5.10	115.21	122.41
1	A	4	LEU	C-N-CA	-5.10	115.21	122.41
1	A	154	ARG	NE-CZ-NH1	-5.08	116.42	121.50

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	299	VAL	Peptide
1	A	301	HIS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2739	0	2687	21	0
2	A	2	0	0	0	0
3	A	45	0	0	1	0
4	A	18	0	23	7	0
5	A	30	0	35	1	0
6	A	657	0	0	7	0
All	All	3491	0	2745	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:411[A]:SO4:O1	6:A:501[A]:HOH:O	1.91	0.89

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:224:GLY:H	4:A:416[A]:GOL:H11	1.41	0.84
1:A:218[A]:LYS:HD3	6:A:1004:HOH:O	1.89	0.73
1:A:320[A]:LYS:HD2	6:A:612:HOH:O	1.91	0.70
1:A:224:GLY:N	4:A:416[A]:GOL:H11	2.12	0.64
1:A:86:THR:HG21	6:A:839:HOH:O	1.99	0.63
1:A:345:TYR:HD1	4:A:415[A]:GOL:HO1	1.45	0.62
1:A:49:LYS:HE3	6:A:867:HOH:O	2.04	0.58
1:A:314:HIS:ND1	4:A:416[A]:GOL:H12	2.20	0.56
1:A:346:PHE:HA	4:A:415[A]:GOL:H32	1.87	0.56
1:A:48[B]:GLU:HG3	6:A:750[B]:HOH:O	2.09	0.52
1:A:224:GLY:H	4:A:416[A]:GOL:C1	2.18	0.52
5:A:413[A]:EPE:C7	5:A:413[A]:EPE:N1	2.73	0.51
1:A:305:GLU:OE2	6:A:502[A]:HOH:O	2.19	0.51
1:A:345:TYR:HD1	4:A:415[A]:GOL:O1	1.96	0.49
1:A:39:VAL:HG11	1:A:89:HIS:CD2	2.51	0.45
1:A:235:ASN:OD1	1:A:287[B]:LEU:HG	2.16	0.45
1:A:252:ILE:HD12	1:A:297:ALA:HB3	2.00	0.43
1:A:10:ARG:NH1	1:A:12:ASP:OD2	2.51	0.43
1:A:43[A]:LYS:HE3	1:A:54:GLN:NE2	2.34	0.43
1:A:218[B]:LYS:HG2	1:A:219:ALA:O	2.18	0.43
1:A:15:ALA:HA	1:A:16[B]:PRO:HD3	1.90	0.41
1:A:261[A]:GLU:OE2	1:A:270:GLU:OE1	2.39	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/347 (104%)	354 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	298/283 (105%)	293 (98%)	5 (2%)	53 18

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	5	LYS
1	A	225[A]	LYS
1	A	225[B]	LYS
1	A	242	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	40	GLN
1	A	226	ASN
1	A	328	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 16 ligands modelled in this entry, 2 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	405[A]	-	4,4,4	0.60	0	6,6,6	0.14	0
5	EPE	A	414[A]	-	15,15,15	1.38	3 (20%)	19,20,20	1.47	3 (15%)
3	SO4	A	411[A]	-	4,4,4	0.63	0	6,6,6	0.19	0
4	GOL	A	412	-	5,5,5	0.90	0	5,5,5	2.24	3 (60%)
3	SO4	A	409[A]	-	4,4,4	0.53	0	6,6,6	0.42	0
3	SO4	A	410[A]	-	4,4,4	0.66	0	6,6,6	0.19	0
3	SO4	A	403[A]	-	4,4,4	0.64	0	6,6,6	0.16	0
4	GOL	A	416[A]	-	5,5,5	1.49	1 (20%)	5,5,5	3.31	3 (60%)
3	SO4	A	408[B]	-	4,4,4	0.65	0	6,6,6	0.14	0
3	SO4	A	407[A]	-	4,4,4	0.53	0	6,6,6	0.14	0
3	SO4	A	404[A]	-	4,4,4	0.67	0	6,6,6	0.15	0
3	SO4	A	406[A]	-	4,4,4	0.62	0	6,6,6	0.09	0
4	GOL	A	415[A]	-	5,5,5	0.66	0	5,5,5	1.46	2 (40%)
5	EPE	A	413[A]	-	15,15,15	1.54	4 (26%)	19,20,20	2.71	4 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EPE	A	414[A]	-	-	0/9/19/19	0/1/1/1
4	GOL	A	412	-	-	2/4/4/4	-
4	GOL	A	416[A]	-	-	2/4/4/4	-
4	GOL	A	415[A]	-	-	4/4/4/4	-
5	EPE	A	413[A]	-	-	7/9/19/19	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	413[A]	EPE	O1S-S	2.95	1.53	1.45
5	A	413[A]	EPE	C7-N4	2.67	1.53	1.47
5	A	413[A]	EPE	C9-N1	-2.63	1.41	1.47
4	A	416[A]	GOL	O1-C1	-2.52	1.31	1.42
5	A	414[A]	EPE	O3S-S	-2.32	1.38	1.47
5	A	414[A]	EPE	O1S-S	2.27	1.51	1.45
5	A	414[A]	EPE	O2S-S	2.04	1.50	1.45
5	A	413[A]	EPE	C3-N4	2.04	1.52	1.46

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	413[A]	EPE	C10-C9-N1	7.46	140.60	112.36
5	A	413[A]	EPE	O1S-S-C10	-7.26	95.76	106.73
4	A	416[A]	GOL	O1-C1-C2	5.46	134.97	110.38
5	A	414[A]	EPE	O1S-S-C10	-4.49	99.94	106.73
4	A	412	GOL	O2-C2-C3	-3.58	94.35	109.18
5	A	413[A]	EPE	O3S-S-O2S	3.57	120.33	111.40
4	A	416[A]	GOL	C3-C2-C1	-3.33	99.58	111.80
4	A	416[A]	GOL	O2-C2-C1	-3.27	95.62	109.18
5	A	414[A]	EPE	C6-N1-C2	-2.49	103.47	108.84
4	A	412	GOL	C3-C2-C1	2.25	120.06	111.80
5	A	413[A]	EPE	O3S-S-O1S	2.18	116.86	111.40
4	A	415[A]	GOL	O3-C3-C2	2.15	120.06	110.38
5	A	414[A]	EPE	C5-C6-N1	-2.06	106.49	110.65
4	A	415[A]	GOL	O1-C1-C2	2.03	119.51	110.38
4	A	412	GOL	O3-C3-C2	-2.01	101.32	110.38

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	412	GOL	C1-C2-C3-O3
4	A	415[A]	GOL	C1-C2-C3-O3
4	A	416[A]	GOL	C1-C2-C3-O3
4	A	416[A]	GOL	O2-C2-C3-O3
5	A	413[A]	EPE	C10-C9-N1-C2
4	A	415[A]	GOL	O1-C1-C2-C3
4	A	412	GOL	O2-C2-C3-O3
4	A	415[A]	GOL	O2-C2-C3-O3
5	A	413[A]	EPE	C8-C7-N4-C3
5	A	413[A]	EPE	C9-C10-S-O3S

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	413[A]	EPE	C8-C7-N4-C5
4	A	415[A]	GOL	O1-C1-C2-O2
5	A	413[A]	EPE	C9-C10-S-O1S
5	A	413[A]	EPE	C9-C10-S-O2S
5	A	413[A]	EPE	N4-C7-C8-O8

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	411[A]	SO4	1	0
4	A	416[A]	GOL	4	0
4	A	415[A]	GOL	3	0
5	A	413[A]	EPE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	345/347 (99%)	-0.35	10 (2%) 53 58	4, 10, 23, 39	34 (9%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	ASP	7.4
1	A	346	PHE	3.6
1	A	347	GLN	3.1
1	A	4	LEU	3.1
1	A	338	GLY	2.3
1	A	337	THR	2.3
1	A	344	LEU	2.3
1	A	5	LYS	2.2
1	A	340	THR	2.1
1	A	343	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	415[A]	6/6	0.86	0.13	15,29,35,46	6
4	GOL	A	412	6/6	0.88	0.14	42,47,55,56	0
3	SO4	A	408[B]	5/5	0.89	0.16	30,30,42,53	5
3	SO4	A	410[A]	5/5	0.91	0.13	21,23,29,32	5
3	SO4	A	411[A]	5/5	0.91	0.12	11,15,34,39	5
4	GOL	A	416[A]	6/6	0.91	0.11	22,30,35,38	6
3	SO4	A	403[A]	5/5	0.92	0.15	20,28,45,49	5
3	SO4	A	406[A]	5/5	0.92	0.14	24,29,41,44	5
5	EPE	A	413[A]	15/15	0.92	0.16	11,24,29,30	15
3	SO4	A	404[A]	5/5	0.93	0.10	36,38,54,55	5
3	SO4	A	409[A]	5/5	0.93	0.12	20,21,24,24	5
3	SO4	A	405[A]	5/5	0.95	0.10	30,31,51,53	5
5	EPE	A	414[A]	15/15	0.97	0.07	11,12,15,22	15
3	SO4	A	407[A]	5/5	0.98	0.06	8,10,12,16	5
2	CU	A	401	1/1	1.00	0.01	9,9,9,9	0
2	CU	A	402	1/1	1.00	0.01	10,10,10,10	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around CU A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CU A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.