



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:54 AM UTC

PDB ID : 2ZB5 / pdb_00002zb5
Title : Crystal structure of the measles virus hemagglutinin (complex-sugar-type)
Authors : Hashiguchi, T.; Kajikawa, M.; Maita, N.; Takeda, M.; Kuroki, K.; Sasaki, K.; Kohda, D.; Yanagi, Y.; Maenaka, K.
Deposited on : 2007-10-16
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	10% 58% 26% • 12%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3269 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	424	Total	C	N	O	S	0	0	0
			3241	2075	539	602	25			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	146	GLU	-	expression tag	UNP Q83625
A	147	THR	-	expression tag	UNP Q83625
A	148	GLY	-	expression tag	UNP Q83625
A	484	THR	ASN	engineered mutation	UNP Q83625
A	492	GLY	GLU	engineered mutation	UNP Q83625
A	618	GLY	-	expression tag	UNP Q83625
A	619	THR	-	expression tag	UNP Q83625
A	620	LYS	-	expression tag	UNP Q83625
A	621	HIS	-	expression tag	UNP Q83625
A	622	HIS	-	expression tag	UNP Q83625
A	623	HIS	-	expression tag	UNP Q83625
A	624	HIS	-	expression tag	UNP Q83625
A	625	HIS	-	expression tag	UNP Q83625
A	626	HIS	-	expression tag	UNP Q83625

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Hemagglutinin protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	134.52Å 134.52Å 99.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.00 30.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-3.00) 99.5 (30.00-3.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.61 (at 3.00Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.227 , 0.271 0.228 , 0.271	Depositor DCC
R_{free} test set	1283 reflections (6.87%)	wwPDB-VP
Wilson B-factor (Å ²)	51.4	Xtriage
Anisotropy	0.903	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3269	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/3322	0.95	7/4524 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	483	PHE	N-CA-C	7.04	121.25	112.24
1	A	404	ASP	N-CA-C	-6.54	103.72	112.94
1	A	559	ILE	N-CA-C	-5.77	107.88	113.53
1	A	354	HIS	N-CA-C	5.38	117.85	108.76
1	A	522	LEU	N-CA-C	5.24	117.77	109.96
1	A	471	GLU	N-CA-C	-5.10	101.38	109.59
1	A	203	LEU	N-CA-C	-5.06	104.29	110.61

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3241	0	3150	92	0
2	A	28	0	26	3	0
All	All	3269	0	3176	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:GLY:HA2	1:A:289:VAL:HG11	1.58	0.85
1:A:364:LYS:HZ2	1:A:472:TRP:CD1	1.99	0.81
1:A:593:HIS:H	2:A:11:NAG:H81	1.50	0.76
1:A:393:LEU:H	1:A:393:LEU:HD12	1.57	0.69
1:A:397:PRO:HG2	1:A:403:LYS:HB3	1.75	0.69
1:A:222:MET:HE2	1:A:291:LEU:HB2	1.76	0.67
1:A:544:SER:HB3	1:A:547:ARG:O	1.94	0.67
1:A:391:GLN:O	1:A:395:GLU:HG2	1.94	0.66
1:A:537:ALA:HA	1:A:556:ARG:CG	2.25	0.66
1:A:235:GLU:HG2	1:A:249:LEU:HD23	1.79	0.65
1:A:362:GLN:HE21	1:A:364:LYS:HE2	1.61	0.65
1:A:233:LEU:HD13	1:A:251:MET:HE3	1.80	0.63
1:A:565:GLU:HB3	1:A:584:VAL:HB	1.83	0.60
1:A:231:THR:HG21	1:A:287:CYS:HB2	1.83	0.60
1:A:364:LYS:HE3	1:A:472:TRP:CE2	2.36	0.60
1:A:249:LEU:O	1:A:250:SER:HB2	2.02	0.60
1:A:557:LEU:N	1:A:558:PRO:HD2	2.16	0.59
2:A:11:NAG:H3	2:A:11:NAG:H82	1.84	0.59
1:A:593:HIS:N	2:A:11:NAG:H81	2.18	0.59
1:A:343:ASP:HB3	1:A:346:ILE:HG12	1.85	0.58
1:A:259:VAL:HB	1:A:261:ARG:NH1	2.19	0.58
1:A:514:LEU:HD11	1:A:526:LEU:HD13	1.86	0.58
1:A:261:ARG:HD2	1:A:271:HIS:ND1	2.20	0.57
1:A:456:ILE:HB	1:A:466:VAL:HB	1.87	0.56
1:A:362:GLN:NE2	1:A:364:LYS:HE2	2.21	0.56
1:A:289:VAL:HG13	1:A:296:LEU:HD11	1.88	0.56
1:A:517:LEU:HD11	1:A:525:VAL:CG2	2.35	0.56
1:A:310:TYR:H	1:A:313:SER:HB2	1.71	0.55
1:A:385:ALA:HB2	1:A:487:ILE:HG13	1.89	0.55
1:A:535:GLU:HG3	1:A:552:PHE:HZ	1.72	0.55
1:A:283:ASP:C	1:A:284:LEU:HD22	2.31	0.55
1:A:530:ASP:OD2	1:A:532:SER:HB2	2.07	0.54
1:A:254:VAL:HG22	1:A:277:GLU:HG3	1.89	0.53
1:A:522:LEU:O	1:A:545:PRO:HD3	2.09	0.53
1:A:339:LEU:HD11	1:A:427:ILE:HD11	1.91	0.52
1:A:419:LEU:HA	1:A:423:LEU:HG	1.92	0.50
1:A:306:ILE:HG21	1:A:317:VAL:HG11	1.93	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:334:GLN:HG3	1:A:334:GLN:O	2.12	0.49
1:A:517:LEU:HD11	1:A:525:VAL:HG22	1.94	0.49
1:A:283:ASP:HB2	1:A:336:TRP:HH2	1.77	0.49
1:A:536:HIS:CD2	1:A:559:ILE:HA	2.48	0.49
1:A:321:LEU:HD22	1:A:415:VAL:HG11	1.94	0.49
1:A:364:LYS:HZ2	1:A:472:TRP:NE1	2.09	0.49
1:A:604:VAL:O	1:A:604:VAL:HG13	2.12	0.49
1:A:554:PRO:HG2	1:A:555:PHE:CD1	2.47	0.48
1:A:563:PRO:HA	1:A:585:LEU:HD23	1.96	0.48
1:A:556:ARG:CG	1:A:556:ARG:O	2.60	0.48
1:A:587:ASP:O	1:A:589:GLU:N	2.46	0.48
1:A:483:PHE:CD2	1:A:483:PHE:N	2.82	0.47
1:A:553:TYR:HE1	1:A:555:PHE:HB2	1.78	0.47
1:A:365:TRP:O	1:A:412:VAL:HA	2.15	0.47
1:A:233:LEU:HD13	1:A:251:MET:CE	2.44	0.46
1:A:556:ARG:C	1:A:558:PRO:HD2	2.40	0.46
1:A:553:TYR:CE1	1:A:555:PHE:HB2	2.50	0.46
1:A:418:SER:C	1:A:420:THR:H	2.23	0.46
1:A:445:LYS:HA	1:A:452:TYR:HD1	1.81	0.46
1:A:284:LEU:HD11	1:A:320:GLN:OE1	2.16	0.46
1:A:364:LYS:HE3	1:A:472:TRP:CZ2	2.51	0.46
1:A:364:LYS:NZ	1:A:472:TRP:CD1	2.78	0.45
1:A:363:ALA:O	1:A:414:SER:HA	2.17	0.45
1:A:432:GLY:O	1:A:433:PRO:C	2.59	0.45
1:A:436:THR:HG23	1:A:497:PRO:HG2	1.99	0.44
1:A:389:LYS:C	1:A:391:GLN:H	2.25	0.44
1:A:488:LYS:O	1:A:489:GLU:HB2	2.18	0.44
1:A:555:PHE:O	1:A:556:ARG:CG	2.66	0.44
1:A:216:VAL:HG22	1:A:234:VAL:HG12	1.99	0.44
1:A:296:LEU:HD12	1:A:297:ALA:H	1.84	0.43
1:A:376:LEU:HD23	1:A:377:ARG:HG3	2.00	0.43
1:A:557:LEU:N	1:A:558:PRO:CD	2.81	0.43
1:A:488:LYS:HD2	1:A:488:LYS:N	2.32	0.43
1:A:452:TYR:CE2	1:A:472:TRP:CZ3	3.07	0.42
1:A:366:ALA:HA	1:A:411:GLY:O	2.19	0.42
1:A:554:PRO:O	1:A:555:PHE:O	2.38	0.42
1:A:393:LEU:HD11	1:A:499:TYR:CE1	2.54	0.42
1:A:288:MET:HE2	1:A:299:LEU:HB3	2.02	0.42
1:A:294:LEU:C	1:A:326:VAL:HG23	2.43	0.42
1:A:418:SER:C	1:A:420:THR:N	2.78	0.42
1:A:221:THR:HB	1:A:569:GLU:OE1	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:350:TYR:CE1	1:A:401:PRO:HB3	2.55	0.41
1:A:484:THR:HG22	1:A:497:PRO:HA	2.02	0.41
1:A:525:VAL:HG13	1:A:542:VAL:HG22	2.01	0.41
1:A:323:LYS:HD3	1:A:417:LEU:HD22	2.02	0.41
1:A:531:THR:HG22	1:A:536:HIS:CE1	2.55	0.41
1:A:195:ARG:HH11	1:A:195:ARG:HG2	1.85	0.41
1:A:462:LEU:HD21	1:A:499:TYR:CE1	2.55	0.41
1:A:237:PRO:HA	1:A:249:LEU:HD11	2.02	0.41
1:A:371:ARG:HB3	1:A:373:ASP:OD1	2.20	0.41
1:A:422:GLU:O	1:A:423:LEU:C	2.62	0.41
1:A:375:LYS:HA	1:A:379:GLU:HB2	2.03	0.40
1:A:490:ALA:C	1:A:492:GLY:H	2.29	0.40
1:A:374:ASP:HA	1:A:431:PHE:CZ	2.56	0.40
1:A:580:ARG:NH1	1:A:599:MET:HE3	2.37	0.40
1:A:364:LYS:NZ	1:A:472:TRP:NE1	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	416/481 (86%)	354 (85%)	46 (11%)	16 (4%)	2	15

All (16) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	420	THR
1	A	555	PHE
1	A	282	ASN
1	A	302	GLY
1	A	449	ASN
1	A	250	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	419	LEU
1	A	421	VAL
1	A	536	HIS
1	A	313	SER
1	A	422	GLU
1	A	423	LEU
1	A	535	GLU
1	A	588	SER
1	A	591	GLY
1	A	400	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	354/420 (84%)	342 (97%)	12 (3%)	32 66

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	206	LEU
1	A	221	THR
1	A	251	MET
1	A	315	LYS
1	A	333	MET
1	A	376	LEU
1	A	488	LYS
1	A	510	LEU
1	A	521	ASP
1	A	525	VAL
1	A	536	HIS
1	A	577	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	225	GLN
1	A	282	ASN
1	A	362	GLN
1	A	384	GLN
1	A	449	ASN
1	A	450	ASN
1	A	461	ASN
1	A	575	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1	1	14,14,15	0.63	0	17,19,21	0.71	0
2	NAG	A	11	1	14,14,15	0.73	1 (7%)	17,19,21	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1	1	-	2/6/23/26	0/1/1/1
2	NAG	A	11	1	-	4/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	11	NAG	C1-C2	2.06	1.55	1.52

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	NAG	C8-C7-N2-C2
2	A	1	NAG	O7-C7-N2-C2
2	A	11	NAG	C8-C7-N2-C2
2	A	11	NAG	O7-C7-N2-C2
2	A	11	NAG	C1-C2-N2-C7
2	A	11	NAG	C3-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	11	NAG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	424/481 (88%)	0.36	48 (11%) 10 6	30, 51, 107, 133	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	421	VAL	7.7
1	A	187	ASN	6.5
1	A	505	ASP	5.5
1	A	607	THR	5.4
1	A	501	PRO	4.9
1	A	557	LEU	4.9
1	A	420	THR	4.7
1	A	313	SER	4.6
1	A	168	ASN	4.6
1	A	161	GLU	4.6
1	A	419	LEU	4.6
1	A	155	ALA	4.4
1	A	558	PRO	4.0
1	A	164	ASN	3.6
1	A	166	LEU	3.6
1	A	281	SER	3.4
1	A	590	SER	3.4
1	A	163	MET	3.3
1	A	499	TYR	3.3
1	A	422	GLU	3.2
1	A	157	VAL	3.1
1	A	591	GLY	3.0
1	A	506	GLY	3.0
1	A	282	ASN	3.0
1	A	160	GLU	2.9
1	A	560	LYS	2.9
1	A	165	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	500	LEU	2.6
1	A	286	ASN	2.6
1	A	188	CYS	2.6
1	A	534	VAL	2.6
1	A	489	GLU	2.6
1	A	248	GLN	2.5
1	A	536	HIS	2.5
1	A	390	ILE	2.5
1	A	284	LEU	2.5
1	A	162	LEU	2.5
1	A	316	GLY	2.4
1	A	462	LEU	2.4
1	A	156	ASP	2.3
1	A	158	ALA	2.3
1	A	423	LEU	2.3
1	A	494	CYS	2.3
1	A	314	GLY	2.2
1	A	592	GLY	2.2
1	A	556	ARG	2.1
1	A	593	HIS	2.1
1	A	312	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	A	1	14/15	0.38	0.25	88,94,96,96	0
2	NAG	A	11	14/15	0.73	0.14	79,83,85,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.