



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:06 AM UTC

PDB ID : 4ZB4 / pdb_00004zb4
Title : Spliceosome component
Authors : Rocha de Moura, T.; Pena, P.
Deposited on : 2015-04-14
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	
1	B	360	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5665 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-processing factor 19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	354	Total	C	N	O	S	0	0	0
			2811	1784	461	552	14			
1	B	342	Total	C	N	O	S	0	0	0
			2723	1727	446	536	14			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	77	Total	O	0	0
			77	77		
2	B	54	Total	O	0	0
			54	54		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pre-mRNA-processing factor 19



• Molecule 1: Pre-mRNA-processing factor 19



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.22Å 100.14Å 65.34Å 90.00° 92.33° 90.00°	Depositor
Resolution (Å)	48.99 – 2.30 48.99 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.3 (48.99-2.30) 99.3 (48.99-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 2.29Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.196 , 0.239 0.198 , 0.242	Depositor DCC
R_{free} test set	1626 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.429	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.039 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5665	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/2869	0.97	16/3889 (0.4%)
1	B	0.38	0/2774	0.96	11/3756 (0.3%)
All	All	0.37	0/5643	0.96	27/7645 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	B	0	4
All	All	0	9

There are no bond length outliers.

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	329	SER	O-C-N	-18.11	98.50	122.59
1	B	472	ALA	CA-C-N	-15.55	101.07	122.99
1	B	472	ALA	C-N-CA	-15.55	101.07	122.99
1	A	418	GLU	CA-C-N	-14.79	100.21	122.09
1	A	418	GLU	C-N-CA	-14.79	100.21	122.09
1	B	467	LEU	CA-C-N	-13.52	104.31	122.16
1	B	467	LEU	C-N-CA	-13.52	104.31	122.16
1	A	330	ASP	N-CA-CB	-13.12	88.32	110.49
1	B	441	GLU	O-C-N	-12.08	106.53	122.59
1	B	440	ASN	CA-C-N	11.33	143.18	121.54
1	B	440	ASN	C-N-CA	11.33	143.18	121.54
1	A	419	PHE	CA-C-N	-9.78	108.78	122.87
1	A	419	PHE	C-N-CA	-9.78	108.78	122.87
1	B	461	GLU	CA-C-N	-8.17	109.34	122.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	461	GLU	C-N-CA	-8.17	109.34	122.66
1	A	420	LYS	CA-C-N	-7.84	107.58	121.70
1	A	420	LYS	C-N-CA	-7.84	107.58	121.70
1	B	217	ASP	O-C-N	-7.42	111.13	123.00
1	B	473	ASP	O-C-N	7.14	131.80	123.30
1	A	420	LYS	N-CA-C	6.78	118.83	109.54
1	A	418	GLU	O-C-N	-6.35	112.84	123.00
1	A	417	PRO	CA-C-N	6.25	132.95	121.70
1	A	417	PRO	C-N-CA	6.25	132.95	121.70
1	A	423	THR	N-CA-C	5.78	118.58	109.39
1	A	327	ALA	CA-C-N	5.13	131.34	121.54
1	A	327	ALA	C-N-CA	5.13	131.34	121.54
1	A	328	LYS	O-C-N	5.13	129.41	122.59

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	329	SER	Mainchain
1	A	417	PRO	Peptide
1	A	420	LYS	Mainchain,Peptide
1	A	421	THR	Mainchain
1	B	440	ASN	Peptide
1	B	441	GLU	Mainchain,Peptide
1	B	473	ASP	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2811	0	2753	27	0
1	B	2723	0	2651	54	0
2	A	77	0	0	1	0
2	B	54	0	0	3	0
All	All	5665	0	5404	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:440:ASN:ND2	1:B:473:ASP:OD1	1.70	1.24
1:B:440:ASN:ND2	1:B:473:ASP:CG	1.97	1.23
1:B:468:GLN:O	1:B:469:SER:C	1.87	1.15
1:B:455:LYS:O	1:B:455:LYS:HG3	1.48	1.09
1:B:468:GLN:O	1:B:469:SER:O	1.72	1.06
1:A:329:SER:O	1:A:330:ASP:C	1.95	1.05
1:A:419:PHE:O	1:A:420:LYS:HD3	1.59	1.03
1:B:470:ASP:HB3	1:B:472:ALA:N	1.76	1.00
1:A:329:SER:O	1:A:331:VAL:N	1.93	1.00
1:B:440:ASN:ND2	1:B:473:ASP:OD2	1.99	0.95
1:B:468:GLN:O	1:B:470:ASP:OD1	1.88	0.92
1:B:439:SER:OG	1:B:441:GLU:HB2	1.71	0.90
1:A:419:PHE:C	1:A:420:LYS:HD3	2.00	0.87
1:B:224:GLN:OE1	1:B:251:ARG:NH1	2.10	0.84
1:B:455:LYS:O	1:B:455:LYS:CG	2.29	0.81
1:B:251:ARG:NH2	2:B:602:HOH:O	2.15	0.79
1:B:440:ASN:CG	1:B:473:ASP:OD2	2.31	0.73
1:B:217:ASP:O	1:B:219:ALA:N	2.22	0.72
1:B:470:ASP:CB	1:B:472:ALA:N	2.54	0.68
1:B:174:TRP:CE3	1:B:449:LYS:NZ	2.62	0.67
1:B:440:ASN:CG	1:B:473:ASP:CG	2.65	0.65
1:B:462:GLU:N	1:B:462:GLU:OE1	2.29	0.65
1:A:225:LEU:HD21	1:A:489:ILE:HD11	1.77	0.65
1:B:184:GLN:NE2	1:B:186:GLN:HG3	2.13	0.64
1:B:190:ARG:HH11	1:B:190:ARG:HG3	1.63	0.63
1:B:217:ASP:O	1:B:218:GLY:C	2.39	0.62
1:A:419:PHE:O	1:A:420:LYS:CD	2.41	0.62
1:B:229:LYS:NZ	2:B:601:HOH:O	2.12	0.61
1:A:321:GLN:HG3	1:B:403:ARG:HG2	1.81	0.60
1:A:240:ARG:NH2	1:A:265:GLN:OE1	2.35	0.59
1:A:420:LYS:C	1:A:422:GLY:HA3	2.29	0.56
1:B:222:PHE:HE1	1:B:249:ILE:HG21	1.70	0.56
1:A:180:LEU:HD23	1:A:465:LEU:HB3	1.90	0.54
1:B:180:LEU:HD23	1:B:465:LEU:HB3	1.88	0.54
1:A:328:LYS:HD2	1:A:357:TYR:OH	2.08	0.54
1:B:288:ILE:HA	1:B:306:ASP:HA	1.89	0.53
1:A:394:ASP:O	1:A:419:PHE:CZ	2.62	0.53
1:B:190:ARG:HG3	1:B:190:ARG:NH1	2.25	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:184:GLN:HE22	1:B:186:GLN:CD	2.14	0.52
1:A:420:LYS:HA	1:A:421:THR:CG2	2.37	0.52
1:A:288:ILE:HA	1:A:306:ASP:HA	1.93	0.51
1:B:163:PHE:HZ	1:B:206:TYR:CE1	2.29	0.51
1:A:240:ARG:NH1	1:A:244:GLU:OE1	2.44	0.50
1:A:265:GLN:HG2	1:A:281:GLU:HG2	1.94	0.50
1:B:451:ASP:OD1	1:B:454:THR:N	2.45	0.50
1:B:240:ARG:HB3	1:B:244:GLU:OE1	2.12	0.49
1:B:424:VAL:N	2:B:605:HOH:O	2.45	0.49
1:B:270:ASP:OD1	1:B:272:LYS:HG3	2.12	0.49
1:B:469:SER:O	1:B:470:ASP:OD1	2.31	0.48
1:A:427:ASP:HB3	1:A:479:VAL:HG23	1.95	0.48
1:B:172:PRO:HD3	1:B:205:TYR:CE1	2.48	0.48
1:B:405:ASP:OD1	1:B:406:VAL:N	2.47	0.47
1:B:240:ARG:NH2	1:B:265:GLN:OE1	2.47	0.47
1:B:438:TYR:HE1	1:B:467:LEU:HD12	1.79	0.47
1:A:383:ALA:HB3	1:A:388:TRP:HB2	1.97	0.47
1:B:427:ASP:HB3	1:B:479:VAL:HG23	1.98	0.46
1:A:147:ILE:HG13	1:B:358:ASN:HB2	1.98	0.45
1:B:169:LEU:HD13	1:B:206:TYR:HB2	1.97	0.45
1:A:419:PHE:HZ	2:A:663:HOH:O	1.98	0.45
1:A:157:LEU:O	1:A:161:ARG:HD3	2.17	0.45
1:B:378:LYS:HE3	1:B:392:GLU:HB3	1.98	0.44
1:A:308:ARG:HA	1:A:331:VAL:CG2	2.48	0.44
1:A:423:THR:HG22	1:A:440:ASN:HD22	1.83	0.44
1:B:428:ILE:HD11	1:B:435:MET:HE2	2.00	0.43
1:B:198:LYS:NZ	1:B:430:ASP:OD2	2.46	0.42
1:B:469:SER:C	1:B:470:ASP:OD1	2.62	0.42
1:B:476:ASP:OD1	1:B:477:MET:N	2.52	0.42
1:B:174:TRP:CB	1:B:449:LYS:HZ1	2.32	0.42
1:A:417:PRO:HA	1:A:418:GLU:HA	1.74	0.42
1:B:190:ARG:HD2	1:B:493:ASN:HA	2.01	0.42
1:B:182:LEU:HD12	1:B:182:LEU:C	2.44	0.41
1:A:182:LEU:N	1:A:182:LEU:CD2	2.83	0.41
1:A:493:ASN:O	1:A:493:ASN:OD1	2.39	0.41
1:B:222:PHE:O	1:B:233:THR:HA	2.20	0.41
1:B:439:SER:HG	1:B:441:GLU:HB2	1.76	0.41
1:B:174:TRP:CD2	1:B:449:LYS:NZ	2.89	0.41
1:A:421:THR:O	1:A:421:THR:OG1	2.39	0.40
1:B:182:LEU:HD12	1:B:183:LEU:N	2.36	0.40
1:B:197:TYR:CD1	1:B:251:ARG:HB3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	348/360 (97%)	330 (95%)	16 (5%)	2 (1%)	21	27
1	B	326/360 (91%)	306 (94%)	18 (6%)	2 (1%)	21	27
All	All	674/720 (94%)	636 (94%)	34 (5%)	4 (1%)	21	27

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	329	SER
1	B	441	GLU
1	B	469	SER
1	A	328	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/320 (98%)	311 (99%)	3 (1%)	68	82
1	B	305/320 (95%)	295 (97%)	10 (3%)	33	50
All	All	619/640 (97%)	606 (98%)	13 (2%)	47	66

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	182	LEU
1	A	289	ILE
1	A	331	VAL
1	B	149	ARG
1	B	173	LYS
1	B	182	LEU
1	B	190	ARG
1	B	244	GLU
1	B	272	LYS
1	B	332	GLU
1	B	397	VAL
1	B	455	LYS
1	B	459	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	224	GLN
1	A	295	ASN
1	B	184	GLN
1	B	186	GLN
1	B	187	ASN
1	B	497	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/360 (98%)	0.25	25 (7%) 22 24	22, 40, 78, 203	0
1	B	342/360 (95%)	0.37	27 (7%) 18 20	24, 48, 90, 217	0
All	All	696/720 (96%)	0.31	52 (7%) 20 22	22, 43, 86, 217	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	217	ASP	5.9
1	A	183	LEU	5.1
1	A	472	ALA	4.8
1	A	327	ALA	4.7
1	A	419	PHE	4.7
1	B	472	ALA	4.0
1	A	329	SER	3.9
1	A	145	VAL	3.7
1	A	331	VAL	3.7
1	B	375	ALA	3.6
1	A	241	THR	3.5
1	B	206	TYR	3.5
1	A	144	ALA	3.4
1	A	421	THR	3.4
1	B	470	ASP	3.3
1	A	418	GLU	3.3
1	B	145	VAL	3.3
1	B	464	ALA	3.2
1	A	467	LEU	3.2
1	A	182	LEU	3.1
1	B	183	LEU	3.1
1	B	218	GLY	3.0
1	A	184	GLN	3.0
1	A	217	ASP	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	473	ASP	3.0
1	A	420	LYS	3.0
1	B	442	SER	2.9
1	A	375	ALA	2.9
1	A	473	ASP	2.9
1	B	330	ASP	2.9
1	B	462	GLU	2.8
1	A	372	ASP	2.6
1	B	463	SER	2.6
1	B	144	ALA	2.5
1	B	272	LYS	2.4
1	B	184	GLN	2.4
1	B	283	ASP	2.4
1	A	464	ALA	2.4
1	B	185	ALA	2.4
1	B	469	SER	2.4
1	B	493	ASN	2.3
1	A	422	GLY	2.3
1	B	285	ALA	2.3
1	B	235	THR	2.2
1	A	147	ILE	2.2
1	A	328	LYS	2.2
1	A	417	PRO	2.2
1	B	424	VAL	2.1
1	B	182	LEU	2.1
1	B	371	VAL	2.1
1	B	244	GLU	2.1
1	A	453	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.