



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:57 PM UTC

PDB ID : 6ZC9 / pdb_00006zc9
Title : Structure of 14-3-3 gamma in complex with Nedd4-2 14-3-3 binding motif Ser448
Authors : Pohl, P.; Kalabova, D.; Obsil, T.; Obsilova, V.
Deposited on : 2020-06-10
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	234	6% 93% 6%
1	B	234	6% 92%
1	C	234	6% 94% 5%
1	D	234	12% 91% 8%
2	E	10	20% 40% 30% 30%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	10	10%40%10%50%
2	G	10	20%40%10%50%
2	H	10	40%60%10%30%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7979 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 14-3-3 protein gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	S	0	0	0
			1809	1131	313	356	9			
1	B	225	Total	C	N	O	S	0	0	0
			1783	1117	307	351	8			
1	C	230	Total	C	N	O	S	0	0	0
			1819	1137	313	360	9			
1	D	232	Total	C	N	O	S	0	0	0
			1802	1128	314	351	9			

- Molecule 2 is a protein called E3 ubiquitin-protein ligase NEDD4-like.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	7	Total	C	N	O	P	0	0	0
			55	31	10	13	1			
2	F	5	Total	C	N	O	P	0	0	0
			41	21	8	11	1			
2	G	5	Total	C	N	O	P	0	0	0
			35	18	5	11	1			
2	H	7	Total	C	N	O	P	0	0	0
			49	28	7	13	1			

- Molecule 3 is 1,1,1,3,3,3-hexafluoropropan-2-ol (CCD ID: CFH) (formula: C₃H₂F₆O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	F	O	0	0
			10	3	6	1		
3	B	1	Total	C	F	O	0	0
			10	3	6	1		
3	D	1	Total	C	F	O	0	0
			10	3	6	1		
3	D	1	Total	C	F	O	0	0
			10	3	6	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	144	Total	O	0	0
			144	144		
4	B	156	Total	O	0	0
			156	156		
4	C	119	Total	O	0	0
			119	119		
4	D	113	Total	O	0	0
			113	113		
4	E	4	Total	O	0	0
			4	4		
4	F	5	Total	O	0	0
			5	5		
4	G	2	Total	O	0	0
			2	2		
4	H	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 14-3-3 protein gamma




- Molecule 1: 14-3-3 protein gamma



- Molecule 1: 14-3-3 protein gamma




- Molecule 1: 14-3-3 protein gamma





- Molecule 2: E3 ubiquitin-protein ligase NEDD4-like








● Molecule 2: E3 ubiquitin-protein ligase NEDD4-like



● Molecule 2: E3 ubiquitin-protein ligase NEDD4-like



● Molecule 2: E3 ubiquitin-protein ligase NEDD4-like



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	205.71 Å 205.71 Å 74.65 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.77 – 1.90 25.77 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (25.77-1.90) 99.8 (25.77-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.01 (at 1.89 Å)	Xtriage
Refinement program	PHENIX 1.12_2829, PHENIX 1.12_2829	Depositor
R, R_{free}	0.212 , 0.239 0.214 , 0.241	Depositor DCC
R_{free} test set	2100 reflections (2.26%)	wwPDB-VP
Wilson B-factor (Å ²)	39.6	Xtriage
Anisotropy	0.396	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.001 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7979	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CFH, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.17	0/1835	0.37	0/2478
1	B	0.16	0/1810	0.36	0/2449
1	C	0.18	0/1846	0.38	0/2497
1	D	0.18	0/1829	0.37	0/2475
2	E	0.21	0/45	0.43	0/58
2	F	0.18	0/29	0.37	0/35
2	G	0.12	0/23	0.22	0/28
2	H	0.14	0/39	0.28	0/51
All	All	0.17	0/7456	0.37	0/10071

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1809	0	1746	8	0
1	B	1783	0	1722	7	0
1	C	1819	0	1753	16	0
1	D	1802	0	1728	13	0
2	E	55	0	51	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	41	0	36	0	0
2	G	35	0	25	0	0
2	H	49	0	40	0	0
3	B	20	0	4	0	0
3	D	20	0	4	0	0
4	A	144	0	0	0	0
4	B	156	0	0	0	0
4	C	119	0	0	1	0
4	D	113	0	0	0	0
4	E	4	0	0	0	0
4	F	5	0	0	0	0
4	G	2	0	0	0	0
4	H	3	0	0	0	0
All	All	7979	0	7109	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:205:ILE:HA	1:C:208:LEU:CD1	1.79	1.10
1:C:205:ILE:HA	1:C:208:LEU:HD12	1.42	0.99
1:C:205:ILE:CA	1:C:208:LEU:HD12	2.16	0.75
1:C:205:ILE:HG21	1:D:66:ILE:HG12	1.75	0.68
1:C:208:LEU:HD21	1:C:223:MET:SD	2.34	0.67
1:A:13:LEU:HD11	1:C:82:VAL:HG22	1.80	0.63
1:C:205:ILE:HA	1:C:208:LEU:HD11	1.81	0.60
1:D:205:ILE:HA	1:D:208:LEU:HG	1.85	0.57
1:C:75:ASN:ND2	4:C:302:HOH:O	2.38	0.57
1:D:208:LEU:HD21	1:D:223:MET:HE1	1.86	0.56
1:B:82:VAL:HG22	1:D:13:LEU:HD11	1.87	0.56
1:A:66:ILE:HG12	1:B:205:ILE:HG21	1.91	0.52
1:D:165:MET:HE1	1:D:171:ILE:HB	1.91	0.51
1:C:165:MET:HE1	1:C:171:ILE:HB	1.93	0.51
1:B:234:THR:HG22	2:E:444:PRO:HB2	1.94	0.50
1:C:208:LEU:HD11	1:C:223:MET:HE1	1.93	0.50
1:D:64:SER:O	1:D:68:GLN:HG2	2.10	0.50
1:D:100:VAL:HG21	1:D:130:TYR:CD2	2.48	0.49
1:B:151:GLU:OE1	1:B:187:GLN:NE2	2.45	0.49
1:D:100:VAL:HG21	1:D:130:TYR:CG	2.48	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:ILE:HA	1:A:208:LEU:HG	1.95	0.47
1:C:211:LEU:HD23	1:C:215:SER:HB2	1.99	0.45
1:C:205:ILE:CA	1:C:208:LEU:CD1	2.71	0.44
1:C:204:ALA:O	1:C:208:LEU:HD12	2.18	0.44
1:A:168:THR:O	1:A:219:SER:OG	2.26	0.44
1:D:28:LYS:HG3	1:D:103:LEU:HD11	2.01	0.43
1:B:165:MET:HE1	1:B:171:ILE:HB	2.01	0.43
1:A:165:MET:HE1	1:A:171:ILE:HB	2.00	0.43
1:B:50:LYS:HB3	1:B:50:LYS:HE2	1.59	0.43
1:D:80:GLU:OE1	1:D:83:ARG:NH2	2.40	0.43
1:C:166:GLN:H	1:C:166:GLN:HG3	1.61	0.42
1:B:28:LYS:NZ	1:B:99:ASP:OD1	2.46	0.42
1:D:23:MET:HG2	1:D:48:ALA:HB2	2.01	0.42
1:A:208:LEU:HD21	1:A:223:MET:HE1	2.02	0.41
1:D:97:CYS:O	1:D:100:VAL:HG22	2.21	0.41
1:C:211:LEU:HD23	1:C:215:SER:CB	2.51	0.41
1:C:208:LEU:HA	1:C:211:LEU:CD1	2.50	0.41
1:A:112:CYS:HB3	1:A:120:LYS:HG2	2.02	0.40
1:D:77:LYS:HB2	1:D:77:LYS:HE2	1.89	0.40
2:E:444:PRO:HB2	2:E:445:ARG:H	1.75	0.40
1:A:12:ARG:HH21	1:A:40:GLU:CD	2.30	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/234 (97%)	223 (99%)	3 (1%)	0	100	100
1	B	221/234 (94%)	219 (99%)	2 (1%)	0	100	100
1	C	226/234 (97%)	223 (99%)	3 (1%)	0	100	100
1	D	230/234 (98%)	228 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	4/10 (40%)	4 (100%)	0	0	100	100
2	F	2/10 (20%)	2 (100%)	0	0	100	100
2	G	2/10 (20%)	2 (100%)	0	0	100	100
2	H	4/10 (40%)	4 (100%)	0	0	100	100
All	All	915/976 (94%)	905 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/206 (91%)	187 (100%)	0	100	100
1	B	187/206 (91%)	187 (100%)	0	100	100
1	C	191/206 (93%)	191 (100%)	0	100	100
1	D	183/206 (89%)	183 (100%)	0	100	100
2	E	6/9 (67%)	6 (100%)	0	100	100
2	F	4/9 (44%)	4 (100%)	0	100	100
2	G	3/9 (33%)	3 (100%)	0	100	100
2	H	5/9 (56%)	5 (100%)	0	100	100
All	All	766/860 (89%)	766 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	164	HIS
1	B	16	GLN
1	B	166	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	16	GLN
1	C	98	GLN
1	D	16	GLN
1	D	68	GLN
1	D	164	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SEP	E	448	2	8,9,10	1.50	1 (12%)	7,12,14	1.02	0
2	SEP	H	448	2	8,9,10	1.54	1 (12%)	7,12,14	0.80	0
2	SEP	G	448	2	8,9,10	1.59	1 (12%)	7,12,14	1.31	1 (14%)
2	SEP	F	448	2	8,9,10	1.55	1 (12%)	7,12,14	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	E	448	2	-	0/6/8/10	-
2	SEP	H	448	2	-	0/6/8/10	-
2	SEP	G	448	2	-	0/6/8/10	-
2	SEP	F	448	2	-	0/6/8/10	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	448	SEP	P-O1P	3.43	1.61	1.50
2	G	448	SEP	P-O1P	3.41	1.61	1.50
2	F	448	SEP	P-O1P	3.38	1.61	1.50
2	E	448	SEP	P-O1P	3.28	1.60	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	448	SEP	OG-CB-CA	2.44	110.52	108.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CFH	D	301	-	9,9,9	1.15	0	15,15,15	0.80	0
3	CFH	D	302	-	9,9,9	1.15	0	15,15,15	0.76	0
3	CFH	B	301	-	9,9,9	1.18	0	15,15,15	0.89	0
3	CFH	B	302	-	9,9,9	1.16	0	15,15,15	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CFH	D	301	-	-	6/12/12/12	-
3	CFH	D	302	-	-	6/12/12/12	-
3	CFH	B	301	-	-	0/12/12/12	-
3	CFH	B	302	-	-	0/12/12/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	301	CFH	O4-C2-C3-F10
3	D	301	CFH	C1-C2-C3-F10
3	D	301	CFH	C1-C2-C3-F8
3	D	301	CFH	C1-C2-C3-F9
3	D	301	CFH	O4-C2-C3-F8
3	D	301	CFH	O4-C2-C3-F9
3	D	302	CFH	O4-C2-C3-F9
3	D	302	CFH	O4-C2-C3-F8
3	D	302	CFH	C1-C2-C3-F10
3	D	302	CFH	O4-C2-C3-F10
3	D	302	CFH	C1-C2-C3-F8
3	D	302	CFH	C1-C2-C3-F9

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/234 (98%)	0.58	15 (6%) 25 26	31, 41, 62, 96	0
1	B	225/234 (96%)	0.49	14 (6%) 26 28	30, 38, 54, 74	0
1	C	230/234 (98%)	0.73	15 (6%) 25 26	32, 44, 61, 74	0
1	D	232/234 (99%)	0.92	27 (11%) 9 10	34, 47, 62, 84	0
2	E	6/10 (60%)	1.72	2 (33%) 1 1	38, 42, 57, 63	0
2	F	4/10 (40%)	1.12	1 (25%) 2 1	40, 44, 48, 64	0
2	G	4/10 (40%)	1.83	2 (50%) 0 0	44, 51, 55, 65	0
2	H	6/10 (60%)	2.34	4 (66%) 0 0	45, 52, 65, 66	0
All	All	937/976 (96%)	0.70	80 (8%) 16 17	30, 43, 62, 96	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	208	LEU	5.3
1	C	211	LEU	5.1
1	D	211	LEU	4.7
1	A	72	ALA	4.6
1	C	79	ILE	4.5
1	B	70	THR	4.5
2	E	450	PRO	4.3
1	A	212	ASN	4.3
1	D	216	TYR	4.2
1	A	73	ASP	4.2
1	C	206	ALA	4.1
1	A	215	SER	4.1
2	H	444	PRO	3.9
1	B	79	ILE	3.8
1	D	115	THR	3.7
1	D	210	THR	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	444	PRO	3.5
1	A	214	ASP	3.4
1	B	115	THR	3.4
1	C	73	ASP	3.4
1	C	205	ILE	3.3
1	D	139	THR	3.3
2	H	450	PRO	3.3
1	D	212	ASN	3.2
1	D	33	LEU	3.1
2	H	447	LEU	3.1
1	B	206	ALA	3.1
1	D	233	TRP	3.0
2	G	445	ARG	2.9
1	B	78	LYS	2.9
1	C	234	THR	2.9
1	D	214	ASP	2.9
1	A	213	GLU	2.9
1	B	208	LEU	2.8
1	C	70	THR	2.8
1	D	205	ILE	2.8
1	D	183	TYR	2.7
1	D	202	ASP	2.7
1	D	232	LEU	2.7
1	B	214	ASP	2.6
1	B	234	THR	2.6
1	D	213	GLU	2.6
1	A	211	LEU	2.6
1	A	79	ILE	2.6
1	D	185	GLU	2.5
1	A	69	LYS	2.5
2	H	445	ARG	2.5
1	B	81	MET	2.5
1	D	184	TYR	2.5
1	D	209	ASP	2.5
1	D	39	ASN	2.5
1	D	215	SER	2.5
1	D	208	LEU	2.4
1	A	3	ASP	2.3
1	A	83	ARG	2.3
1	B	68	GLN	2.3
1	D	70	THR	2.3
1	D	61	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	7	LEU	2.3
1	D	168	THR	2.3
2	G	447	LEU	2.2
2	F	449	SER	2.2
1	A	216	TYR	2.2
1	C	171	ILE	2.2
1	A	234	THR	2.1
1	D	234	THR	2.1
1	C	68	GLN	2.1
1	B	66	ILE	2.1
1	C	216	TYR	2.1
1	C	215	SER	2.1
1	D	114	GLU	2.1
1	B	211	LEU	2.1
1	A	183	TYR	2.1
1	B	69	LYS	2.1
1	C	81	MET	2.1
1	C	212	ASN	2.1
1	D	144	ALA	2.1
1	B	205	ILE	2.0
1	C	69	LYS	2.0
1	D	98	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SEP	H	448	10/11	0.96	0.08	36,38,43,44	0
2	SEP	F	448	10/11	0.98	0.05	31,34,39,42	0
2	SEP	G	448	10/11	0.98	0.05	36,40,44,46	0
2	SEP	E	448	10/11	0.98	0.06	26,32,36,37	0

6.3 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CFH	D	302	10/10	0.81	0.13	56,63,69,75	0
3	CFH	D	301	10/10	0.87	0.11	54,63,65,67	0
3	CFH	B	302	10/10	0.92	0.08	41,49,52,53	0
3	CFH	B	301	10/10	0.93	0.08	46,52,55,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.