



wwPDB EM Validation Summary Report ⓘ

Mar 30, 2026 – 11:36 AM UTC

PDB ID : 6ZCE / pdb_00006zce
EMDB ID : EMD-11160
Title : Structure of a yeast ABCE1-bound 43S pre-initiation complex
Authors : Kratzat, H.; Mackens-Kiani, T.; Cheng, J.; Berninghausen, O.; Becker, T.; Beckmann, R.
Deposited on : 2020-06-10
Resolution : 5.30 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY



The reported resolution of this entry is 5.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
RNA backbone	8273	3508	-
Q-score	-	25397	625 (4.80 - 5.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	l	347	
2	r	274	
3	A	1800	
4	B	252	






Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	C	255	
6	D	254	
7	E	240	
8	F	261	
9	G	225	
10	H	236	
11	I	190	
12	J	200	
13	K	197	
14	L	105	
15	M	155	
16	N	143	
17	O	151	
18	P	136	
19	Q	141	
20	R	143	
21	S	136	
22	T	146	
23	U	144	
24	V	121	
25	W	87	
26	X	130	
27	Y	145	
28	Z	135	
29	a	108	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
30	b	119	
31	c	82	
32	d	67	
33	e	56	
34	f	63	
35	g	152	
36	h	319	
37	o	964	
38	p	763	
39	q	812	
40	i	153	
41	m	108	
42	s	265	
42	t	265	
43	j	77	
44	k	608	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
46	ADP	k	701	-	-	X	-

2 Entry composition

There are 49 unique types of molecules in this entry. The entry contains 76380 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Eukaryotic translation initiation factor 3 subunit I.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	l	330	Total	C	N	O	0	0
			1624	964	330	330		

- Molecule 2 is a protein called Eukaryotic translation initiation factor 3 subunit G.

Mol	Chain	Residues	Atoms				AltConf	Trace
2	r	53	Total	C	N	O	0	0
			261	155	53	53		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
r	139	SER	ASN	conflict	UNP Q04067

- Molecule 3 is a RNA chain called 18S ribosomal RNA (1719-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
3	A	1719	Total	C	N	O	P	0	0
			36643	16382	6499	12043	1719		

- Molecule 4 is a protein called 40S ribosomal protein S0-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
4	B	206	Total	C	N	O	0	0
			1020	608	206	206		

- Molecule 5 is a protein called 40S ribosomal protein S1-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
5	C	214	Total	C	N	O	0	0
			1061	633	214	214		

- Molecule 6 is a protein called 40S ribosomal protein S2.

Mol	Chain	Residues	Atoms				AltConf	Trace
6	D	217	Total	C	N	O	0	0
			1063	629	217	217		

- Molecule 7 is a protein called 40S ribosomal protein S3.

Mol	Chain	Residues	Atoms				AltConf	Trace
7	E	223	Total	C	N	O	0	0
			1098	652	223	223		

- Molecule 8 is a protein called 40S ribosomal protein S4-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
8	F	260	Total	C	N	O	0	0
			1276	756	260	260		

- Molecule 9 is a protein called 40S ribosomal protein S5.

Mol	Chain	Residues	Atoms				AltConf	Trace
9	G	206	Total	C	N	O	0	0
			1020	608	206	206		

- Molecule 10 is a protein called 40S ribosomal protein S6-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
10	H	226	Total	C	N	O	0	0
			1113	661	226	226		

- Molecule 11 is a protein called 40S ribosomal protein S7-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
11	I	184	Total	C	N	O	0	0
			913	545	184	184		

- Molecule 12 is a protein called 40S ribosomal protein S8-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
12	J	188	Total	C	N	O	0	0
			924	548	188	188		

- Molecule 13 is a protein called 40S ribosomal protein S9-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
13	K	184	Total	C	N	O	0	0
			910	542	184	184		

- Molecule 14 is a protein called 40S ribosomal protein S10-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
14	L	92	Total	C	N	O	0	0
			456	272	92	92		

- Molecule 15 is a protein called 40S ribosomal protein S11-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
15	M	142	Total	C	N	O	0	0
			702	418	142	142		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	147	ALA	GLY	conflict	UNP P0CX47

- Molecule 16 is a protein called 40S ribosomal protein S12.

Mol	Chain	Residues	Atoms				AltConf	Trace
16	N	120	Total	C	N	O	0	0
			590	350	120	120		

- Molecule 17 is a protein called 40S ribosomal protein S13.

Mol	Chain	Residues	Atoms				AltConf	Trace
17	O	150	Total	C	N	O	0	0
			742	442	150	150		

- Molecule 18 is a protein called 40S ribosomal protein S14-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
18	P	127	Total	C	N	O	0	0
			620	366	127	127		

- Molecule 19 is a protein called 40S ribosomal protein S15.

Mol	Chain	Residues	Atoms				AltConf	Trace
19	Q	122	Total	C	N	O	0	0
			601	357	122	122		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Q	137	SER	ARG	conflict	UNP Q01855

- Molecule 20 is a protein called 40S ribosomal protein S16-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
20	R	141	Total	C	N	O	0	0
			693	411	141	141		

- Molecule 21 is a protein called 40S ribosomal protein S17-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
21	S	117	Total	C	N	O	0	0
			579	345	117	117		

- Molecule 22 is a protein called 40S ribosomal protein S18-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
22	T	145	Total	C	N	O	0	0
			715	425	145	145		

- Molecule 23 is a protein called 40S ribosomal protein S19-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
23	U	143	Total	C	N	O	0	0
			700	414	143	143		

- Molecule 24 is a protein called 40S ribosomal protein S20.

Mol	Chain	Residues	Atoms				AltConf	Trace
24	V	105	Total	C	N	O	0	0
			521	311	105	105		

- Molecule 25 is a protein called 40S ribosomal protein S21-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
25	W	87	Total	C	N	O	0	0
			429	255	87	87		

- Molecule 26 is a protein called 40S ribosomal protein S22-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
26	X	129	Total	C	N	O	0	0
			634	376	129	129		

- Molecule 27 is a protein called 40S ribosomal protein S23-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
27	Y	144	Total	C	N	O	0	0
			704	416	144	144		

- Molecule 28 is a protein called 40S ribosomal protein S24-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
28	Z	134	Total	C	N	O	0	0
			661	393	134	134		

- Molecule 29 is a protein called 40S ribosomal protein S25-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
29	a	70	Total	C	N	O	0	0
			347	207	70	70		

- Molecule 30 is a protein called 40S ribosomal protein S26-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
30	b	97	Total	C	N	O	0	0
			482	288	97	97		

- Molecule 31 is a protein called 40S ribosomal protein S27-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
31	c	81	Total	C	N	O	0	0
			400	238	81	81		

- Molecule 32 is a protein called 40S ribosomal protein S28-B.

Mol	Chain	Residues	Atoms				AltConf	Trace
32	d	62	Total	C	N	O	0	0
			305	181	62	62		

- Molecule 33 is a protein called 40S ribosomal protein S29-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
33	e	53	Total	C	N	O	0	0
			260	154	53	53		

- Molecule 34 is a protein called 40S ribosomal protein S30-A.

Mol	Chain	Residues	Atoms				AltConf	Trace
34	f	53	Total	C	N	O	0	0
			261	155	53	53		

- Molecule 35 is a protein called Ubiquitin-40S ribosomal protein S31.

Mol	Chain	Residues	Atoms				AltConf	Trace
35	g	71	Total	C	N	O	0	0
			351	209	71	71		

- Molecule 36 is a protein called Guanine nucleotide-binding protein subunit beta-like protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
36	h	318	Total	C	N	O	0	0
			1568	932	318	318		

- Molecule 37 is a protein called Eukaryotic translation initiation factor 3 subunit A.

Mol	Chain	Residues	Atoms				AltConf	Trace
37	o	529	Total	C	N	O	0	0
			2631	1573	529	529		

- Molecule 38 is a protein called Eukaryotic translation initiation factor 3 subunit B.

Mol	Chain	Residues	Atoms				AltConf	Trace
38	p	646	Total	C	N	O	0	0
			3201	1909	646	646		

- Molecule 39 is a protein called Eukaryotic translation initiation factor 3 subunit C.

Mol	Chain	Residues	Atoms				AltConf	Trace
39	q	636	Total	C	N	O	0	0
			3169	1897	636	636		

- Molecule 40 is a protein called Eukaryotic translation initiation factor 1A.

Mol	Chain	Residues	Atoms				AltConf	Trace
40	i	97	Total	C	N	O	0	0
			476	282	97	97		

- Molecule 41 is a protein called Eukaryotic translation initiation factor eIF-1.

Mol	Chain	Residues	Atoms				AltConf	Trace
41	m	90	Total	C	N	O	0	0
			443	263	90	90		

- Molecule 42 is a protein called Eukaryotic translation initiation factor 3 subunit J.

Mol	Chain	Residues	Atoms				AltConf	Trace
42	t	84	Total	C	N	O	0	0
			418	251	84	83		
42	s	95	Total	C	N	O	0	0
			470	280	95	95		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
t	254	ALA	PRO	conflict	UNP Q05775
s	254	ALA	PRO	conflict	UNP Q05775

- Molecule 43 is a protein called RNA recognition motif (unknown).

Mol	Chain	Residues	Atoms				AltConf	Trace
43	j	77	Total	C	N	O	0	0
			385	231	77	77		

- Molecule 44 is a protein called Translation initiation factor RLI1.

Mol	Chain	Residues	Atoms				AltConf	Trace
44	k	579	Total	C	N	O	0	0
			2860	1702	579	579		


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
k	350	VAL	ALA	conflict	UNP Q03195

- Molecule 45 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
45	A	1	Total	Zn	0
			1	1	
45	b	1	Total	Zn	0
			1	1	
45	c	1	Total	Zn	0
			1	1	
45	g	1	Total	Zn	0
			1	1	

- Molecule 46 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).




Mol	Chain	Residues	Atoms					AltConf
46	k	1	Total	C	N	O	P	0
			27	10	5	10	2	

- Molecule 47 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		AltConf
47	k	2	Total	Mg	0
			2	2	

- Molecule 48 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					AltConf
48	k	1	Total	C	N	O	P	0
			31	10	5	13	3	

- Molecule 49 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			AltConf
49	k	1	Total	Fe	S	0
			8	4	4	
49	k	1	Total	Fe	S	0
			8	4	4	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Eukaryotic translation initiation factor 3 subunit I




• Molecule 2: Eukaryotic translation initiation factor 3 subunit G



CYS	TYR	SER
PRO	ARG	SER
PHE	ASP	GLU
LYS	SER	GLU
THR	ARG	VAL
ILE	GLU	ALA
LEU	ARG	GLU
SER	ASP	GLN
GLU	ASP	ALA
LEU	MET	LEU
SER	CYS	THR
ALA	THR	LEU
LEU	LEU	LEU
GLU	LYS	ASP
GLU	ILE	GLY
PRO	MET	ASP
ALA	GLN	ALA
THR	VAL	THR
SER	ASN	MET
GLU	GLU	ASN
GLY	ASN	A26
VAL	ALA	U27
GLU	ASP	A28
GLU	GLU	U29
ALA	ASN	ARG
ALA	VAL	G30
ALA	SER	C31
SER	LEU	GLU
GLU	ARG	TRP
GLY	GLU	SER
LYS	GLU	LYS
LYS	PRO	PRO
ALA	LEU	LYS
GLY	LEU	VAL
VAL	PHE	GLN
GLY	PHE	VAL
GLY	ALA	GLY
ALA	ALA	GLY
ALA	PRO	PRO
GLY	ILE	ILE
SER	PRO	SER
ILE	ARG	THR
VAL	VAL	ARG
GLY	SER	GLY
GLY	GLY	LYS
ALA	ARG	ALA
ARG	GLY	ARG
ARG	LEU	ASP
ALA	ALA	PRO
GLY	PHE	SER
LYS	VAL	SER
ALA	THR	ASP
	PHE	ALA


• Molecule 3: 18S ribosomal RNA (1719-MER)

Chain A:  38% 45% 12% .







- Molecule 4: 40S ribosomal protein S0-A




- Molecule 5: 40S ribosomal protein S1-A





- Molecule 6: 40S ribosomal protein S2





- Molecule 7: 40S ribosomal protein S3





- Molecule 8: 40S ribosomal protein S4-A





- Molecule 9: 40S ribosomal protein S5





- Molecule 10: 40S ribosomal protein S6-A






- Molecule 11: 40S ribosomal protein S7-A




- Molecule 12: 40S ribosomal protein S8-A




Chain K:




- Molecule 18: 40S ribosomal protein S14-A




- Molecule 19: 40S ribosomal protein S15




- Molecule 20: 40S ribosomal protein S16-A




- Molecule 21: 40S ribosomal protein S17-A




- Molecule 22: 40S ribosomal protein S18-A




- Molecule 23: 40S ribosomal protein S19-A

Chain U: 





- Molecule 24: 40S ribosomal protein S20

Chain V: 




- Molecule 25: 40S ribosomal protein S21-A

Chain W: 



- Molecule 26: 40S ribosomal protein S22-A

Chain X: 



- Molecule 27: 40S ribosomal protein S23-A

Chain Y: 




- Molecule 28: 40S ribosomal protein S24-A


Chain Z: 




- Molecule 29: 40S ribosomal protein S25-A

Chain a: 







- Molecule 30: 40S ribosomal protein S26-A




- Molecule 31: 40S ribosomal protein S27-A




- Molecule 32: 40S ribosomal protein S28-B





- Molecule 33: 40S ribosomal protein S29-A




- Molecule 34: 40S ribosomal protein S30-A





- Molecule 35: Ubiquitin-40S ribosomal protein S31







• Molecule 36: Guanine nucleotide-binding protein subunit beta-like protein





• Molecule 37: Eukaryotic translation initiation factor 3 subunit A







• Molecule 39: Eukaryotic translation initiation factor 3 subunit C





• Molecule 40: Eukaryotic translation initiation factor 1A




- Molecule 43: RNA recognition motif (unknown)



- Molecule 44: Translation initiation factor RLI1





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	20618	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	2.5	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON II (4k x 4k)	Depositor
Maximum map value	0.054	Depositor
Minimum map value	-0.016	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.004	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	390.24, 390.24, 390.24	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.084, 1.084, 1.084	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SF4, ATP, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	l	0.49	0/1622	0.83	0/2252
2	r	0.41	0/260	0.81	0/360
3	A	0.38	2/40986 (0.0%)	0.50	7/63859 (0.0%)
4	B	0.30	0/1019	0.78	4/1419 (0.3%)
5	C	0.31	0/1060	0.79	0/1477
6	D	0.29	0/1062	0.74	1/1472 (0.1%)
7	E	0.31	0/1097	0.65	0/1524
8	F	0.29	0/1275	0.72	1/1769 (0.1%)
9	G	0.29	0/1019	0.67	0/1419
10	H	0.24	0/1112	0.59	0/1545
11	I	0.37	0/912	0.89	4/1271 (0.3%)
12	J	0.31	0/922	0.72	0/1278
13	K	0.28	0/908	0.69	0/1262
14	L	0.33	0/455	1.03	3/633 (0.5%)
15	M	0.35	0/701	0.82	1/975 (0.1%)
16	N	0.35	0/589	0.83	0/817
17	O	0.32	0/741	0.71	2/1031 (0.2%)
18	P	0.33	0/619	0.73	0/856
19	Q	0.47	1/600 (0.2%)	0.82	0/833
20	R	0.34	0/692	0.91	3/960 (0.3%)
21	S	0.30	0/577	0.73	0/801
22	T	0.30	0/714	0.82	5/992 (0.5%)
23	U	0.32	0/699	0.72	0/968
24	V	0.31	0/520	0.71	0/724
25	W	0.32	0/428	0.61	0/594
26	X	0.32	0/633	0.87	3/878 (0.3%)
27	Y	0.33	0/703	0.90	0/973
28	Z	0.26	0/660	0.77	3/917 (0.3%)
29	a	0.37	0/346	0.87	0/481
30	b	0.72	1/481 (0.2%)	0.88	2/670 (0.3%)
31	c	0.30	0/399	0.73	0/554
32	d	0.29	0/304	0.64	0/421

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
33	e	0.37	0/259	0.95	2/358 (0.6%)
34	f	0.33	0/260	0.64	0/360
35	g	0.45	0/350	1.27	5/486 (1.0%)
36	h	0.24	0/1567	0.58	1/2179 (0.0%)
37	o	0.71	0/2626	1.33	2/3660 (0.1%)
38	p	0.72	0/3197	1.05	2/4452 (0.0%)
39	q	0.84	0/3165	1.37	7/4418 (0.2%)
40	i	0.62	0/475	1.01	0/658
41	m	0.72	0/442	1.04	1/613 (0.2%)
42	s	0.90	0/469	1.67	8/652 (1.2%)
42	t	0.95	0/417	1.83	9/581 (1.5%)
43	j	0.36	0/384	0.66	0/536
44	k	0.94	2/2858 (0.1%)	1.50	6/3977 (0.2%)
All	All	0.47	6/80584 (0.0%)	0.77	82/118915 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	B	0	1
5	C	0	4
6	D	0	1
8	F	0	1
9	G	0	3
11	I	0	3
15	M	0	1
16	N	0	3
18	P	0	1
19	Q	0	2
20	R	0	3
21	S	0	1
22	T	0	2
26	X	0	1
27	Y	0	1
28	Z	0	2
29	a	0	2
30	b	0	2
31	c	0	1
35	g	0	4
36	h	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
38	p	0	2
All	All	0	42

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
30	b	59	TYR	C-N	13.47	1.65	1.33
19	Q	67	ALA	C-N	8.17	1.52	1.33
44	k	379	GLU	C-O	-6.50	1.16	1.23
44	k	379	GLU	C-N	5.97	1.42	1.33
3	A	649	U	O3'-P	5.87	1.70	1.61

The worst 5 of 82 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
42	t	175	SER	N-CA-C	10.81	124.21	111.02
20	R	40	GLU	CA-C-N	-10.80	106.34	119.84
20	R	40	GLU	C-N-CA	-10.80	106.34	119.84
14	L	87	VAL	CA-C-N	10.76	133.28	119.84
14	L	87	VAL	C-N-CA	10.76	133.28	119.84

There are no chirality outliers.

5 of 42 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	B	94	GLY	Peptide
5	C	131	ASP	Peptide
5	C	205	PHE	Peptide
5	C	206	PRO	Peptide
5	C	36	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	1624	0	727	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	r	261	0	122	1	0
3	A	36643	0	18433	860	0
4	B	1020	0	474	6	0
5	C	1061	0	473	10	0
6	D	1063	0	499	11	0
7	E	1098	0	525	16	0
8	F	1276	0	576	11	0
9	G	1020	0	482	11	0
10	H	1113	0	510	11	0
11	I	913	0	400	15	0
12	J	924	0	452	14	0
13	K	910	0	419	8	0
14	L	456	0	196	5	0
15	M	702	0	304	7	0
16	N	590	0	286	11	0
17	O	742	0	345	7	0
18	P	620	0	311	14	0
19	Q	601	0	277	6	0
20	R	693	0	323	12	0
21	S	579	0	246	2	0
22	T	715	0	318	19	0
23	U	700	0	334	11	0
24	V	521	0	217	3	0
25	W	429	0	201	2	0
26	X	634	0	289	3	0
27	Y	704	0	324	24	0
28	Z	661	0	312	5	0
29	a	347	0	158	15	0
30	b	482	0	223	8	0
31	c	400	0	180	9	0
32	d	305	0	133	7	0
33	e	260	0	112	2	0
34	f	261	0	113	8	0
35	g	351	0	158	65	0
36	h	1568	0	754	17	0
37	o	2631	0	1207	6	0
38	p	3201	0	1397	9	0
39	q	3169	0	1372	17	0
40	i	476	0	215	14	0
41	m	443	0	193	6	0
42	s	470	0	223	16	0
42	t	418	0	191	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
43	j	385	0	387	11	0
44	k	2860	0	1246	29	0
45	A	1	0	0	0	0
45	b	1	0	0	1	0
45	c	1	0	0	0	0
45	g	1	0	0	0	0
46	k	27	0	12	19	0
47	k	2	0	0	0	0
48	k	31	0	12	3	0
49	k	16	0	0	0	0
All	All	76380	0	36661	1090	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 1090 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1253:U:H4'	35:g:143:LYS:CA	1.27	1.55
3:A:1219:A:C2	42:s:224:ALA:HB1	1.50	1.44
34:f:9:ALA:HB2	40:i:83:ASP:CB	1.48	1.43
3:A:1756:A:N1	40:i:66:ARG:CA	1.79	1.34
3:A:321:C:N4	39:q:192:ALA:HB3	1.47	1.29

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	l	326/347 (94%)	318 (98%)	8 (2%)	0	100	100
2	r	51/274 (19%)	48 (94%)	3 (6%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	B	204/252 (81%)	174 (85%)	26 (13%)	4 (2%)	6	30
5	C	212/255 (83%)	162 (76%)	45 (21%)	5 (2%)	4	25
6	D	215/254 (85%)	194 (90%)	19 (9%)	2 (1%)	14	49
7	E	221/240 (92%)	202 (91%)	15 (7%)	4 (2%)	6	32
8	F	258/261 (99%)	226 (88%)	30 (12%)	2 (1%)	16	52
9	G	204/225 (91%)	178 (87%)	21 (10%)	5 (2%)	4	25
10	H	224/236 (95%)	209 (93%)	9 (4%)	6 (3%)	4	24
11	I	182/190 (96%)	153 (84%)	20 (11%)	9 (5%)	1	15
12	J	184/200 (92%)	160 (87%)	23 (12%)	1 (0%)	24	63
13	K	180/197 (91%)	160 (89%)	19 (11%)	1 (1%)	21	58
14	L	90/105 (86%)	77 (86%)	10 (11%)	3 (3%)	3	20
15	M	140/155 (90%)	128 (91%)	10 (7%)	2 (1%)	9	38
16	N	118/143 (82%)	85 (72%)	29 (25%)	4 (3%)	3	20
17	O	148/151 (98%)	134 (90%)	12 (8%)	2 (1%)	9	38
18	P	125/136 (92%)	112 (90%)	12 (10%)	1 (1%)	16	52
19	Q	120/141 (85%)	100 (83%)	17 (14%)	3 (2%)	4	25
20	R	139/143 (97%)	121 (87%)	14 (10%)	4 (3%)	3	22
21	S	113/136 (83%)	100 (88%)	12 (11%)	1 (1%)	14	49
22	T	143/146 (98%)	122 (85%)	18 (13%)	3 (2%)	5	28
23	U	141/144 (98%)	125 (89%)	16 (11%)	0	100	100
24	V	103/121 (85%)	98 (95%)	5 (5%)	0	100	100
25	W	85/87 (98%)	72 (85%)	11 (13%)	2 (2%)	4	25
26	X	127/130 (98%)	115 (91%)	11 (9%)	1 (1%)	16	52
27	Y	142/145 (98%)	119 (84%)	19 (13%)	4 (3%)	4	23
28	Z	132/135 (98%)	121 (92%)	8 (6%)	3 (2%)	5	26
29	a	68/108 (63%)	52 (76%)	14 (21%)	2 (3%)	3	22
30	b	95/119 (80%)	67 (70%)	17 (18%)	11 (12%)	0	5
31	c	79/82 (96%)	70 (89%)	8 (10%)	1 (1%)	9	40
32	d	60/67 (90%)	51 (85%)	9 (15%)	0	100	100
33	e	51/56 (91%)	49 (96%)	2 (4%)	0	100	100
34	f	51/63 (81%)	47 (92%)	2 (4%)	2 (4%)	2	18

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
35	g	69/152 (45%)	44 (64%)	18 (26%)	7 (10%)	0	7
36	h	316/319 (99%)	298 (94%)	18 (6%)	0	100	100
37	o	519/964 (54%)	483 (93%)	30 (6%)	6 (1%)	10	42
38	p	638/763 (84%)	583 (91%)	43 (7%)	12 (2%)	6	30
39	q	628/812 (77%)	568 (90%)	48 (8%)	12 (2%)	6	30
40	i	95/153 (62%)	82 (86%)	13 (14%)	0	100	100
41	m	88/108 (82%)	77 (88%)	8 (9%)	3 (3%)	3	20
42	s	93/265 (35%)	87 (94%)	5 (5%)	1 (1%)	11	45
42	t	82/265 (31%)	74 (90%)	7 (8%)	1 (1%)	10	42
43	j	75/77 (97%)	68 (91%)	5 (7%)	2 (3%)	4	24
44	k	575/608 (95%)	522 (91%)	47 (8%)	6 (1%)	12	47
All	All	7909/9930 (80%)	7035 (89%)	736 (9%)	138 (2%)	9	34

5 of 138 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	C	207	LEU
10	H	153	VAL
11	I	109	VAL
14	L	87	VAL
14	L	88	PRO

5.3.2 Protein sidechains ⓘ

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	A	1713/1800 (95%)	501 (29%)	35 (2%)

5 of 501 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	A	2	A
3	A	4	C

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	A	17	C
3	A	25	C
3	A	26	A

5 of 35 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
3	A	1250	U
3	A	1339	C
3	A	1493	A
3	A	507	U
3	A	417	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
49	SF4	k	705	-	0,12,12	-	-	-		
48	ATP	k	703	47	32,33,33	1.39	4 (12%)	48,52,52	1.77	9 (18%)
49	SF4	k	706	-	0,12,12	-	-	-		
46	ADP	k	701	47	28,29,29	1.40	4 (14%)	43,45,45	1.82	9 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
49	SF4	k	705	-	-	-	0/6/5/5
48	ATP	k	703	47	-	2/22/38/38	0/3/3/3
49	SF4	k	706	-	-	-	0/6/5/5
46	ADP	k	701	47	-	4/16/32/32	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
48	k	703	ATP	C5-C4	4.74	1.47	1.39
46	k	701	ADP	C5-C4	4.69	1.47	1.39
48	k	703	ATP	C5-C6	2.77	1.48	1.41
46	k	701	ADP	C5-C6	2.71	1.48	1.41
48	k	703	ATP	C8-N7	2.41	1.36	1.31

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
48	k	703	ATP	C5-C4-N3	-5.86	118.65	126.72
46	k	701	ADP	C5-C4-N3	-5.77	118.77	126.72
46	k	701	ADP	N3-C4-N9	4.59	134.98	127.17
48	k	703	ATP	N3-C4-N9	4.57	134.94	127.17
48	k	703	ATP	C2-N3-C4	3.75	120.99	111.83

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
46	k	701	ADP	C5'-O5'-PA-O1A
46	k	701	ADP	C5'-O5'-PA-O2A
46	k	701	ADP	C5'-O5'-PA-O3A
48	k	703	ATP	O4'-C4'-C5'-O5'
46	k	701	ADP	PA-O3A-PB-O2B


There are no ring outliers.

2 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
48	k	703	ATP	3	0
46	k	701	ADP	19	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
30	b	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	b	59:TYR	C	60:PRO	N	1.65


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-11160. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 175




Y Index: 159




Z Index: 189


6.3.2 Raw map



X Index: 175



Y Index: 159




Z Index: 187


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 925 nm³; this corresponds to an approximate mass of 836 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.189 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.189 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	5.30	-	-
Author-provided FSC curve	5.23	7.09	5.52
Unmasked-calculated*	6.68	9.82	7.08

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.68 differs from the reported value 5.3 by more than 10 %


9 Map-model fit ⓘ

This section contains information regarding the fit between EMDB map EMD-11160 and PDB model 6ZCE. Per-residue inclusion information can be found in section 3 on page 14.


9.1 Map-model overlay ⓘ



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).

9.4 Atom inclusion [i](#)



At the recommended contour level, 70% of all backbone atoms, 73% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7340	 0.2310
A	 0.9030	 0.2370
B	 0.8190	 0.2900
C	 0.7970	 0.2700
D	 0.7250	 0.2860
E	 0.6370	 0.2750
F	 0.7910	 0.2840
G	 0.7530	 0.2640
H	 0.8110	 0.2590
I	 0.8170	 0.2710
J	 0.8620	 0.2650
K	 0.7820	 0.2710
L	 0.8580	 0.2530
M	 0.7410	 0.2930
N	 0.3360	 0.0530
O	 0.8050	 0.2740
P	 0.7740	 0.2340
Q	 0.6390	 0.2110
R	 0.7750	 0.2540
S	 0.6820	 0.2760
T	 0.8030	 0.2530
U	 0.8230	 0.2390
V	 0.7270	 0.2600
W	 0.7810	 0.2850
X	 0.7370	 0.2680
Y	 0.8030	 0.2870
Z	 0.8320	 0.2640
a	 0.7520	 0.2380
b	 0.6560	 0.2730
c	 0.6530	 0.2650
d	 0.5740	 0.2590
e	 0.8650	 0.2460
f	 0.6940	 0.2870
g	 0.4720	 0.0900
h	 0.7600	 0.2200



Continued on next page...

Continued from previous page...

Chain	Atom inclusion	Q-score
i	 0.0290	 0.2280
j	 0.0030	 0.0520
k	 0.1880	 0.2140
l	 0.0000	 0.0820
m	 0.4920	 0.2080
o	 0.5090	 0.1870
p	 0.4120	 0.1670
q	 0.5340	 0.1930
r	 0.0000	 0.1130
s	 0.0300	 0.1930
t	 0.0000	 0.2500