



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 05:07 AM UTC

PDB ID : 4ZDK / pdb_00004zdk
Title : Crystal structure of the M. tuberculosis CTP synthase PyrG in complex with UTP, AMP-PCP and oxonorleucine
Authors : Bellinzoni, M.; Barilone, N.; Alzari, P.M.
Deposited on : 2015-04-17
Resolution : 3.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1085 (3.54-3.46)
Clashscore	190562	1140 (3.54-3.46)
Ramachandran outliers	187476	1113 (3.54-3.46)
Sidechain outliers	187428	1114 (3.54-3.46)
RSRZ outliers	180081	1084 (3.54-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	586	5% 85% 5% • 9%
1	B	586	6% 85% 5% • 9%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 7986 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CTP synthase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	534	Total	C	N	O	S	0	0	0
			3969	2502	701	756	10			
1	B	534	Total	C	N	O	S	0	0	0
			3875	2431	688	746	10			

- Molecule 2 is 5-OXO-L-NORLEUCINE (CCD ID: ONL) (formula: $C_6H_{11}NO_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			10	6	1	3		
2	B	1	Total	C	N	O	0	0
			10	6	1	3		

- Molecule 3 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
3	B	1	Total	C	N	O	P	0	0
			31	11	5	12	3		

- Molecule 4 is URIDINE 5'-TRIPHOSPHATE (CCD ID: UTP) (formula: $C_9H_{15}N_2O_{15}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			29	9	2	15	3		
4	B	1	Total	C	N	O	P	0	0
			29	9	2	15	3		


- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Mg 1	0	0
5	B	1	Total 1	Mg 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CTP synthase



• Molecule 1: CTP synthase



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	121.73Å 194.72Å 207.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.68 – 3.49 48.68 – 3.49	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.68-3.49) 99.8 (48.68-3.49)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 3.48Å)	Xtriage
Refinement program	BUSTER 2.10.1	Depositor
R, R_{free}	0.203 , 0.214 0.216 , 0.227	Depositor DCC
R_{free} test set	1601 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	133.4	Xtriage
Anisotropy	0.375	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 116.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7986	wwPDB-VP
Average B, all atoms (Å ²)	143.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ACP, ONL, UTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	2/4057 (0.0%)	1.21	12/5539 (0.2%)
1	B	0.76	3/3958 (0.1%)	1.22	7/5410 (0.1%)
All	All	0.76	5/8015 (0.1%)	1.22	19/10949 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	58	MET	SD-CE	-5.95	1.64	1.79
1	B	58	MET	SD-CE	-5.72	1.65	1.79
1	B	259	ILE	CA-C	5.45	1.57	1.52
1	A	259	ILE	CA-C	5.22	1.57	1.52
1	B	311	VAL	CA-C	5.20	1.59	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	241	ASP	CA-CB-CG	9.52	122.12	112.60
1	B	310	TYR	CA-C-N	6.94	132.32	121.35
1	B	310	TYR	C-N-CA	6.94	132.32	121.35
1	A	71	ASP	CA-CB-CG	6.69	119.29	112.60
1	A	312	GLU	CA-C-N	6.55	134.05	121.54
1	A	312	GLU	C-N-CA	6.55	134.05	121.54
1	A	314	SER	CA-C-N	6.51	133.98	121.54
1	A	314	SER	C-N-CA	6.51	133.98	121.54
1	A	372	ILE	N-CA-CB	-6.33	103.61	112.35
1	B	71	ASP	CA-CB-CG	6.23	118.83	112.60
1	B	277	ASN	CA-CB-CG	6.09	118.69	112.60
1	A	277	ASN	CA-CB-CG	6.06	118.66	112.60
1	B	314	SER	N-CA-C	5.65	118.68	109.24
1	A	409	LEU	CA-C-N	5.41	127.79	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	409	LEU	C-N-CA	5.41	127.79	120.38
1	A	140	ASP	CA-CB-CG	5.36	117.96	112.60
1	A	313	LEU	CA-C-N	5.13	130.30	120.97
1	A	313	LEU	C-N-CA	5.13	130.30	120.97
1	B	140	ASP	CA-CB-CG	5.01	117.61	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3969	0	3811	17	0
1	B	3875	0	3634	11	0
2	A	10	0	9	1	0
2	B	10	0	8	0	0
3	A	31	0	14	1	0
3	B	31	0	14	1	0
4	A	29	0	11	1	0
4	B	29	0	11	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
All	All	7986	0	7512	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:LEU:O	1:A:223:ARG:HD3	1.78	0.82
1:B:22:LEU:O	1:B:223:ARG:HD3	1.82	0.78
1:B:23:GLY:HA2	3:B:602:ACP:O2A	1.90	0.72
1:B:42:VAL:HG22	1:B:148:VAL:CG1	2.26	0.65
1:A:42:VAL:HG22	1:A:148:VAL:CG1	2.27	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:LEU:O	1:A:223:ARG:CD	2.50	0.60
1:B:160:ILE:H	4:B:603:UTP:H3	1.51	0.59
1:B:52:ASN:CG	1:B:58:MET:HE3	2.32	0.54
1:A:52:ASN:CG	1:A:58:MET:HE3	2.33	0.53
1:A:394:LEU:HB2	2:A:601:ONL:HG2	1.91	0.53
1:A:160:ILE:H	4:A:603:UTP:H3	1.56	0.52
1:A:173:ARG:HE	1:A:183:PHE:HE2	1.59	0.51
1:B:42:VAL:HG22	1:B:148:VAL:HG13	1.94	0.49
1:A:42:VAL:HG22	1:A:148:VAL:HG13	1.94	0.49
1:B:347:THR:HG23	1:B:350:GLY:H	1.79	0.47
1:A:23:GLY:HA3	3:A:602:ACP:H8	1.98	0.45
1:A:530:ARG:HH11	1:A:530:ARG:HD3	1.65	0.45
1:A:201:LYS:HE3	1:A:204:GLN:CD	2.43	0.44
1:A:201:LYS:HE3	1:A:204:GLN:OE1	2.18	0.43
1:B:342:SER:O	1:B:374:GLY:HA3	2.18	0.43
1:A:347:THR:HG23	1:A:350:GLY:H	1.84	0.42
1:A:184:LEU:HD11	1:A:221:ILE:HG12	2.01	0.42
1:A:342:SER:O	1:A:374:GLY:HA3	2.18	0.42
1:A:207:VAL:HG11	1:A:240:CYS:HB3	2.01	0.42
1:B:184:LEU:HD11	1:B:221:ILE:HG12	2.01	0.42
1:B:201:LYS:HE3	1:B:204:GLN:CD	2.46	0.41
1:A:363:ILE:CG2	1:A:375:LYS:HE3	2.50	0.41
1:B:207:VAL:HG11	1:B:240:CYS:HB3	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	530/586 (90%)	517 (98%)	11 (2%)	2 (0%)	30 62
1	B	530/586 (90%)	521 (98%)	8 (2%)	1 (0%)	43 74

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1060/1172 (90%)	1038 (98%)	19 (2%)	3 (0%)	36 67

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	LEU
1	B	393	CYS
1	A	393	CYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	404/480 (84%)	398 (98%)	6 (2%)	57 71
1	B	380/480 (79%)	371 (98%)	9 (2%)	43 64
All	All	784/960 (82%)	769 (98%)	15 (2%)	50 68

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	HIS
1	A	201	LYS
1	A	223	ARG
1	A	255	SER
1	A	372	ILE
1	A	471	SER
1	B	41	HIS
1	B	130	LYS
1	B	201	LYS
1	B	223	ARG
1	B	241	ASP
1	B	255	SER
1	B	312	GLU
1	B	372	ILE
1	B	471	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	234	ASN
1	A	264	HIS
1	B	205	HIS
1	B	234	ASN
1	B	264	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	UTP	A	603	-	29,30,30	0.59	0	43,47,47	0.65	0
3	ACP	A	602	5	31,33,33	0.71	2 (6%)	47,52,52	0.89	2 (4%)
3	ACP	B	602	5	31,33,33	0.96	3 (9%)	47,52,52	0.94	3 (6%)
4	UTP	B	603	-	29,30,30	0.67	0	43,47,47	0.66	0
2	ONL	B	601	1	8,9,9	1.47	2 (25%)	8,11,11	0.96	0
2	ONL	A	601	1	8,9,9	1.54	2 (25%)	8,11,11	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UTP	A	603	-	-	5/22/38/38	0/2/2/2
3	ACP	A	602	5	-	4/19/38/38	0/3/3/3
3	ACP	B	602	5	-	1/19/38/38	0/3/3/3
4	UTP	B	603	-	-	4/22/38/38	0/2/2/2
2	ONL	B	601	1	-	5/9/9/9	-
2	ONL	A	601	1	-	6/9/9/9	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	ACP	PB-O3A	3.74	1.62	1.58
2	B	601	ONL	O-C	3.22	1.31	1.22
2	A	601	ONL	O-C	3.09	1.31	1.22
2	B	601	ONL	OXT-C	-2.59	1.22	1.30
3	A	602	ACP	PG-O1G	2.55	1.55	1.50
2	A	601	ONL	OXT-C	-2.50	1.22	1.30
3	B	602	ACP	PG-O1G	2.42	1.55	1.50
3	A	602	ACP	PG-O3G	-2.26	1.49	1.55
3	B	602	ACP	PG-O3G	-2.23	1.50	1.55

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	ACP	O1B-PB-C3B	4.26	120.27	109.05
3	B	602	ACP	O1B-PB-C3B	3.76	118.94	109.05
3	B	602	ACP	O2G-PG-O1G	-2.42	106.13	112.39
3	B	602	ACP	O3G-PG-C3B	2.22	111.78	106.40
3	A	602	ACP	O3G-PG-C3B	2.06	111.39	106.40

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	ONL	N-CA-CB-CG
2	A	601	ONL	CA-CB-CG-CD
2	A	601	ONL	OD-CD-CG-CB

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	601	ONL	CE-CD-CG-CB
2	B	601	ONL	N-CA-CB-CG
2	B	601	ONL	C-CA-CB-CG
2	B	601	ONL	CA-CB-CG-CD
2	B	601	ONL	OD-CD-CG-CB
2	B	601	ONL	CE-CD-CG-CB
3	A	602	ACP	PG-C3B-PB-O1B
3	A	602	ACP	PG-C3B-PB-O3A
4	B	603	UTP	C5'-O5'-PA-O1A
4	B	603	UTP	PB-O3B-PG-O1G
4	B	603	UTP	C3'-C4'-C5'-O5'
4	A	603	UTP	PB-O3B-PG-O2G
2	A	601	ONL	OXT-C-CA-CB
2	A	601	ONL	O-C-CA-CB
3	A	602	ACP	PG-C3B-PB-O2B
3	A	602	ACP	C5'-O5'-PA-O1A
4	A	603	UTP	C5'-O5'-PA-O1A
4	B	603	UTP	C5'-O5'-PA-O3A
3	B	602	ACP	C4'-C5'-O5'-PA
4	A	603	UTP	PB-O3B-PG-O1G
4	A	603	UTP	PB-O3B-PG-O3G
4	A	603	UTP	PB-O3A-PA-O1A


There are no ring outliers.


5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	603	UTP	1	0
3	A	602	ACP	1	0
3	B	602	ACP	1	0
4	B	603	UTP	1	0
2	A	601	ONL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	534/586 (91%)	0.21	29 (5%) 31 18	94, 125, 152, 180	0
1	B	534/586 (91%)	0.32	37 (6%) 23 14	115, 156, 193, 211	0
All	All	1068/1172 (91%)	0.27	66 (6%) 26 16	94, 140, 186, 211	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	429	MET	6.4
1	B	369	ILE	5.1
1	A	369	ILE	4.7
1	B	367	PHE	4.5
1	A	367	PHE	4.1
1	B	495	PHE	4.0
1	B	531	PRO	3.9
1	B	425	VAL	3.8
1	A	410	THR	3.8
1	A	310	TYR	3.7
1	B	532	THR	3.7
1	B	429	MET	3.4
1	A	312	GLU	3.3
1	B	348	THR	3.2
1	B	398	CYS	3.2
1	B	366	GLY	3.2
1	B	443	GLY	3.2
1	A	366	GLY	3.1
1	B	411	ASN	3.1
1	B	32	GLY	3.0
1	A	314	SER	3.0
1	A	371	GLY	3.0
1	A	64	GLY	3.0
1	A	411	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	123	PRO	2.9
1	B	496	SER	2.9
1	A	240	CYS	2.9
1	A	553	LEU	2.9
1	A	311	VAL	2.8
1	A	231	ALA	2.8
1	A	55	PRO	2.8
1	A	198	LEU	2.8
1	A	468	THR	2.7
1	B	480	VAL	2.7
1	B	43	THR	2.6
1	A	357	ASP	2.6
1	B	463	GLN	2.6
1	B	236	ILE	2.6
1	B	198	LEU	2.5
1	B	507	PHE	2.5
1	B	199	LYS	2.4
1	B	7	THR	2.4
1	A	257	TYR	2.3
1	B	516	PRO	2.3
1	B	497	GLY	2.3
1	A	58	MET	2.3
1	A	372	ILE	2.3
1	A	163	GLN	2.3
1	B	18	VAL	2.2
1	B	521	THR	2.2
1	A	284	ASP	2.2
1	B	410	THR	2.2
1	A	65	GLU	2.2
1	B	343	ASP	2.2
1	A	370	ARG	2.2
1	B	508	VAL	2.2
1	A	66	VAL	2.1
1	B	313	LEU	2.1
1	B	468	THR	2.1
1	B	458	ASP	2.1
1	B	211	ARG	2.1
1	A	365	GLY	2.0
1	B	61	PHE	2.0
1	B	456	GLU	2.0
1	A	484	TYR	2.0
1	B	317	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ONL	B	601	10/10	0.81	0.13	150,157,163,166	0
4	UTP	A	603	29/29	0.83	0.11	210,213,216,217	0
4	UTP	B	603	29/29	0.84	0.09	162,164,167,168	0
2	ONL	A	601	10/10	0.86	0.13	119,122,131,132	0
3	ACP	B	602	31/31	0.89	0.10	177,183,185,187	0
3	ACP	A	602	31/31	0.90	0.09	134,136,146,147	0
5	MG	B	604	1/1	0.98	0.08	113,113,113,113	0
5	MG	A	604	1/1	1.00	0.04	94,94,94,94	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UTP A 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around UTP B 603:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ACP B 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ACP A 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.